

## ON SOME NEW FRACTIONAL HERMITE-HADAMARD TYPE INEQUALITIES FOR CONVEX AND CO-ORDINATED CONVEX FUNCTIONS

MUHAMMAD AAMIR ALI, HÜSEYİN BUDAK, AND SADIA SAKHI

ABSTRACT. In this study, some new inequalities of Hermite-Hadamard type for convex and co-ordinated convex functions via Riemann-Liouville fractional integrals are derived. It is also shown that the results obtained in this paper are the extension of some earlier ones.

### 1. Introduction

The Hermite-Hadamard inequality, which is the first fundamental result for convex mappings with a natural geometrical interpretation and many applications have drawn attention much interest in elementary mathematics. Several mathematicians have devoted their efforts to generalize, refine, counterpart and extend it for different classes of functions such as using convex mappings.

The inequalities discovered by C. Hermite and J. Hadamard for convex functions are considerable significance in the literature (see, e.g., [16, p.137], [9]). These inequalities state that if  $f : I \rightarrow \mathbb{R}$  is a convex

---

Received July 31, 2020. Revised December 21, 2020. Accepted December 22, 2020.

2010 Mathematics Subject Classification: 26B25, 26A51, 26D15.

Key words and phrases: Hermite-Hadamard inequality, fractional integrals, Convex functions, co-ordinated convex functions.

This work was partially supported by the National Natural Foundation of China (No. 11971241).

© The Kangwon-Kyungki Mathematical Society, 2020.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

function on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ , then

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}.$$

Both inequalities hold in the reversed direction if  $f$  is concave. For the further study of this area, one can consult [1]- [7], [13], [14].

In [10], the authors gave an inequality (1.1) for twice differentiable functions and they raised the succeeding problem:

do there exist real numbers  $q, Q$  such that

$$f\left(\frac{a+b}{2}\right) \leq q \leq \frac{1}{b-a} \int_a^b f(x)dx \leq Q \leq \frac{f(a)+f(b)}{2}?$$

where  $f$  is convex function.

After that, in [11], Farissi gave a favorable answer to the above-given problem and found the following values of  $q$  and  $Q$ :

$$(1.2) \quad f\left(\frac{a+b}{2}\right) \leq q(\omega) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq Q(\omega) \leq \frac{f(a)+f(b)}{2}$$

where

$$q(\omega) = \omega f\left(\frac{\omega b + (2-\omega)a}{2}\right) + (1-\omega) f\left(\frac{(1+\omega)b + (1-\omega)a}{2}\right),$$

$$Q(\omega) = \frac{1}{2}(f(\omega b + (1-\omega)a) + \omega f(a) + (1-\omega)f(b)).$$

Inspired by this work of Farissi, Chen gave these values of Hermite-Hadamard inequalities for co-ordinated convex functions as follows:

**THEOREM 1.** [8] *Let  $f : \Delta = [a, b] \times [c, d] \rightarrow \mathbb{R}$  be a co-ordinated convex function, then we have following inequality for all  $\omega, \mu \in [0, 1]$*

$$(1.3) \quad f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \leq q(\omega, \mu) \leq \int_a^b \int_c^d f(x, y) dy dx \leq Q(\omega, \mu)$$

$$\leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4},$$

where

$$\begin{aligned}
 & q(\omega, \mu) \\
 &= \omega\mu f\left(\frac{\omega b + (2 - \omega)a}{2}, \frac{\mu d + (2 - \mu)c}{2}\right) \\
 &\quad + \omega(1 - \mu) f\left(\frac{\omega b + (2 - \omega)a}{2}, \frac{(1 + \mu)d + (1 - \mu)c}{2}\right) \\
 &\quad + (1 - \omega)\mu f\left(\frac{(1 + \omega)b + (1 - \omega)a}{2}, \frac{\mu d + (2 - \mu)c}{2}\right) \\
 &\quad + (1 - \omega)(1 - \mu) f\left(\frac{(1 + \omega)b + (1 - \omega)a}{2}, \frac{(1 + \mu)d + (1 - \mu)c}{2}\right)
 \end{aligned}$$

and

$$\begin{aligned}
 & Q(\omega, \mu) \\
 &= \frac{\omega\mu}{4} f(a, c) + \frac{\omega(1 - \mu)}{4} f(a, d) + \frac{(1 - \omega)\mu}{4} f(b, c) + \frac{(1 - \omega)(1 - \mu)}{4} f(b, d) \\
 &\quad + \frac{f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c)}{4} + \frac{\omega}{4} f(a, \mu d + (1 - \mu)c) \\
 &\quad + \frac{1 - \omega}{4} f(b, \mu d + (1 - \mu)c) + \frac{\mu}{4} f(\omega b + (1 - \omega)a, c) \\
 &\quad + \frac{1 - \mu}{4} f(\omega b + (1 - \omega)a, d).
 \end{aligned}$$

The main objective of this paper is to give the fractional variant of inequalities (1.2) and (1.3).

## 2. Preliminaries

In this section, we review the definitions of Riemann Liouville fractional integrals for single and two variables functions.

DEFINITION 1. [12] Let  $f \in L_1[a, b]$ . The Riemann-Liouville integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha$  of order  $\alpha > 0$  with  $a \geq 0$  are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x - t)^{\alpha-1} f(t) dt, \quad a < x$$

and

$$J_{b-}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively. Here  $\Gamma(\alpha)$  is Gamma function and  $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$ .

In [18], Sarikaya et al. gave the following Hermite-Hadamard inequalities concerned with the last fractional integrals.

**THEOREM 2.** [18] *Let  $f : [a, b] \rightarrow R$  be a positive function with  $0 \leq a < b$  and  $f \in L_1[a, b]$ . If  $f$  is a convex function on  $[a, b]$ , then the following inequalities for fractional integrals hold:*

$$(2.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}} [J_{a+}^{\alpha} f(b) + J_{b-}^{\alpha} f(a)] \leq \frac{f(a) + f(b)}{2},$$

where  $\alpha > 0$ .

**EXAMPLE 1.** A function  $f(x) = x^2$  is a convex function. The above inequality (2.1) holds for the given  $f(x)$ .

**SOLUTION 1.** For  $\alpha = \frac{1}{2}$ ,  $a = 1$ , and  $b = 2$ , we have

$$\begin{aligned} \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}} [J_{a+}^{\alpha} f(b) + J_{b-}^{\alpha} f(a)] &= 2.36, \\ f\left(\frac{a+b}{2}\right) &= 2.25, \end{aligned}$$

and

$$\frac{f(a) + f(b)}{2} = 2.5.$$

Thus, the inequality (2.1) is true.

In [17], Sarikaya offered the following Riemann-Liouville fractional integrals and associated inequalities of Hermite-Hadamard type:

**DEFINITION 2.** [17] Let  $f \in L_1([a, b] \times [c, d])$ . Then Riemann-Liouville integrals  $J_{a+,c+}^{\alpha,\beta}$ ,  $J_{a+,d-}^{\alpha,\beta}$ ,  $J_{b-,c+}^{\alpha,\beta}$  and  $J_{b-,d-}^{\alpha,\beta}$  of order  $\alpha, \beta > 0$  with  $a, c \geq 0$

are defined by

$$J_{a+,c+}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y (x-t)^{\alpha-1} (y-s)^{\beta-1} f(t,s) dt ds,$$

$$x > a, y > c,$$

$$J_{a+,d-}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_y^d (x-t)^{\alpha-1} (s-y)^{\beta-1} f(t,s) dt ds,$$

$$x > a, y < d,$$

$$J_{b-,c+}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_c^y (t-x)^{\alpha-1} (y-s)^{\beta-1} f(t,s) dt ds,$$

$$x < b, y > c$$

and

$$J_{b-,d-}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_y^d (t-x)^{\alpha-1} (s-y)^{\beta-1} f(t,s) dt ds,$$

$$x < b, y < d,$$

respectively. Here  $\Gamma$  is a gamma function,

$$J_{a+,c+}^{0,0} f(x,y) = J_{a+,d-}^{0,0} f(x,y) = J_{b-,c+}^{0,0} f(x,y) = J_{b-,d-}^{0,0} f(x,y) = f(x,y).$$

**THEOREM 3.** [17] *Let  $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  be a co-ordinated convex function on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $0 \leq a < b, 0 \leq c < d$  and  $f \in L_1(\Delta)$ . Then we have following inequalities for double fractional integrals:*

$$\begin{aligned} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) &\leq \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \left[ J_{a+,c+}^{\alpha,\beta} f(b,d) + J_{a+,d-}^{\alpha,\beta} f(b,c) \right. \\ (2.2) \quad &\quad \left. + J_{b-,c+}^{\alpha,\beta} f(a,d) + J_{b-,d-}^{\alpha,\beta} f(a,c) \right] \\ &\leq \frac{f(a,c) + f(a,d) + f(b,c) + f(b,d)}{4}. \end{aligned}$$

### 3. Key Results

For brevity, we use the following notations in upcoming new results:

$$\Delta_1 = [a, \omega b + (1 - \omega) a], \quad \Delta_2 = [\omega b + (1 - \omega) a, b].$$

and

$$\Delta = [a, b] \times [c, d] = \Delta_3 \cup \Delta_4 \cup \Delta_5 \cup \Delta_6$$

where

$$\begin{aligned}\Delta_3 &= [a, \omega b + (1 - \omega) a] \times [c, \mu d + (1 - \mu) c], \\ \Delta_4 &= [a, \omega b + (1 - \omega) a] \times [\mu d + (1 - \mu) c, d], \\ \Delta_5 &= [\omega b + (1 - \omega) a, b] \times [c, \mu d + (1 - \mu) c], \\ \Delta_6 &= [\omega b + (1 - \omega) a, b] \times [\mu d + (1 - \mu) c, d].\end{aligned}$$

**THEOREM 4.** *Suppose that  $f : I \rightarrow \mathbb{R}$  is a convex function, then following inequalities hold for all  $\omega \in [0, 1]$ ,*

$$f\left(\frac{a+b}{2}\right) \leq q(\omega) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} I^\alpha(f) \leq Q(\omega) \leq \frac{f(a)+f(b)}{2},$$

where

$$\begin{aligned}I^\alpha(f) &= \frac{1}{\omega^{\alpha-1}} [J_{a+}^\alpha f(\omega b + (1 - \omega) a) + J_{(\omega b + (1 - \omega) a)-}^\alpha f(a)] \\ &\quad + \frac{1}{(1 - \omega)^{\alpha-1}} [J_{b-}^\alpha f(\omega b + (1 - \omega) a) + J_{(\omega b + (1 - \omega) a)+}^\alpha f(b)], \\ q(\omega) &= \omega \left( f\left(\frac{\omega b + (2 - \omega) a}{2}\right) + (1 - \omega) f\left(\frac{(1 + \omega) b + (1 - \omega) a}{2}\right) \right), \\ Q(\omega) &= \frac{1}{2} (f(\omega b + (1 - \omega) a) + \omega f(a) + (1 - \omega) f(b))\end{aligned}$$

and  $\alpha > 0$ .

*Proof.* From inequalities in (2.1) over the  $\Delta_1$ , we have

$$\begin{aligned}(3.1) \quad & f\left(\frac{\omega b + (2 - \omega) a}{2}\right) \\ & \leq \frac{\Gamma(\alpha+1)}{2\omega^\alpha(b-a)^\alpha} [J_{a+}^\alpha f(\omega b + (1 - \omega) a) + J_{(\omega b + (1 - \omega) a)-}^\alpha f(a)] \\ & \leq \frac{f(a) + f(\omega b + (1 - \omega) a)}{2}.\end{aligned}$$

Again from inequalities in (2.1) over  $\Delta_2$ , we find that

$$\begin{aligned}(3.2) \quad & f\left(\frac{(1 + \omega) b + (1 - \omega) a}{2}\right) \\ & \leq \frac{\Gamma(\alpha+1)}{2(1 - \omega)^\alpha(b-a)^\alpha} [J_{(\omega b + (1 - \omega) a)+}^\alpha f(b) + J_{b-}^\alpha f(\omega b + (1 - \omega) a)] \\ & \leq \frac{f(b) + f(\omega b + (1 - \omega) a)}{2}.\end{aligned}$$

Multiplying (3.1), (3.2) by  $\omega$  and  $(1 - \omega)$ , respectively. After that, adding the resultant inequalities, we obtain that

$$(3.3) \quad q(\omega) \leq \frac{\Gamma(\alpha + 1)}{2(b - a)^\alpha} I^\alpha(f) \leq Q(\omega).$$

Since  $f$  is convex function, so we have

$$(3.4) \quad \begin{aligned} & f\left(\frac{a + b}{2}\right) \\ &= f\left(\omega\left(\frac{\omega b + (1 - \omega)a + a}{2}\right) + (1 - \omega)\left(\frac{\omega b + (1 - \omega)a + b}{2}\right)\right) \\ &\leq \omega f\left(\frac{\omega b + (1 - \omega)a + a}{2}\right) + (1 - \omega) f\left(\frac{\omega b + (1 - \omega)a + b}{2}\right) \\ &\leq \frac{1}{2}(f(\omega b + (1 - \omega)a) + \omega f(a) + (1 - \omega) f(b)) \\ &\leq \frac{f(a) + f(b)}{2}. \end{aligned}$$

From (3.3) and (3.4), we conclude the desired inequality. □

REMARK 1. Under the hypothesis of Theorem 4 with  $\alpha = 1$ , we have [11, Theorem 1.1].

COROLLARY 1. Under the same conditions and notations stated in Theorem 4, we have the following new inequalities

$$f\left(\frac{a + b}{2}\right) \leq \sup_{\omega \in [0,1]} q(\omega) \leq \frac{\Gamma(\alpha + 1)}{2(b - a)^\alpha} I^\alpha(f) \leq \inf_{\omega \in [0,1]} Q(\omega) \leq \frac{f(a) + f(b)}{2}.$$

THEOREM 5. Let  $f : \Delta \rightarrow \mathbb{R}$  be a co-ordinated convex function and  $f \in L(\Delta)$ , then the following inequalities satisfy for all  $\omega, \mu \in [0, 1]$  :

$$\begin{aligned} & f\left(\frac{a + b}{2}, \frac{c + d}{2}\right) \\ &\leq q(\omega, \mu) \leq \frac{\Gamma(\alpha + 1)\Gamma(\beta + 1)}{4(b - a)^\alpha(d - c)^\beta} I^{\alpha,\beta}(f) \leq Q(\omega, \mu) \\ &\leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \end{aligned}$$

where

$$\begin{aligned}
I^{\alpha, \beta}(f) &= \frac{1}{\omega^{\alpha-1} \mu^{\beta-1}} \left\{ J_{a^+, c^+}^{\alpha, \beta} f(\omega b + (1-\omega)a, \mu d + (1-\mu)c) \right. \\
&+ J_{a^+, (\mu d + (1-\mu)c)^-}^{\alpha, \beta} f(\omega b + (1-\omega)a, c) + J_{(\omega b + (1-\omega)a)^-, c^+}^{\alpha, \beta} f(a, \mu d + (1-\mu)c) \\
&+ \left. J_{(\omega b + (1-\omega)a)^-, (\mu d + (1-\mu)c)^-}^{\alpha, \beta} f(a, c) \right\} \\
&+ \frac{1}{\omega^{\alpha-1} (1-\mu)^{\beta-1}} \left\{ J_{a^+, (\mu d + (1-\mu)c)^+}^{\alpha, \beta} f(\omega b + (1-\omega)a, d) \right. \\
&+ J_{a^+, d^-}^{\alpha, \beta} f(\omega b + (1-\omega)a, \mu d + (1-\mu)c) + J_{(\omega b + (-\omega)a)^-, (\mu d + (1-\mu)c)^+}^{\alpha, \beta} f(a, d) \\
&+ \left. J_{(\omega b + (-\omega)a)^-, d^-}^{\alpha, \beta} f(a, \mu d + (1-\mu)c) \right\} \\
&+ \frac{1}{(1-\omega)^{\alpha-1} \mu^{\beta-1}} \left\{ J_{(\omega b + (1-\omega)a)^+, c^+}^{\alpha, \beta} f(b, \mu d + (1-\mu)c) \right. \\
&+ J_{(\omega b + (1-\omega)a)^+, (\mu d + (1-\mu)c)^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(\omega b + (1-\omega)a, \mu d + (1-\mu)c) \\
&+ \left. J_{b^-, (\mu d + (1-\mu)c)^-}^{\alpha, \beta} f(\omega b + (1-\omega)a, c) \right\} \\
&+ \frac{1}{(1-\omega)^{\alpha-1} (1-\mu)^{\beta-1}} \left\{ J_{(\omega b + (1-\omega)a)^+, (\mu d + (1-\mu)c)^+}^{\alpha, \beta} f(b, d) \right. \\
&+ J_{(\omega b + (1-\omega)a)^+, d^-}^{\alpha, \beta} f(b, \mu d + (1-\mu)c) + J_{b^-, (\mu d + (1-\mu)c)^+}^{\alpha, \beta} f(\omega b + (1-\omega)a, d) \\
&+ \left. J_{b^-, d^-}^{\alpha, \beta} f(\omega b + (1-\omega)a, \mu d + (1-\mu)c) \right\},
\end{aligned}$$

and  $\alpha, \beta > 0$ .

*Proof.* From inequalities given in(2.2) for  $\Delta_3, \Delta_4, \Delta_5, \Delta_6$  with  $\omega \neq 0, 1$  and  $\mu \neq 0, 1$ , we get that

$$(3.5) \quad f\left(\frac{\omega b + (2-\omega)a}{2}, \frac{\mu d + (2-\mu)c}{2}\right)$$



$$\begin{aligned}
 &\leq \frac{\Gamma(\alpha + 1)\Gamma(\beta + 1)}{4\omega^\alpha\mu^\beta(b-a)^\alpha(d-c)^\beta} \\
 &\times \left[ J_{a+,c+}^{\alpha,\beta} f(\omega b + (1-\omega)a, \mu d + (1-\mu)c) \right. \\
 &+ J_{a+,( \mu d + (1-\mu)c)-}^{\alpha,\beta} f(\omega b + (1-\omega)a, c) \\
 &+ J_{(\omega b + (1-\omega)a)-,c+}^{\alpha,\beta} f(a, \mu d + (1-\mu)c) \\
 &\left. + J_{(\omega b + (1-\omega)a)-,( \mu d + (1-\mu)c)-}^{\alpha,\beta} f(a, c) \right] \\
 &\leq \frac{1}{4} [f(a, c) + f(a, \mu d + (1-\mu)c) + f(\omega b + (1-\omega)a, c) \\
 &+ f(\omega b + (1-\omega)a, \mu d + (1-\mu)c)], \\
 (3.6) \quad &f\left(\frac{\omega b + (2-\omega)a}{2}, \frac{(1+\mu)d + (1-\mu)c}{2}\right) \\
 &\leq \frac{\Gamma(\alpha + 1)\Gamma(\beta + 1)}{4\omega^\alpha(1-\mu)^\beta(b-a)^\alpha(d-c)^\beta} \\
 &\times \left[ J_{a+,( \mu d + (1-\mu)c)+}^{\alpha,\beta} f(\omega b + (1-\omega)a, d) \right. \\
 &+ J_{a+,d-}^{\alpha,\beta} f(\omega b + (1-\omega)a, \mu d + (1-\mu)c) \\
 &+ J_{(\omega b + (-\omega)a)-,( \mu d + (1-\mu)c)+}^{\alpha,\beta} f(a, d) \\
 &\left. + J_{(\omega b + (-\omega)a)-,d-}^{\alpha,\beta} f(a, \mu d + (1-\mu)c) \right] \\
 &\leq \frac{1}{4} [f(a, \mu d + (1-\mu)c) + f(a, d) \\
 &+ f(\omega b + (-\omega)a, \mu d + (1-\mu)c) + f(\omega b + (-\omega)a, d)],
 \end{aligned}$$

$$\begin{aligned}
(3.7) \quad & f\left(\frac{(1+\omega)b+(1-\omega)a}{2}, \frac{\mu d+(2-\mu)c}{2}\right) \\
& \leq \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(1-\omega)^\alpha \mu^\beta (b-a)^\alpha (d-c)^\beta} \\
& \quad \times \left[ J_{(\omega b+(1-\omega)a)_+, c+}^{\alpha, \beta} f(b, \mu d+(1-\mu)c) \right. \\
& \quad + J_{(\omega b+(1-\omega)a)_+, (\mu d+(1-\mu)c)-}^{\alpha, \beta} f(b, c) \\
& \quad + J_{b^-, c+}^{\alpha, \beta} f(\omega b+(1-\omega)a, \mu d+(1-\mu)c) \\
& \quad \left. + J_{b^-, (\mu d+(1-\mu)c)-}^{\alpha, \beta} f(\omega b+(1-\omega)a, c) \right] \\
& \leq \frac{1}{4} [f(\omega b+(1-\omega)a, c) + f(\omega b+(1-\omega)a, \mu d+(1-\mu)c) \\
& \quad + f(b, c) + f(b, \mu d+(1-\mu)c)],
\end{aligned}$$

$$\begin{aligned}
(3.8) \quad & f\left(\frac{(1+\omega)b+(1-\omega)a}{2}, \frac{(1+\mu)d+(1-\mu)c}{2}\right) \\
& \leq \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(1-\omega)^\alpha (1-\mu)^\beta (b-a)^\alpha (d-c)^\beta} \\
& \quad \times \left[ J_{(\omega b+(1-\omega)a)_+, (\mu d+(1-\mu)c)+}^{\alpha, \beta} f(b, d) \right. \\
& \quad + J_{(\omega b+(1-\omega)a)_+, d-}^{\alpha, \beta} f(b, \mu d+(1-\mu)c) \\
& \quad + J_{b^-, (\mu d+(1-\mu)c)+}^{\alpha, \beta} f(\omega b+(1-\omega)a, d) \\
& \quad \left. + J_{b^-, d-}^{\alpha, \beta} f(\omega b+(1-\omega)a, \mu d+(1-\mu)c) \right]
\end{aligned}$$

$$\leq \frac{1}{4} [f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c) + f(\omega b + (1 - \omega)a, d) + f(b, \mu d + (1 - \mu)c) + f(b, d)].$$

Multiplying (3.5), (3.6), (3.7) and (3.8) by  $\omega\mu$ ,  $\omega(1 - \mu)$ ,  $(1 - \omega)\mu$  and  $(1 - \omega)(1 - \mu)$ , respectively. After that, adding the resultant inequalities, we found that

$$\begin{aligned} (3.9) \quad & \omega\mu f\left(\frac{\omega b + (2 - \omega)a}{2}, \frac{\mu d + (2 - \mu)c}{2}\right) \\ & + \omega(1 - \mu) f\left(\frac{\omega b + (2 - \omega)a}{2}, \frac{(1 + \mu)d + (1 - \mu)c}{2}\right) \\ & + (1 - \omega)\mu f\left(\frac{(1 + \omega)b + (1 - \omega)a}{2}, \frac{\mu d + (2 - \mu)c}{2}\right) \\ & + (1 - \omega)(1 - \mu) f\left(\frac{(1 + \omega)b + (1 - \omega)a}{2}, \frac{(1 + \mu)d + (1 - \mu)c}{2}\right) \\ = & q(\omega, \mu) \\ \leq & \frac{\Gamma(\alpha + 1)\Gamma(\beta + 1)}{4(b - a)^\alpha(d - c)^\beta} \\ & \times \left[ \frac{1}{\omega^{\alpha-1}\mu^{\beta-1}} \left\{ J_{a^+,c^+}^{\alpha,\beta} f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c) \right. \right. \\ & + J_{a^+, (\mu d + (1 - \mu)c)^-}^{\alpha,\beta} f(\omega b + (1 - \omega)a, c) \\ & + J_{(\omega b + (1 - \omega)a)^-, c^+}^{\alpha,\beta} f(a, \mu d + (1 - \mu)c) \\ & \left. \left. + J_{(\omega b + (1 - \omega)a)^-, (\mu d + (1 - \mu)c)^-}^{\alpha,\beta} f(a, c) \right\} \right] \\ & + \frac{1}{\omega^{\alpha-1}(1 - \mu)^{\beta-1}} \left\{ J_{a^+, (\mu d + (1 - \mu)c)^+}^{\alpha,\beta} f(\omega b + (1 - \omega)a, d) \right. \\ & + J_{a^+, d^-}^{\alpha,\beta} f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c) \\ & \left. + J_{(\omega b + (-\omega)a)^-, (\mu d + (1 - \mu)c)^+}^{\alpha,\beta} f(a, d) \right\} \end{aligned}$$

$$\begin{aligned}
& + J_{(\omega b + (-\omega)a)_-, d-}^{\alpha, \beta} f(a, \mu d + (1 - \mu)c) \Big\} \\
& + \frac{1}{(1 - \omega)^{\alpha-1} \mu^{\beta-1}} \left\{ J_{(\omega b + (1-\omega)a)_+, c+}^{\alpha, \beta} f(b, \mu d + (1 - \mu)c) \right. \\
& + J_{(\omega b + (1-\omega)a)_+, (\mu d + (1-\mu)c)-}^{\alpha, \beta} f(b, c) \\
& + J_{b-, c+}^{\alpha, \beta} f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c) \\
& \left. + J_{b-, (\mu d + (1-\mu)c)-}^{\alpha, \beta} f(\omega b + (1 - \omega)a, c) \right\} \\
& + \frac{1}{(1 - \omega)^{\alpha-1} (1 - \mu)^{\beta-1}} \left\{ J_{(\omega b + (1-\omega)a)_+, (\mu d + (1-\mu)c)+}^{\alpha, \beta} f(b, d) \right. \\
& + J_{(\omega b + (1-\omega)a)_+, d-}^{\alpha, \beta} f(b, \mu d + (1 - \mu)c) \\
& + J_{b-, (\mu d + (1-\mu)c)+}^{\alpha, \beta} f(\omega b + (1 - \omega)a, d) \\
& \left. + J_{b-, d-}^{\alpha, \beta} f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c) \right\} \\
\leq & \frac{\omega \mu}{4} [f(a, c) + f(a, \mu d + (1 - \mu)c) + f(\omega b + (1 - \omega)a, c) \\
& + f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c)] \\
& + \frac{\omega(1 - \mu)}{4} [f(a, \mu d + (1 - \mu)c) + f(a, d) \\
& + f(\omega b + (-\omega)a, \mu d + (1 - \mu)c) + f(\omega b + (-\omega)a, d)] \\
& + \frac{(1 - \omega)\mu}{4} [f(\omega b + (1 - \omega)a, c) \\
& f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c)
\end{aligned}$$

$$\begin{aligned}
 & + f(b, c) + f(b, \mu d + (1 - \mu)c)] \\
 & + \frac{(1 - \omega)(1 - \mu)}{4} [f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c) \\
 & + f(\omega b + (1 - \omega)a, d) \\
 & + f(b, \mu d + (1 - \mu)c) + f(b, d)] \\
 = & \frac{\omega\mu}{4} f(a, c) + \frac{\omega(1 - \mu)}{4} f(a, d) \\
 & + \frac{(1 - \omega)\mu}{4} f(b, c) + \frac{(1 - \omega)(1 - \mu)}{4} f(b, d) \\
 & + \frac{f(\omega b + (1 - \omega)a, \mu d + (1 - \mu)c)}{4} \\
 & + \frac{\omega}{4} f(a, \mu d + (1 - \mu)c) \\
 & + \frac{1 - \omega}{4} f(b, \mu d + (1 - \mu)c) + \frac{\mu}{4} f(\omega b + (1 - \omega)a, c) \\
 & + \frac{1 - \mu}{4} f(\omega b + (1 - \omega)a, d) \\
 = & Q(\omega, \mu).
 \end{aligned}$$

Since  $f$  is co-ordinated convex function, so we obtain that

$$\begin{aligned}
 (3.10) \quad & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 = & f\left(\omega \frac{\omega b + (2 - \omega)a}{2} + (1 - \omega) \frac{(1 + \omega)b + (1 - \omega)a}{2}, \right. \\
 & \left. \mu \frac{\mu d + (2 - \mu)c}{2} + (1 - \mu) \frac{(1 + \mu)d + (1 - \mu)c}{2}\right) \\
 \leq & \omega\mu f\left(\frac{\omega b + (2 - \omega)a}{2}, \frac{\mu d + (2 - \mu)c}{2}\right) \\
 & + \omega(1 - \mu) f\left(\frac{\omega b + (2 - \omega)a}{2}, \frac{(1 + \mu)d + (1 - \mu)c}{2}\right) \\
 & + (1 - \omega)\mu f\left(\frac{(1 + \omega)b + (1 - \omega)a}{2}, \frac{\mu d + (2 - \mu)c}{2}\right)
 \end{aligned}$$

$$\begin{aligned}
& + (1 - \omega)(1 - \mu) f \left( \frac{(1 + \omega)b + (1 - \omega)a}{2}, \frac{(1 + \mu)d + (1 - \mu)c}{2} \right) \\
& = q(\omega, \mu).
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
(3.11) \quad & Q(\omega, \mu) \\
& \leq \frac{\omega\mu}{4} f(a, c) + \frac{\omega(1 - \mu)}{4} f(a, d) \\
& \quad + \frac{(1 - \omega)\mu}{4} f(b, c) + \frac{(1 - \omega)(1 - \mu)}{4} f(b, d) \\
& \quad + \frac{(1 - \omega)(1 - \mu)}{4} f(a, c) + \frac{(1 - \omega)\mu}{4} f(a, d) \\
& \quad + \frac{\omega(1 - \mu)}{4} f(b, c) + \frac{\omega\mu}{4} f(b, d) \\
& \quad + \frac{\omega\mu}{4} f(a, d) + \frac{\omega(1 - \mu)}{4} f(a, c) \\
& \quad + \frac{(1 - \omega)\mu}{4} f(b, d) + \frac{(1 - \omega)(1 - \mu)}{4} f(b, c) \\
& \quad + \frac{\mu\omega}{4} f(b, c) + \frac{(1 - \omega)\mu}{4} f(a, c) \\
& \quad + \frac{\omega(1 - \mu)}{4} f(b, d) + \frac{(1 - \omega)(1 - \mu)}{4} f(a, d) \\
& = \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}.
\end{aligned}$$

By (3.9), (3.10) and (3.11), we conclude the required inequality.  $\square$

**REMARK 2.** Under the same assumptions given stated in Theorem 5 with  $\alpha = \beta = 1$ , then we have [8, Theorem 2.1].

**REMARK 3.** Under the same assumptions stated in Theorem 5 with  $\omega = \mu = \frac{1}{2}$  and  $\alpha = \beta = 1$ , then we have result of [15, Theorem 2.6].

COROLLARY 2. *Under the same conditions and notations stated in Theorem 5, we have following inequalities*

$$\begin{aligned}
 f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) &\leq \sup_{\omega, \mu \in [0,1]} q(\omega, \mu) \leq I^{\alpha, \beta}(f) \\
 &\leq \inf_{\omega, \mu \in [0,1]} Q(\omega, \mu) \\
 &\leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}.
 \end{aligned}$$

#### 4. Conclusion

In this investigation, a new fractional version of Hermite-Hadamard type inequalities for convex and co-ordinated convex functions is derived. Some existing and new inequalities are also obtained in the special cases of the main results. The authors hope that this work may stimulate further research in different areas of pure and applied sciences.

#### Acknowledgment

The first author is thankful to the Chinese Scholarship Council for offering a full scholarship in his Ph.D. studies at Nanjing Normal University, Nanjing, China. The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

#### References

- [1] M. A. Ali, H. Budak, M. Abbas, M. Z. Sarikaya, and A. Kashuri, *Hermite-Hadamard Type Inequalities for h-convex Functions via Generalized Fractional Integrals*. JOURNAL OF MATHEMATICAL EXTENSION, **14** (1) (2019).
- [2] M. A. Ali, H. Budak, Z. Zhang, and H. Yildirim, *Some new Simpson's type inequalities for co-ordinated convex functions in quantum calculus*, Mathematical Methods in the Applied Sciences, <https://doi.org/10.1002/mma.7048>.
- [3] M. A. Ali, H. Budak, M. Abbas, and Y.-M. Chu, *Quantum Hermite-Hadamard type inequalities for functions whose second  $q^b$ -derivatives absolute value are convex*, Advances in Difference Equation, In press, 2020.
- [4] H. Budak, *Some trapezoid and midpoint type inequalities for newly defined quantum integrals*, Proyecciones Journal of Mathematics, in press.

- [5] H. Budak, S. Erden, and M. A. Ali, *Simpson and Newton type inequalities for convex functions via newly defined quantum integrals*, Mathematical Methods in the Applied Sciences (2020).
- [6] H. Budak, M. A. Ali, and M. Tarhanaci, *Some New Quantum Hermite–Hadamard-Like Inequalities for Coordinated Convex Functions*, Journal of Optimization Theory and Applications (2020): 1–12.
- [7] H. Budak, M. A. Ali, and T. Tunç, *Quantum Ostrowski type integral inequalities for functions of two variables*, Mathematical Methods in the Applied Sciences, In press, 2020.
- [8] F. Chen, *A note on the Hermite-Hadamard inequality for convex functions on the co-ordinates*, Journal of Mathematical Inequalities **8** (4) (2014), 915–923.
- [9] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000.
- [10] A. E. Farissi, Z. Latreuch and B. Belaidi, *Hadamard-Type Inequalities for Twice Differentiable Functions*, RGMIA Research Report collection, 12, 1 (2009), Art. 6.
- [11] A. E. Farissi, *Simple proof and refinements of Hermite-Hadamard inequality*, Journal of Mathematical Inequalities. **4** (3) (2010), 365–369.
- [12] R. Gorenflo, F. Mainardi, *Fractional Calculus, Integral and Differential Equations of Fractional Order*, Springer Verlag, Wien, 1997, 223–276.
- [13] A. Kashuri, M. A. Ali, M. Abbas, and H. Budak, *New inequalities for generalized  $m$ -convex functions via generalized fractional integral operators and their applications*, International Journal of Nonlinear Analysis and Applications **10** (2) (2019), 275–299.
- [14] A. Kashuri, M. A. Ali, M. Abbas, H. Budak, and M. Z. Sarikaya, *Fractional integral inequalities for generalized convexity*, Tbilisi Mathematical Journal **13** (3) (2020), 63–83.
- [15] M. E. Özdemir, Ş. Yildiz and A. O. Akdemir, *On some new Hadamard-type inequalities for co-ordinated quasi-convex functions*, Hacettepe Journal of Mathematics and Statistics **41** (5) (2012), 697–707.
- [16] J.E. Pečarić, F. Proschan and Y.L. Tong, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, Boston, 1992.
- [17] M. Z. Sarikaya, *On the Hermite-Hadamard-type inequalities for co-ordinated convex functions via fractional integrals*, Integral Transforms and Special Functions **25** (2) (2014), 134–147.
- [18] M. Z. Sarikaya, E. Set, H. Yildiz and N. Başak, *Hermite-Hadamard’s inequalities for fractional integrals and related inequalities*, Mathematical and Computer Modelling **57** (2013), 2403–2407.



**Muhammad Aamir Ali**

Jiangsu Key Laboratory for NSLSCS  
School of Mathematical Sciences,  
Nanjing Normal University, Nanjing, China.  
*E-mail:* mahr.muhammad.aamir@gmail.com

**Hüseyin Budak**

Department of Mathematics  
Faculty of Science and Arts, Düzce  
University, Düzce-TURKEY  
*E-mail:* hsyn.budak@gmail.com

**Sadia Sakhi**

Department of Mathematics  
Institute of Southern Punjab, Multan, Pakistan  
*E-mail:* sadiasakhi111@gmail.com