

A FIXED POINT THEOREM IN HILBERT C^* -MODULES

HASAN RANJBAR AND ASADOLLAH NIKNAM

ABSTRACT. Fixed point theory has many useful applications in applied sciences. The object of this paper is to obtain fixed point for continuous self mappings in Hilbert C^* -module with rational conditions.

1. Introduction

Let \mathcal{A} be a C^* -algebra. A (right) inner-product \mathcal{A} -module is a linear space E , which is a right \mathcal{A} -module and $\lambda(xa) = (\lambda x)a = x(\lambda a)$ for all $x \in E$, $a \in \mathcal{A}$, $\lambda \in \mathbb{C}$, together with an inner product $\langle \cdot, \cdot \rangle : E \times E \rightarrow \mathcal{A}$ satisfying the following conditions:

- (i) $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0$ if and only if $x = 0$,
- (ii) $\langle x, \lambda y + z \rangle = \lambda \langle x, y \rangle + \langle x, z \rangle$,
- (iii) $\langle x, ya \rangle = \langle x, y \rangle a$,
- (iv) $\langle x, y \rangle^* = \langle y, x \rangle$,

for all $x, y, z \in E$, $a \in \mathcal{A}$ and $\lambda \in \mathbb{C}$. A Hilbert \mathcal{A} -module (Hilbert C^* -module) is an inner product \mathcal{A} -module E which is complete under the norm $\|x\| = \|\langle x, x \rangle\|^{1/2}$. Let E be a Hilbert \mathcal{A} -module. A map $T : E \rightarrow E$ is called adjointable if there is a map $T^* : E \rightarrow E$ such that $\langle Tx, y \rangle = \langle x, T^*y \rangle$ for all $x, y \in E$. It is easy to see that T must be \mathcal{A} -linear (i.e., $T(xa) = T(x)a$ for all $x \in E$ and $a \in \mathcal{A}$) and bounded [9, P 8]. The set of all adjointable maps is denoted by $\mathcal{L}(E)$, which is a C^* -algebra. For every pair of vectors $x, y \in E$, we use $\theta_{x,y}$ to denote the rank 1 linear operator on E , defined by $\theta_{x,y}(z) = x\langle y, z \rangle$ for any $z \in E$. The closed linear subspace of $\mathcal{L}(E)$ spanned by $\{\theta_{x,y} : x, y \in E\}$ is denoted by $\mathcal{K}(E)$. In fact, $\mathcal{K}(E)$ is a closed ideal of $\mathcal{L}(E)$ and is called the algebra of “compact” operators.

In mathematics, a fixed point (sometimes shortened to fixpoint, also known as an invariant point) of a function is an element of the function’s domain that is mapped to itself by the function. That is to say, c is a fixed point of the function f if $f(c) = c$. A topological space X is said to have the fixed point property (briefly FPP) if for any continuous function $f : X \rightarrow X$ there exists $x \in X$ such that $f(x) = x$. Banach space is said to have the fixed point property (briefly, FPP) if every nonexpansive a self mapping defined on a nonempty closed convex bounded subset has a fixed point. In 1965, Browder presented a fundamental fixed point theorem that states every Hilbert space has FPP [1]. In the same year, Browder [2] and Göhde [4] independently showed even more that every uniformly convex Banach space has FPP and Kirk established

Received July 10, 2021. Revised April 24, 2022. Accepted April 27, 2022.

2010 Mathematics Subject Classification: 47H10, 46L08.

Key words and phrases: Fixed point, Hilbert C^* -module, continuous self mapping.

© The Kangwon-Kyungki Mathematical Society, 2022.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

this property for a larger class of reflexive Banach spaces with normal structure [7]. As pointed out, the first known Banach spaces with the FPP were the Hilbert spaces [1]. On the other hand, Hilbert C-modules as a powerful tool in operator and operator algebra theory generalize the notion of a Hilbert space. Thus, a natural question raises that which classes of Hilbert C*-modules have FPP. In this paper, we aim to investigate FPP for Hilbert modules.

2. Preliminaries

The main result of this paper is that every Hilbert C*-module over C*-algebra has FPP. First we see some fundamental results:

THEOREM 2.1. [5] *If f is self mapping of a complete metric space X into itself satisfying:*

$$d(x, y) \leq \alpha(d(Tx, x) + d(Ty, y)),$$

for all $x, y \in X$ and $\alpha \in [0, 1/2)$, then f has a unique fixed point in X .

THEOREM 2.2. [3] *If $d(x, y) \leq \alpha(d(Tx, x) + d(Ty, y)) + \beta d(x, y)$, for all $x, y \in X$ and $\alpha, \beta \in [0, 1/2)$, then f has a unique fixed point in X .*

THEOREM 2.3. [8] *If T is a self mapping on a closed subset S of a Hilbert Space H with Kannan type condition*

$$\|Tx - Ty\|^2 \leq \alpha(\|Tx - x\|^2 + \|Ty - y\|^2),$$

for all $x, y \in S$ and $\alpha \in [0, 1/2)$, then T has a unique fixed point in S .

Sharma et.al [11] have proved the common fixed point theorems in Hilbert space with following condition

$$\|Tx - Ty\| \leq \alpha \left(\frac{\|Tx - x\|^2 + \|Ty - y\|^2}{\|Tx - x\| + \|Ty - y\|} \right) + \beta \|x - y\|,$$

for all $x, y \in S$ and $x \neq y$, $\alpha \in [0, 1/2)$, $\beta \geq 0$ and $2\alpha + \beta < 1$.

3. main theorem

Now we express the main theorem in the following

THEOREM 3.1. *Let S be a non empty closed subset of a Hilbert module Space E . Let T be self mapping on S satisfying the condition,*

$$\begin{aligned} \|Tx - Ty\| &\leq a_1 \left(\frac{\|Tx - x\|^2 + \|Ty - y\|^2 + \|Ty - x\|^2 + \|Tx - y\|^2}{\|Tx - x\| + \|Ty - y\| + \|Ty - x\| + \|Tx - y\|} \right) \\ &+ a_2 \left(\frac{\|Tx - x\|^2 + \|Ty - y\|^2}{\|Tx - x\| + \|Ty - y\|} \right) \\ &+ a_3 \left(\frac{\|Ty - x\|^2 + \|Tx - y\|^2}{\|Ty - x\| + \|Tx - y\|} \right) + a_4 \|x - y\|, \end{aligned}$$

for all $x, y \in S$ and $a_1, a_2, a_3, a_4 \geq 0$ and $4a_1 + 2a_2 + 2a_3 + a_4 < 1$, then T has a unique fixed point in S .

Proof. Let S be a non empty closed subset of a Hilbert module Space E . Let T be a self mapping on S . Let x_0 be any arbitrary point in S . We define a sequence $\{x_n\}_{n=1}^\infty$ in S by

$$x_{n+1} = Tx_n = T^{n+1}x_0, \text{ for } n = 0, 1, 2, \dots$$

Suppose that $x_{n+1} \neq x_n$, for $n = 0, 1, 2, \dots$

It follows from assumptions that for any integer $n \geq 1$

$$\begin{aligned} \|x_{n+1} - x_n\| &= \|Tx_n - Tx_{n-1}\| \\ &\leq a_1 \left(\frac{\|x_n - Tx_n\|^2 + \|x_{n-1} - Tx_{n-1}\|^2 + \|x_n - Tx_{n-1}\|^2 + \|x_{n-1} - Tx_n\|^2}{\|x_n - Tx_n\| + \|x_{n-1} - Tx_{n-1}\| + \|x_n - Tx_{n-1}\| + \|x_{n-1} - Tx_n\|} \right) \\ &+ a_2 \left(\frac{\|x_n - Tx_n\|^2 + \|x_{n-1} - Tx_{n-1}\|^2}{\|x_n - Tx_n\| + \|x_{n-1} - Tx_{n-1}\|} \right) \\ &+ a_3 \left(\frac{\|x_n - Tx_{n-1}\|^2 + \|x_{n-1} - Tx_n\|^2}{\|x_n - Tx_{n-1}\| + \|x_{n-1} - Tx_n\|} \right) + a_4 \|x_{n+1} - x_n\| \\ &\leq a_1 \left(\frac{\|x_n - x_{n+1}\|^2 + \|x_{n-1} - x_n\|^2 + \|x_n - x_n\|^2 + \|x_{n-1} - x_{n+1}\|^2}{\|x_n - x_{n+1}\| + \|x_{n-1} - x_n\| + \|x_n - x_n\| + \|x_{n-1} - x_{n+1}\|} \right) \\ &+ a_2 \left(\frac{\|x_n - x_{n+1}\|^2 + \|x_{n-1} - x_n\|^2}{\|x_n - x_{n+1}\| + \|x_{n-1} - x_n\|} \right) \\ &+ a_3 \left(\frac{\|x_n - x_n\|^2 + \|x_{n-1} - x_{n+1}\|^2}{\|x_n - x_n\| + \|x_{n-1} - x_{n+1}\|} \right) + a_4 \|x_n - x_{n-1}\| \\ &\leq a_1 (\|x_n - x_{n+1}\| + \|x_{n-1} - x_n\| + \|x_{n-1} - x_{n+1}\|) \\ &+ a_2 (\|x_n - x_{n+1}\| + \|x_{n-1} - x_n\|) \\ &+ a_3 (\|x_{n-1} - x_{n+1}\|) + a_4 \|x_n - x_{n-1}\| \\ &\leq a_1 (\|x_n - x_{n+1}\| + \|x_{n-1} - x_n\| + \|x_{n-1} - x_n\| + \|x_n - x_{n+1}\|) \\ &+ a_2 (\|x_n - x_{n+1}\| + \|x_{n-1} - x_n\|) \\ &+ a_3 (\|x_{n-1} - x_n\| + \|x_n - x_{n+1}\|) + a_4 \|x_n - x_{n-1}\|, \end{aligned}$$

therefore

$$\|x_{n+1} - x_n\| \leq (2a_1 + a_2 + a_3 + a_4) \|x_n - x_{n-1}\| + (2a_1 + a_2 + a_3) \|x_{n+1} - x_n\|,$$

so we have

$$\|x_{n+1} - x_n\| \leq k \|x_n - x_{n-1}\|,$$

where $k = \frac{2a_1 + a_2 + a_3 + a_4}{1 - 2a_1 - a_2 - a_3}$ and according to the assumption of the theorem $4a_1 + 2a_2 + 2a_3 + a_4 \leq 1$. Similarly to above proof we have $\|x_n - x_{n-1}\| \leq k \|x_{n-1} - x_{n-2}\|, \dots, \|x_2 - x_1\| \leq k \|x_1 - x_0\|$, therefore

$$\|x_{n+1} - x_n\| \leq k \|x_n - x_{n-1}\| \leq k^2 \|x_{n-1} - x_{n-2}\| \leq k^3 \|x_{n-2} - x_{n-3}\| \leq \dots \leq k^n \|x_1 - x_0\|.$$

Now for any positive integer $p \geq n \geq 1$ we have

$$\begin{aligned} \|x_n - x_p\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - x_{n+2}\| + \dots + \|x_{p-1} - x_p\| \\ &\leq k^n \|x_1 - x_0\| + k^{n+1} \|x_1 - x_0\| + \dots + k^{p-1} \|x_1 - x_0\| \\ &\leq (k^n + k^{n+1} + \dots + k^{p-1}) \|x_1 - x_0\| \\ &\leq k^n (1 + k^n + \dots + k^{p-n-1}) \|x_1 - x_0\|, \end{aligned}$$

thus

$$\|x_n - x_p\| \leq \frac{k^n}{1-k} \|x_1 - x_0\| \longrightarrow 0 \text{ as } n \longrightarrow \infty.$$

So $\{x_n\}_{n=1}^{\infty}$ is a Cauchy sequence. Since S is closed subset of Hilbert module E , the sequence $\{x_n\}_{n=1}^{\infty}$ converges to point $\nu \in S$. Now we show that ν is a fixed point of self mapping T . Let $T\nu \neq \nu$. Consider

$$\begin{aligned} \|\nu - T\nu\| &= \|\nu - x_n + x_n - T\nu\| \leq \|\nu - x_n\| + \|x_n - T\nu\| \leq \|x_n - T\nu\| \\ &\leq \|Tx_{n-1} - T\nu\| \quad (\text{as } n \longrightarrow \infty, x \longrightarrow \nu) \\ &\leq a_1 \left(\frac{\|x_{n-1} - Tx_{n-1}\|^2 + \|\nu - T\nu\|^2 + \|x_{n-1} - T\nu\|^2 + \|\nu - Tx_{n-1}\|^2}{\|x_{n-1} - Tx_{n-1}\| + \|\nu - T\nu\| + \|x_{n-1} - T\nu\| + \|\nu - Tx_{n-1}\|} \right) \\ &+ a_2 \left(\frac{\|x_{n-1} - Tx_{n-1}\|^2 + \|\nu - T\nu\|^2}{\|x_{n-1} - Tx_{n-1}\| + \|\nu - T\nu\|} \right) \\ &+ a_3 \left(\frac{\|x_{n-1} - T\nu\|^2 + \|\nu - Tx_{n-1}\|^2}{\|x_{n-1} - T\nu\| + \|\nu - Tx_{n-1}\|} \right) + a_4 \|x_{n-1} - \nu\| \\ &\leq a_1 \left(\frac{\|x_{n-1} - x_n\|^2 + \|\nu - T\nu\|^2 + \|x_{n-1} - T\nu\|^2 + \|\nu - x_n\|^2}{\|x_{n-1} - x_n\|^2 + \|\nu - T\nu\|^2 + \|x_{n-1} - T\nu\|^2 + \|\nu - x_n\|^2} \right) \\ &+ a_2 \left(\frac{\|x_{n-1} - x_n\|^2 + \|\nu - T\nu\|^2}{\|x_{n-1} - x_n\| + \|\nu - T\nu\|} \right) + a_3 \left(\frac{\|x_{n-1} - T\nu\|^2 + \|\nu - x_n\|^2}{\|x_{n-1} - T\nu\| + \|\nu - x_n\|} \right) \\ &+ a_4 \|x_{n-1} - \nu\| \end{aligned}$$

where we have used the assumptions, hence

$$\begin{aligned} \|\nu - T\nu\| &\leq a_1 (\|x_{n-1} - x_n\| + \|\nu - T\nu\| + \|x_{n-1} - T\nu\| + \|\nu - x_n\|) \\ &+ a_2 (\|x_{n-1} - x_n\| + \|\nu - T\nu\|) + a_3 (\|x_{n-1} - T\nu\| + \|\nu - x_n\|) \\ &+ a_4 \|x_{n-1} - \nu\|, \end{aligned}$$

so

$$\begin{aligned} \|\nu - T\nu\| &\leq a_1 (\|\nu - T\nu\| + \|\nu - T\nu\|) + a_2 \|\nu - T\nu\| \\ &+ a_3 \|\nu - T\nu\| \quad x_{n-1}, x_n \longrightarrow \nu \text{ as } n \longrightarrow \infty \end{aligned}$$

thus

$$\begin{aligned} \|\nu - T\nu\| &\leq (2a_1 + a_2 + a_3) \|\nu - T\nu\| \\ &\rightarrow (1 - 2a_1 - a_2 - a_3) \|\nu - T\nu\| \leq 0 \\ &\rightarrow \nu = T\nu. \end{aligned}$$

Hence ν is fixed point for the self mapping T .

Uniqueness: Let $\nu \neq w$ such that $Tw = w$, take

$$\begin{aligned} \|\nu - w\| = \|T\nu - Tw\| &\leq a_1 \left(\frac{\|\nu - T\nu\|^2 + \|w - Tw\|^2 + \|\nu - Tw\|^2 + \|w - T\nu\|^2}{\|\nu - T\nu\| + \|w - Tw\| + \|\nu - Tw\| + \|w - T\nu\|} \right) \\ &+ a_2 \left(\frac{\|\nu - T\nu\|^2 + \|w - Tw\|^2}{\|\nu - T\nu\| + \|w - Tw\|} \right) + a_3 \left(\frac{\|\nu - Tw\|^2 + \|w - T\nu\|^2}{\|\nu - Tw\| + \|w - T\nu\|} \right) \\ &+ a_4 \|\nu - w\|, \end{aligned}$$

so we have

$$\|\nu - w\| \leq (2a_1 + a_3 + a_4) \|\nu - w\| \Rightarrow \|\nu - w\| = 0 \Rightarrow \nu = w.$$

Hence ν is a unique fixed point. \square

THEOREM 3.2. *Let S be a non empty closed subset of a Hilbert module space E . Let T_1 and T_2 are self mapping on S satisfying the condition,*

$$\begin{aligned} \|T_1x - T_2y\| &\leq a_1\left(\frac{\|x - T_1x\|^2 + \|y - T_2y\|^2 + \|x - T_2y\|^2 + \|y - T_1x\|^2}{\|x - T_1x\| + \|y - T_2y\| + \|x - T_2y\| + \|y - T_1x\|}\right) \\ &+ a_2\left(\frac{\|x - T_1x\|^2 + \|y - T_2y\|^2}{\|x - T_1x\| + \|y - T_2y\|}\right) \\ &+ a_3\left(\frac{\|x - T_2y\|^2 + \|y - T_1x\|^2}{\|x - T_2y\| + \|y - T_1x\|}\right) + a_4\|x - y\|, \end{aligned}$$

for all $x, y \in S$ and $a_1, a_2, a_3, a_4 \geq 0$ and $4a_1 + 2a_2 + 2a_3 + a_4 < 1$, then T_1 and T_2 have a unique common fixed point in S .

Proof. Let S be a non empty closed subset of a Hilbert module Space E . Let T_1 and T_2 are self mappings on S . Let x_o be any arbitrary point in S . We define a sequence $\{x_n\}_{n=1}^\infty$ in S by $x_{2n+1} = T_1x_{2n}$ and $x_{2n+2} = T_2x_{2n+1}$ for $n = 0, 1, 2, \dots$. Suppose that for some n , $x_{n+2} \neq x_{n+1} \neq x_n$, for $n = 0, 1, 2, \dots$. By the assumptions, for any integer $n \geq 1$

$$\begin{aligned} &\|x_{2n+1} - x_{2n}\| \\ &= \|T_1x_{2n} - T_2x_{2n-1}\| \\ &\leq a_1\left(\frac{\|x_{2n} - T_1x_{2n}\|^2 + \|x_{2n-1} - T_2x_{2n-1}\|^2 + \|x_{2n} - T_2x_{2n-1}\|^2 + \|x_{2n-1} - T_1x_{2n}\|^2}{\|x_{2n} - T_1x_{2n}\| + \|x_{2n-1} - T_2x_{2n-1}\| + \|x_{2n} - T_2x_{2n-1}\| + \|x_{2n-1} - T_1x_{2n}\|}\right) \\ &+ a_2\left(\frac{\|x_{2n} - T_1x_{2n}\|^2 + \|x_{2n-1} - T_2x_{2n-1}\|^2}{\|x_{2n} - T_1x_{2n}\| + \|x_{2n-1} - T_2x_{2n-1}\|}\right) \\ &+ a_3\left(\frac{\|x_{2n} - T_2x_{2n-1}\|^2 + \|x_{2n-1} - T_1x_{2n}\|^2}{\|x_{2n} - T_2x_{2n-1}\| + \|x_{2n-1} - T_1x_{2n}\|}\right) + a_4\|x_{2n} - x_{2n-1}\| \\ &\leq a_1\left(\frac{\|x_{2n} - x_{2n+1}\|^2 + \|x_{2n-1} - x_{2n}\|^2 + \|x_{2n} - x_{2n}\|^2 + \|x_{2n-1} - x_{2n+1}\|^2}{\|x_{2n} - x_{2n+1}\| + \|x_{2n-1} - x_{2n}\| + \|x_{2n} - x_{2n}\| + \|x_{2n-1} - x_{2n+1}\|}\right) \\ &+ a_2\left(\frac{\|x_{2n} - x_{2n+1}\|^2 + \|x_{2n-1} - x_{2n}\|^2}{\|x_{2n} - x_{2n+1}\| + \|x_{2n-1} - x_{2n}\|}\right) \\ &+ a_3\left(\frac{\|x_{2n} - x_{2n}\|^2 + \|x_{2n-1} - x_{2n+1}\|^2}{\|x_{2n} - x_{2n}\| + \|x_{2n-1} - x_{2n+1}\|}\right) + a_4\|x_{2n} - x_{2n-1}\| \\ &\leq a_1(\|x_{2n} - x_{2n+1}\| + \|x_{2n-1} - x_{2n}\| + \|x_{2n-1} - x_{2n+1}\|) \\ &+ a_2(\|x_{2n} - x_{2n+1}\| + \|x_{2n-1} - x_{2n}\|) + a_3(\|x_{2n-1} - x_{2n+1}\|) \\ &+ a_4\|x_{2n} - x_{2n-1}\| \\ &\leq a_1(\|x_{2n} - x_{2n+1}\| + \|x_{2n-1} - x_{2n}\| + \|x_{2n-1} - x_{2n}\| + \|x_{2n} - x_{2n+1}\|) \\ &+ a_2(\|x_{2n} - x_{2n+1}\| + \|x_{2n-1} - x_{2n}\|) + a_3(\|x_{2n-1} - x_{2n}\| + \|x_{2n} - x_{2n+1}\|) \\ &+ a_4\|x_{2n} - x_{2n-1}\|, \end{aligned}$$

therefore

$$\|x_{2n+1} - x_{2n}\| \leq (2a_1 + a_2 + a_3 + a_4)\|x_{2n} - x_{2n-1}\| + (2a_1 + a_2 + a_3)\|x_{2n+1} - x_{2n}\|,$$

so we have $\|x_{2n+1} - x_{2n}\| \leq k\|x_{2n} - x_{2n-1}\|$ where $k = \frac{2a_1+a_2+a_3+a_4}{1-2a_1-a_2-a_3}$ and $4a_1 + 2a_2 + 2a_3 + a_4 < 1$. In general

$$\begin{aligned} \|x_{n+1} - x_n\| \leq k\|x_n - x_{n-1}\| &\leq k^2\|x_{n-1} - x_{n-2}\| \leq k^3\|x_{n-2} - x_{n-3}\| \leq \dots \leq k^n\|x_1 - x_0\| \\ \implies \|x_{n+1} - x_n\| &\leq k^n\|x_1 - x_0\|. \end{aligned}$$

Now for any positive integer $p \geq n \geq 1$

$$\begin{aligned} \|x_n - x_p\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - x_{n+2}\| + \dots + \|x_{p-1} - x_p\| \\ &\leq k^n\|x_1 - x_0\| + k^{n+1}\|x_1 - x_0\| + \dots + k^{p-1}\|x_1 - x_0\| \\ &\leq (k^n + k^{n+1} + \dots + k^{p-1})\|x_1 - x_0\| \\ &\leq k^n(1 + k^n + \dots + k^{p-n-1})\|x_1 - x_0\| \\ &\leq \frac{k^n}{1-k}\|x_1 - x_0\| \longrightarrow 0 \text{ as } n \longrightarrow \infty. \end{aligned}$$

$\{x_n\}_{n=1}^\infty$ is a Cauchy sequence. Since S is closed subset of a Hilbert module space E , the sequence $\{x_n\}_{n=1}^\infty$ converges to a point ν in S . Now we show that ν is a common fixed point of T_1 and T_2 . First we see that ν is fixed point of T_1 . Let $T_1\nu \neq \nu$. Consider

$$\begin{aligned} \|\nu - T_1\nu\| &= \|\nu - x_{2n+2} + x_{2n+2} - T_1\nu\| \\ &\leq \|\nu - x_{2n+2}\| + \|x_{2n+2} - T_1\nu\| \text{ as } n \longrightarrow \infty, x_{2n+2} \longrightarrow \nu \\ &\leq \|T_2x_{2n+1} - T_1\nu\| + \|T_1\nu - T_2x_{2n+1}\| \\ &\leq a_1\left(\frac{\|\nu - T_1\nu\|^2 + \|x_{2n+1} - T_2x_{2n+1}\|^2 + \|\nu - T_2x_{2n+1}\|^2 + \|x_{2n+1} - T_1\nu\|^2}{\|\nu - T_1\nu\| + \|x_{2n+1} - T_2x_{2n+1}\| + \|\nu - T_2x_{2n+1}\| + \|x_{2n+1} - T_1\nu\|}\right) \\ &\quad + a_2\left(\frac{\|\nu - T_1\nu\|^2 + \|x_{2n+1} - T_2x_{2n+1}\|^2}{\|\nu - T_1\nu\| + \|x_{2n+1} - T_2x_{2n+1}\|}\right) \\ &\quad + a_3\left(\frac{\|\nu - T_2x_{2n+1}\|^2 + \|x_{2n+1} - T_1\nu\|^2}{\|\nu - T_2x_{2n+1}\| + \|x_{2n+1} - T_1\nu\|}\right) + a_4\|x_{2n+1} - \nu\| \\ &\leq a_1\left(\frac{\|\nu - T_1\nu\|^2 + \|x_{2n+1} - x_{2n+2}\|^2 + \|\nu - x_{2n+2}\|^2 + \|x_{2n+1} - T_1\nu\|^2}{\|\nu - T_1\nu\| + \|x_{2n+1} - x_{2n+2}\| + \|\nu - x_{2n+2}\| + \|x_{2n+1} - T_1\nu\|}\right) \\ &\quad + a_2\left(\frac{\|\nu - T_1\nu\|^2 + \|x_{2n+1} - x_{2n+2}\|^2}{\|\nu - T_1\nu\| + \|x_{2n+1} - x_{2n+2}\|}\right) + a_3\left(\frac{\|\nu - x_{2n+2}\|^2 + \|x_{2n+1} - T_1\nu\|^2}{\|\nu - x_{2n+2}\| + \|x_{2n+1} - T_1\nu\|}\right) \\ &\quad + a_4\|x_{2n+1} - \nu\| \end{aligned}$$

so we have

$$\begin{aligned} \|\nu - T_1\nu\| &\leq a_1(\|\nu - T_1\nu\| + \|\nu - T_1\nu\|) + a_2\|\nu - T_1\nu\| \\ &\quad + a_3\|\nu - T_1\nu\| + a_4\|\nu - x_{2n+1}\|. \end{aligned}$$

Therefore when $n \longrightarrow \infty$ then $x_{2n+1}, x_{2n+2} \longrightarrow \nu$, hence we conclude

$$\begin{aligned} \|\nu - T_1\nu\| &\leq (2a_1 + a_2 + a_3)\|\nu - T_1\nu\| \\ \implies (1 - 2a_1 - a_2 - a_3)\|\nu - T_1\nu\| &\leq 0 \\ \implies \nu &= T_1\nu. \end{aligned}$$

Hence ν is fixed point for the self mapping T_1 . Similarly we can show that ν is a fixed point of T_2 . Hence ν is a common fixed point of T_1 and T_2 .

Uniqueness: Let $w \neq \nu$ be another fixed point such that $T_2w = w$.

Take

$$\begin{aligned} \|\nu - w\| &= \|T_1\nu - T_2w\| \\ &\leq a_1\left(\frac{\|\nu - T_1\nu\|^2 + \|w - T_2w\|^2 + \|\nu - T_2w\|^2 + \|w - T_1\nu\|^2}{\|\nu - T_1\nu\| + \|w - T_2w\| + \|\nu - T_2w\| + \|T_1w - \nu\|}\right) \\ &+ a_2\left(\frac{\|\nu - T_1\nu\|^2 + \|w - T_2w\|^2}{\|\nu - T_1\nu\| + \|w - T_2w\|}\right) + a_3\left(\frac{\|\nu - T_2w\|^2 + \|w - T_1\nu\|^2}{\|\nu - T_2w\| + \|w - T_1\nu\|}\right) \\ &+ a_4\|\nu - w\|. \end{aligned}$$

Thus we arrive to

$$\begin{aligned} \|\nu - w\| &\leq (2a_1 + a_2 + a_4)\|\nu - w\| \\ &\Rightarrow \|\nu - w\| = 0 \Rightarrow \nu = w. \end{aligned}$$

Hence ν is a unique common fixed point of T_1 and T_2 . □

4. Conclusion

In this paper we obtained a unique fixed point for a continuous self mapping T as well as a unique common fixed point for two self mappings T_1 and T_2 satisfying rational conditions in Hilbert module space.

References

- [1] Browder, F.E, *Fixed point theorems for noncompact mappings in Hilbert spaces*, Proc. Natl. Acad. Sci. USA. **43** (1965), 1272–1276.
- [2] Browder, F.E, *Nonexpansive nonlinear operators in a Banach space*, Proc. Nat. Acad. Sei. U.S.A. **54** (1965), 1041–1044.
- [3] Fisher, B, *A fixed point theorem for compact metric space*, Publ. Inst. Math. **25** (1976), 193–194.
- [4] Göhde, D, *Zum Prinzip der kontraktiven Abbildung*, Math. Nachr. **30** (1965), 251–258.
- [5] Kannan, R, *Some results on fixed point theorems*, Bull. Calcutta. Math. Soc. **60** (1969), 71–78.
- [6] Kaplansky, I, *Modules over operator algebras*, Amer. J. Math. **75** (4) (1953), 839–853.
- [7] Kirk, W.A, *A fixed point theorem for mappings which do not increase distance*, Amer. Math. Monthly. **72** (1965), 1004–1006.
- [8] Koparde, P.V. and Waghmode, B.B, *Kannan type mappings in Hilbert spaces*, Scientist Phyl. Sciences **3** (1) (1991), 45–50.
- [9] Lance, E.C, *Hilbert C^* -modules*, A toolkit for operator algebraists, Cambridge Univ. Press, Cambridge 1995.
- [10] Phillips, N.C. and Weaver, N, *Modules with norms which take values in a C^* -algebra*, Pacific J. of Maths, **185** (1) (1998), 163–181.
- [11] Sharma, A.K., Badshah, V.H and Gupta, V.K, *Common fixed point theorems in Hilbert space*, International Academy of Sciences, Engg. And Tech. **3** (2014), 63–70.

Hasan Ranjbar

Department of Mathematics, Mashhad Branch, Islamic Azad University,
Mashhad, Iran.

E-mail: hasan.ranjbar514@gmail.com

Asadollah Niknam

Department of Mathematics, Mashhad Branch, Islamic Azad University,
Mashhad, Iran.

E-mail: dassamankin@yahoo.co.uk