

## HANKEL DETERMINANT PROBLEMS FOR CERTAIN SUBCLASSES OF SAKAGUCHI TYPE FUNCTIONS DEFINED WITH SUBORDINATION

GAGANDEEP SINGH\* AND GURCHARANJIT SINGH

ABSTRACT. The present investigation is concerned with the estimation of initial coefficients, Fekete-Szegő inequality, second Hankel determinants, Zalcman functionals and third Hankel determinants for certain subclasses of Sakaguchi type functions defined with subordination in the open unit disc  $E = \{z \in \mathbb{C} : |z| < 1\}$ . The results derived in this paper will pave the way for the further study in this direction.

### 1. Introduction

Let  $\mathcal{A}$ , be the class of all analytic functions of the form  $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$  defined in the open unit disc  $E = \{z \in \mathbb{C} : |z| < 1\}$  and normalized by the conditions  $f(0) = f'(0) - 1 = 0$ . Let  $\mathcal{S}$  denote the subclass of  $\mathcal{A}$ , consisting of functions which are univalent in  $E$ . For two analytic functions  $f$  and  $g$  in  $E$ ,  $f$  is said to be subordinate to  $g$  (symbolically  $f \prec g$ ) if there exists a function  $w$  with  $w(0) = 0$  and  $|w(z)| < 1$  for  $z \in E$  such that  $f(z) = g(w(z))$ . Further, if  $g$  is univalent in  $E$ , then  $f \prec g$  is equivalent to  $f(0) = g(0)$  and  $f(E) \subset g(E)$ .

A very famous result in the theory of univalent functions was Bieberbach's conjecture established by Bieberbach [5]. According to this conjecture, for  $f \in \mathcal{S}$ ,  $|a_n| \leq n$ ,  $n = 2, 3, \dots$ . This conjecture remained as a challenge for the mathematicians for a long time. Finally, it was L. De-Branges [7], who proved this conjecture in 1985. During the course of proving this conjecture, various results related to the coefficients were established and some new subclasses of  $\mathcal{S}$  were developed. The well-known classes of starlike and convex functions are denoted by  $\mathcal{S}^*$  and  $\mathcal{K}$ , respectively.

Sakaguchi [21] introduced the class  $\mathcal{S}_s^*$  consisting of analytic functions  $f \in \mathcal{A}$  and satisfying the condition

$$\operatorname{Re} \left( \frac{2zf'(z)}{f(z) - f(-z)} \right) > 0 \text{ or } \frac{2zf'(z)}{f(z) - f(-z)} \prec \frac{1+z}{1-z}.$$

---

Received September 8, 2021. Revised January 26, 2022. Accepted February 25, 2022.

2010 Mathematics Subject Classification: 30C45, 30C50.

Key words and phrases: Analytic functions, Sakaguchi type functions, subordination, coefficient problem, Hankel determinant.

\* Corresponding Author.

© The Kangwon-Kyungki Mathematical Society, 2022.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

The functions in the class  $\mathcal{S}_s^*$  are known as the starlike functions with respect to symmetric points.

Later on, Das and Singh [6] introduced the class  $\mathcal{K}_s$  consisting of analytic functions  $f \in \mathcal{A}$ , known as convex functions with respect to symmetric points, which satisfy the condition

$$\operatorname{Re} \left( \frac{2(zf'(z))'}{(f(z) - f(-z))'} \right) > 0 \text{ or } \frac{2(zf'(z))'}{(f(z) - f(-z))'} \prec \frac{1+z}{1-z}.$$

Clearly,  $f \in \mathcal{K}_s$  if and only if  $zf' \in \mathcal{S}_s^*$ .

Sokol and Stankiewicz [25] introduced the class  $\mathcal{SL}^*$  consisting of analytic functions  $f \in \mathcal{A}$  and satisfying the condition

$$\left| \left( \frac{zf'(z)}{f(z)} \right)^2 - 1 \right| < 1 \text{ or } \frac{zf'(z)}{f(z)} \prec \sqrt{1+z}.$$

The superordinate function  $\sqrt{1+z}$  maps the unit disc  $E$  onto the right-half of the lemniscate of Bernoulli which has the equation

$$(x^2 + y^2)^2 - 2(x^2 - y^2) = 0.$$

From time to time, various subclasses of  $\mathcal{S}$  were studied by subordinating to the function  $\sqrt{1+z}$  by various researchers including, Najafzadeh et al. [17], Singh et al. [24], Ali et al. [1], Sokol and Thomas [26] and Ullah et al. [27]. Getting inspired from these works, now we define the following subclasses of Sakaguchi type functions by subordinating to  $\sqrt{1+z}$ .

Let  $\mathcal{S}_s^*(\mathcal{L})$  denote the class which consists of analytic functions  $f \in \mathcal{A}$  satisfying the condition

$$\left( \frac{2zf'(z)}{f(z) - f(-z)} \right) \prec \sqrt{1+z}.$$

By  $\mathcal{K}_s(\mathcal{L})$ , we denote the class which consists of analytic functions  $f \in \mathcal{A}$  satisfying the condition

$$\left( \frac{2(zf'(z))'}{(f(z) - f(-z))'} \right) \prec \sqrt{1+z}.$$

For the complex sequence  $a_n, a_{n+1}, a_{n+2}, \dots$ , the  $q \times q$  Hankel matrix, named after Hermann Hankel (1839-1873), is defined as

$$\begin{pmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ a_{n+q-1} & \cdots & \cdots & a_{n+2q-2} \end{pmatrix} \text{ where } q \in \mathbb{N} - \{1\}.$$

We observe that the Hankel matrix has constant positive slopping diagonals whose entries also satisfy:

$$a_{ij} = a_{i-1, j+1} (i \in \mathbb{N} - \{1\}; j \in \mathbb{N}).$$

For basic properties of the hankel matrix, we refer to [9, 10]. In 1976, Noonan and Thomas [18] stated the  $q^{th}$  Hankel determinant for  $q \geq 1$  and  $n \geq 1$  as

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ a_{n+q-1} & \dots & \dots & a_{n+2q-2} \end{vmatrix}.$$

Particularly, for  $q = 2, n = 1, a_1 = 1$  and  $q = 2, n = 2$ , the Hankel determinant simplifies respectively to

$$H_2(1) = a_3 - a_2^2 \text{ and } H_2(2) = a_2a_4 - a_3^2.$$

Easily, one can observe that the Fekete-Szegő functional is  $H_2(1)$ . Fekete and Szegő [8] then further generalised the estimate  $|a_3 - \mu a_2^2|$  where  $\mu$  is real and  $f \in \mathcal{S}$ . Also  $H_2(2)$  is called the second Hankel determinant.

The functional  $J_{n,m}(f) = a_n a_m - a_{m+n-1}$ ,  $n, m \in \mathbb{N} - \{1\}$ , was investigated by Ma [15] and it is known as generalized Zalcman functional. The functional  $J_{2,3}(f) = a_2 a_3 - a_4$  is a specific case of the generalized Zalcman functional. Various authors computed the upper bound for the functional  $J_{2,3}(f)$  over different subclasses of analytic functions.

The Hankel determinant in the case  $q = 3, n = 1$ ,

$$H_3(1) = \begin{vmatrix} a_1 & a_2 & a_3 \\ a_2 & a_3 & a_4 \\ a_3 & a_4 & a_5 \end{vmatrix}$$

is called the Third Hankel determinant.

For  $f \in \mathcal{S}$  and  $a_1 = 1$ , we have

$$H_3(1) = a_3(a_2a_4 - a_3^2) - a_4(a_4 - a_2a_3) + a_5(a_3 - a_2^2).$$

By using the triangle inequality, the above equation yields

$$(1) \quad |H_3(1)| \leq |a_3||a_2a_4 - a_3^2| + |a_4||a_2a_3 - a_4| + |a_5||a_3 - a_2^2| \dots$$

For various subclasses of  $\mathcal{S}$ , the second Hankel determinants has been extensively studied by various authors including Mehrok and Singh [16], Janteng et al. [11] and many others, while the third Hankel determinants were studied by the authors including Babalola [4], Shanmugam et al. [22], Altinkaya and Yalcin [2] and Singh and Singh [23].

In the present paper, we study Fekete-Szegő inequalities, Second Hankel determinants, Zalcman functionals and third Hankel determinants for the classes  $\mathcal{S}_s^*(\mathcal{L})$  and  $\mathcal{K}_s(\mathcal{L})$ .

To prove our result, we shall make use of the following lemmas:

By  $\mathcal{P}$ , we denote the class of analytic functions  $p$  of the form

$$p(z) = 1 + \sum_{k=1}^{\infty} p_k z^k,$$

whose real parts are positive in  $E$ .

LEMMA 1.1. [12, 19] *If  $p \in \mathcal{P}$ , then*

$$(2) \quad |p_k| \leq 2, k \in \mathbb{N} \dots,$$

$$(3) \quad \left| p_2 - \frac{p_1^2}{2} \right| \leq 2 - \frac{|p_1|^2}{2} \dots,$$

$$(4) \quad |p_{i+j} - \mu p_i p_j| \leq 2, 0 \leq \mu \leq 1 \dots,$$

and for complex number  $\rho$ , we have

$$(5) \quad |p_2 - \rho p_1^2| \leq 2 \max\{1, |2\rho - 1|\} \dots$$

LEMMA 1.2. [13, 14] *If  $p \in \mathcal{P}$ , then*

$$2p_2 = p_1^2 + (4 - p_1^2)x,$$

$$4p_3 = p_1^3 + 2p_1(4 - p_1^2)x - p_1(4 - p_1^2)x^2 + 2(4 - p_1^2)(1 - |x|^2)z,$$

for  $|x| \leq 1$  and  $|z| \leq 1$ .

LEMMA 1.3. [3] *Let  $p \in \mathcal{P}$ , then*

$$|Jp_1^3 - Kp_1p_2 + Lp_3| \leq 2|J| + 2|K - 2J| + 2|J - K + L|.$$

In particular, it is proved in [19] that

$$|p_1^3 - 2p_1p_2 + p_3| \leq 2.$$

LEMMA 1.4. [20] *Let  $m, n, l$  and  $r$  satisfy the inequalities  $0 < m < 1$ ,  $0 < r < 1$  and*

$$8r(1-r) [(mn - 2l)^2 + (m(r+m) - n)^2] + m(1-m)(n - 2rm)^2 \leq 4m^2(1-m)^2r(1-r).$$

If  $p \in \mathcal{P}$ , then

$$\left| lp_1^4 + rp_2^2 + 2mp_1p_3 - \frac{3}{2}np_1^2p_2 - p_4 \right| \leq 2.$$

## 2. The class $\mathcal{S}_s^*(\mathcal{L})$

THEOREM 2.1. *If  $f \in \mathcal{S}_s^*(\mathcal{L})$ , then*

$$(6) \quad |a_2| \leq \frac{1}{4} \dots,$$

$$(7) \quad |a_3| \leq \frac{1}{4} \dots,$$

$$(8) \quad |a_4| \leq \frac{1}{8} \dots,$$

and

$$(9) \quad |a_5| \leq \frac{1}{8} \dots$$

*Proof.* As  $f \in \mathcal{S}_s^*(\mathcal{L})$ , by the principle of subordination, we have

$$(10) \quad \frac{2zf'(z)}{f(z) - f(-z)} = \sqrt{1 + w(z)} \dots$$

Define  $p(z) = \frac{1 + w(z)}{1 - w(z)} = 1 + p_1z + p_2z^2 + p_3z^3 + \dots$ , which implies  $w(z) = \frac{p(z) - 1}{p(z) + 1}$ .

Also  $\frac{2zf'(z)}{f(z) - f(-z)} = 1 + 2a_2z + 2a_3z^2 + (4a_4 - 2a_2a_3)z^3 + (4a_5 - 2a_3^2)z^4 + \dots$

Again  $\sqrt{1 + w(z)} = \left(\frac{2p(z)}{p(z) + 1}\right)^{\frac{1}{2}} = 1 + \frac{1}{4}p_1z + \left(\frac{p_2}{4} - \frac{5p_1^2}{32}\right)z^2 + \left(\frac{13p_1^3}{128} - \frac{5p_1p_2}{16} + \frac{p_3}{4}\right)z^3 + \left(\frac{-141p_1^4}{2048} + \frac{39p_1^2p_2}{128} - \frac{5p_3p_1}{16} - \frac{5p_2^2}{32} + \frac{p_4}{4}\right)z^4 + \dots$

On comparing (10), it yields

$$(11) \quad 1 + 2a_2z + 2a_3z^2 + (4a_4 - 2a_2a_3)z^3 + (4a_5 - 2a_3^2)z^4 + \dots = 1 + \frac{1}{4}p_1z + \left(\frac{p_2}{4} - \frac{5p_1^2}{32}\right)z^2 + \left(\frac{13p_1^3}{128} - \frac{5p_1p_2}{16} + \frac{p_3}{4}\right)z^3 + \left(\frac{-141p_1^4}{2048} + \frac{39p_1^2p_2}{128} - \frac{5p_3p_1}{16} - \frac{5p_2^2}{32} + \frac{p_4}{4}\right)z^4 + \dots$$

Equating the coefficients of  $z, z^2, z^3$  and  $z^4$  in (11) and on simplifying, we obtain

$$(12) \quad a_2 = \frac{p_1}{8} \dots,$$

$$(13) \quad a_3 = \frac{p_2}{8} - \frac{5p_1^2}{64} \dots,$$

$$(14) \quad a_4 = \frac{21p_1^3}{1024} - \frac{9p_1p_2}{128} + \frac{p_3}{16} \dots,$$

and

$$(15) \quad a_5 = -\frac{1}{16} \left[ \frac{29}{128}p_1^4 + \frac{p_2^2}{2} + \frac{5p_3p_1}{4} - \frac{34}{32}p_1^2p_2 - p_4 \right] \dots$$

Using (2) in (12), it gives (6).

From (13), we have

$$|a_3| = \frac{1}{8} \left| p_2 - \frac{5}{8}p_1^2 \right|.$$

By using (5), it yields

$$|a_3| \leq \frac{1}{8} \cdot 2 \max \left\{ 1, \frac{1}{4} \right\},$$

which gives (7).

On applying Lemma 3 in (14), (8) can be easily obtained.

Using Lemma 4 in (15), the result (9) is obvious. □

**THEOREM 2.2.** *If  $f \in \mathcal{S}_s^*(\mathcal{L})$ , then*

$$(16) \quad |a_3 - \rho a_2^2| \leq \frac{1}{4} \max \left\{ 1, \left| \frac{1 + \rho}{4} \right| \right\} \dots$$

*Proof.* From (12) and (13), we have

$$(17) \quad |a_3 - \rho a_2^2| = \frac{1}{8} \left| p_2 - \left( \frac{5}{8} + \frac{1}{8}\rho \right) p_1^2 \right| \dots$$

An application of (5) in (17) leads us to (16).

For  $\rho = 1$ , (16) gives

$$(18) \quad |a_3 - a_2^2| \leq \frac{1}{4} \dots$$

□

**THEOREM 2.3.** *If  $f \in \mathcal{S}_s^*(\mathcal{L})$ , then*

$$(19) \quad |a_2 a_3 - a_4| \leq \frac{1}{8} \dots$$

*Proof.* From (12), (13) and (14), we have

$$(20) \quad |a_2 a_3 - a_4| = \left| \frac{31}{1024} p_1^3 - \frac{11}{128} p_1 p_2 + \frac{1}{16} p_3 \right| \dots$$

By implementing the triangle inequality and Lemma 3 in (20), it leads us to (19).

□

**THEOREM 2.4.** *If  $f \in \mathcal{S}_s^*(\mathcal{L})$ , then*

$$(21) \quad |a_2 a_4 - a_3^2| \leq \frac{1}{16} \dots$$

*The bound is sharp.*

*Proof.* Using (12), (13) and (14), we have

$$|a_2 a_4 - a_3^2| = \frac{1}{8192} \left| -29p_1^4 + 88p_1^2 p_2 + 64p_1 p_3 - 128p_2^2 \right|.$$

Substituting for  $p_2$  and  $p_3$  from Lemma 2 and letting  $p_1 = p$ , we get

$$|a_2 a_4 - a_3^2| = \frac{1}{8192} \left| -p^4 + 12p^2(4-p^2)x + 32p(4-p^2)(1-|x|^2)z - 16(4-p^2)(8-p^2)x^2 \right|.$$

Since  $|p| = |p_1| \leq 2$ , by using (2), we may assume that  $p \in [0, 2]$ . Then by using triangle inequality and  $|z| \leq 1$  with  $|x| = t \in [0, 1]$ , we obtain

$$\begin{aligned} & |a_2 a_4 - a_3^2| \\ & \leq \frac{1}{8192} \left[ p^4 + 32p(4-p^2) + 12p^2(4-p^2)t + 16(4-p^2)(8-2p-p^2)t^2 \right] = F(p, t). \end{aligned}$$

Then

$$\frac{\partial F}{\partial t} = \frac{1}{8192} [12(4-p^2)p^2 + 32t(4-p^2)(8-2p-p^2)] \geq 0.$$

Therefore  $F(p, t)$  is an increasing function of  $t$ .

$$\text{So } \max F(p, t) = F(p, 1) = \frac{1}{8192} [5p^4 - 144p^2 + 512] = G(p).$$

Now  $G'(p) = 0$  gives  $p = 0$ . Also  $G''(p) = \frac{1}{8192} [60p^2 - 288]$  which is negative for each  $p \in [0, 2]$ .

$$\text{This implies } \max G(p) = G(0) = \frac{512}{8192}.$$

Hence  $|a_2a_4 - a_3^2| \leq \frac{1}{16}$ .

The result is sharp for  $p_1 = 0$ ,  $p_2 = \pm 2$  and  $p_3 = 0$ .

□

**THEOREM 2.5.** *If  $f \in \mathcal{S}_s^*(\mathcal{L})$ , then*

$$(22) \quad |H_3(1)| \leq \frac{1}{16} \dots$$

*Proof.* By using (7), (8), (9), (18), (19) and (21) in (1), the result (22) is obvious.

□

### 3. The class $\mathcal{K}_s(\mathcal{L})$

**THEOREM 3.1.** *If  $f(z) \in \mathcal{K}_s(\mathcal{L})$ , then*

$$(23) \quad |a_2| \leq \frac{1}{8} \dots,$$

$$(24) \quad |a_3| \leq \frac{1}{12} \dots,$$

$$(25) \quad |a_4| \leq \frac{1}{32} \dots$$

and

$$(26) \quad |a_5| \leq \frac{1}{40} \dots$$

*Proof.* As  $f \in \mathcal{K}_s(\mathcal{L})$ , therefore

$$(27) \quad \frac{2(zf'(z))'}{(f(z) - f(-z))'} = \sqrt{1 + w(z)} \dots$$

On expanding as in Theorem 2.1, (27) yields

$$(28) \quad 1 + 4a_2z + 6a_3z^2 + (16a_4 - 12a_2a_3)z^3 + (20a_5 - 18a_3^2)z^4 + \dots = 1 + \frac{1}{4}p_1z + \left(\frac{p_2}{4} - \frac{5p_1^2}{32}\right)z^2 + \left(\frac{13p_1^3}{128} - \frac{5p_1p_2}{16} + \frac{p_3}{4}\right)z^3 + \left(\frac{-141p_1^4}{2048} + \frac{39p_1^2p_2}{128} - \frac{5p_3p_1}{16} - \frac{5p_2^2}{32} + \frac{p_4}{4}\right)z^4 + \dots$$

Equating the coefficients of  $z$ ,  $z^2$ ,  $z^3$  and  $z^4$  in (28) and on simplifying, we obtain

$$(29) \quad a_2 = \frac{p_1}{16} \dots,$$

$$(30) \quad a_3 = \frac{p_2}{24} - \frac{5p_1^2}{192} \dots,$$

$$(31) \quad a_4 = \frac{1}{4096} [21p_1^3 - 72p_1p_2 + 64p_3] \dots,$$

and

$$(32) \quad a_5 = \frac{1}{80} \left[ -\frac{29}{128}p_1^4 + \frac{17}{16}p_1^2p_2 - \frac{5p_3p_1}{4} - \frac{p_2^2}{2} + p_4 \right] \dots$$

By using Lemma 1, Lemma 3 and Lemma 4 in (29), (30), (31) and (32) and following the procedure of Theorem 2.1, the results (23), (24), (25) and (26) can be easily obtained. □

**THEOREM 3.2.** *If  $f \in \mathcal{K}_s(\mathcal{L})$ , then*

$$(33) \quad |a_3 - \rho a_2^2| \leq \frac{1}{12} \max \left\{ 1, \left| \frac{4+3\rho}{16} \right| \right\} \dots$$

*Proof.* From (29) and (30), we have

$$(34) \quad |a_3 - \rho a_2^2| = \frac{1}{24} \left| p_2 - \left( \frac{20+3\rho}{32} \right) p_1^2 \right| \dots$$

By using (5) in (34), the result (33) is obvious.

For  $\rho = 1$ , (33) transforms to

$$(35) \quad |a_3 - a_2^2| \leq \frac{1}{12} \dots$$

□

**THEOREM 3.3.** *If  $f \in \mathcal{K}_s(\mathcal{L})$ , then*

$$(36) \quad |a_2 a_3 - a_4| \leq \frac{1}{32} \dots$$

*Proof.* By using (29), (30) and (31) and on the lines of Theorem 2.3, the proof is obvious. □

**THEOREM 3.4.** *If  $f \in \mathcal{K}_s(\mathcal{L})$ , then*

$$(37) \quad |a_2 a_4 - a_3^2| \leq \frac{1}{144} \dots$$

*The result is sharp.*

*Proof.* By using (29), (30) and (31) and following the procedure of Theorem 2.4, the result (37) can be easily obtained.

The bound is sharp for  $p_1 = 0$ ,  $p_2 = \pm 2$  and  $p_3 = 0$ . □

**THEOREM 3.5.** *If  $f \in \mathcal{K}_s(\mathcal{L})$ , then*

$$(38) \quad |H_3(1)| \leq \frac{503}{138240} \dots$$

*Proof.* By using (24), (25), (26), (35), (36) and (37) in (1), it yields result (38). □



## References

- [1] R. M. Ali, N. E. Cho, V. Ravichandran and S. Sivaprasad Kumar, *Differential subordination for functions associated with the lemniscate of Bernoulli*, Taiwanese J. Math. **16** (3) (2012), 1017–1026.
- [2] Sahsene Altinkaya and Sibel Yalcin, *Third Hankel determinant for Bazilevic functions*, Adv. Math., Scientific Journal, **5** (2) (2016), 91–96.
- [3] Muhammad Arif, Mohsan Raza, Huo Tang, Shehzad Hussain and Hassan Khan, *Hankel determinant of order three for familiar subsets of analytic functions related with sine function*, Open Math. **17** (2019), 1615–1630.
- [4] K. O. Babalola, *On  $H_3(1)$  Hankel determinant for some classes of univalent functions*, Ineq. Th. Appl. **6** (2010), 1–7.
- [5] L. Bieberbach, *Über die koeffizienten derjenigen Potenzreihen, welche eine schlichte Abbildung des Einheitskreises vermitteln*, Sitzungsberichte Preussische Akademie der Wissenschaften, **138** (1916), 940–955.
- [6] R. N. Das and P. Singh, *On subclasses of schlicht mappings*, Ind. J. Pure Appl. Math. **8** (1977), 864–872.
- [7] L. De-Branges, *A proof of the Bieberbach conjecture*, Acta Math. **154** (1985), 137–152.
- [8] M. Fekete and G. Szegő, *Eine Bemerkung über ungerade schlichte Funktionen*, J. Lond. Math. Soc. **8** (1933), 85–89.
- [9] R. Horn and C. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
- [10] I. Iohvidov, *Hankel and Toeplitz matrices and forms*, Birkhäuser, Boston, Mass, 1982.
- [11] Aini Janteng, Suzeini Abdul Halim and Maslina Darus, *Hankel determinant for starlike and convex functions*, Int. J. Math. Anal. **1** (13) (2007), 619–625.
- [12] S. R. Keogh and E.P. Merkes, *A coefficient inequality for certain subclasses of analytic functions*, Proc. Amer. Math. Soc. **20** (1969), 8–12.
- [13] R. J. Libera and E-J. Zlotkiewicz, *Early coefficients of the inverse of a regular convex function*, Proc. Amer. Math. Soc. **85** (1982), 225–230.
- [14] R. J. Libera and E-J. Zlotkiewicz, *Coefficient bounds for the inverse of a function with derivative in  $\mathcal{P}$* , Proc. Amer. Math. Soc. **87** (1983), 251–257.
- [15] W. Ma, *Generalized Zalcman conjecture for starlike and typically real functions*, J. Math. Anal. Appl. **234** (1999), 328–329.
- [16] B. S. Mehrotra and Gagandeep Singh, *Estimate of second Hankel determinant for certain classes of analytic functions*, Scientia Magna, **8** (3) (2012), 85–94.
- [17] Sh. Najafzadeh, H. Rahmatan and H. Haji, *Application of subordination for estimating the Hankel determinants for subclass of univalent functions*, Creat. Math. Inform. **30** (1) (2021), 69–74.
- [18] J. W. Noonan and D. K. Thomas, *On the second Hankel determinant of a really mean  $p$ -valent functions*, Trans. Amer. Math. Soc. **223** (2) (1976), 337–346.
- [19] Ch. Pommerenke, *Univalent functions*, Math. Lehrbücher, vandenhoek and Ruprecht, Göttingen, 1975.
- [20] V. Ravichandran and S. Verma, *Bound for the fifth coefficient of certain starlike functions*, Comptes Rendus mathématique, **353** (2015), 505–510.
- [21] K. Sakaguchi, *On a certain univalent mapping*, J. Math. Soc. Japan, **11** (1959), 72–80.
- [22] G. Shanmugam, B. Adolf Stephen and K. O. Babalola, *Third Hankel determinant for  $\alpha$  starlike functions*, Gulf J. Math. **2** (2) (2014), 107–113.
- [23] Gagandeep Singh and Gurcharanjit Singh, *On third Hankel determinant for a subclass of analytic functions*, Open Sci. J. Math. Appl. **3** (6) (2015), 172–175.
- [24] Gurmeet Singh, Gagandeep Singh and Gurcharanjit Singh, *Certain subclasses of multivalent functions defined with generalized Salagean operator and related to sigmoid function and lemniscate of Bernoulli*, J. Frac. Calc. Appl. **13** (1) (2022), 65–81.
- [25] J. Sokol and J. Stankiewicz, *Radius of convexity of some subclasses of strongly starlike functions*, Zeszyty Nauk. Politech. Rzeszowskiej Mat. **19** (1996), 101–105.
- [26] J. Sokol and D. K. Thomas, *Further results on a class of starlike functions related to the Bernoulli lemniscate*, Houston J. Math., **44** (1) (2018), 83–95.

- [27] N. Ullah, I. Ali, S. M. Hussain, Jong-Suk Ro, N. Khan and B. Khan, *Third Hankel determinant for a subclass of univalent functions associated with lemniscate of Bernoulli*, *Fractal and Fractional*, **6** (48) (2022), 1–8.

**Gagandeep Singh**

Department of Mathematics, Khalsa College, Amritsar 143002, Punjab, India

*E-mail*: kamboj.gagandeep@yahoo.in

**Gurcharanjit Singh**

Department of Mathematics, G.N.D.U. College, Chungh 143304, Punjab, India

*E-mail*: dhillongs82@yahoo.com