

## A NEW ITERATION METHOD FOR FIXED POINT OF NONEXPANSIVE MAPPING IN UNIFORMLY CONVEX BANACH SPACE

OMPRAKASH SAHU\*, AMITABH BANERJEE, AND NIYATI GURUDWAN

**ABSTRACT.** The aim of this paper is to introduce a new iterative process and show that our iteration scheme is faster than other existing iteration schemes with the help of numerical examples. Next, we have established convergence and stability results for the approximation of fixed points of the contractive-like mapping in the framework of uniformly convex Banach space. In addition, we have established some convergence results for the approximation of the fixed points of a nonexpansive mapping.

This article is the revised version by the corrigendum. Korean J. Math. **31** (2023), No. 1, pp. 121–122.

### 1. Introduction

Fixed point theory always has an important role in the field of analysis. Banach contraction principle become milestone in the fixed point theory always inspire researchers to obtain fixed point of different mappings over different spaces. The origin of fixed point theory lies in the method of successive approximation used for proving existence of solution of differential equation introduced independently by Joseph Liouville in 1837 and Charles Picard in 1890.

In 1953, Mann [27], introduced an iterative scheme. Later in 1974, Ishikawa [22], introduced an iterative scheme which was two step iterative scheme. In 2000, Noor [15], introduced a three-step iterative scheme for approximating fixed point problems. Later several researchers modified Mann, Ishikawa and Noor iterations.

In 2007, Agarwal et al. [20], developed a new iteration method and proved that this iteration process converges faster than Mann iteration for contraction mapping. In 2014, Abbas and Nazir [14] introduced an new iterative scheme which is converges faster than Agarwal et al. [20]. In 2016, Thakur et al. [3] proposed a new iteration method for Suzuki generalized nonexpansive mapping and proved that this iteration process converges faster than previous process. In 2018, Ullah et al. [10] developed new iteration scheme. Let  $M$  be a nonempty closed convex bounded subset of uniformly

---

Received January 3, 2022. Revised November 20, 2022. Accepted November 24, 2022.

2010 Mathematics Subject Classification: 47H09, 47H10.

Key words and phrases: Fixed Point, Uniformly convex Banach space, Contractive like mapping, Nonexpansive mapping, Stability.

\* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2022.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

Banach space  $K$  for each  $u_0 \in M$  construct a sequence  $\{u_n\}$  by

$$(1.1) \quad \begin{cases} u_{n+1} = Tv_n, \\ v_n = T((1 - \alpha_n)w_n + \alpha_n Tw_n) \\ w_n = (1 - \beta_n)u_n + \beta_n Tu_n \end{cases}$$

$\forall n \geq 0$  and  $\{\alpha_n\}$  and  $\{\beta_n\}$  are sequences in  $(0, 1)$ . This iteration process is converges faster than previous iteration process.

In 2020 Hassan et al. [21] developed new iteration process. Let  $M$  be a nonempty closed convex bounded subset of uniformly Banach space  $K$  for each  $u_0 \in M$  construct a sequence  $\{u_n\}$  by

$$(1.2) \quad \begin{cases} u_{n+1} = T((1 - \alpha_n)v_n + \alpha_n Tv_n) \\ v_n = T((1 - \beta_n)w_n + \beta_n Tw_n) \\ w_n = T((1 - \gamma_n)x_n + \gamma_n Tx_n) \\ x_n = T((1 - \zeta_n)u_n + \zeta_n Tu_n) \end{cases}$$

for all  $n \geq 0$  and  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$  and  $\{\zeta_n\}$  are sequences in  $(0, 1)$ . This iteration process is converges faster than previous iteration process.

Our aim is to introduce a new faster iteration process than those mentioned above and to prove the convergence results for nonexpansive mappings in uniformly convex Banach space.

## 2. Preliminaries

In this section, we recall some definitions and results to be used in establishing the main results.

**DEFINITION 2.1.** [9] A Banach space  $K$  is said to be uniformly convex if for each  $\epsilon \in (0, 2]$  there is a  $\delta > 0$  such that  $x, y \in K$

$$\begin{cases} \|x\| \leq 1, \\ \|y\| \leq 1, \\ \|x - y\| \geq \epsilon \end{cases} \Rightarrow \left\| \frac{x + y}{2} \right\| \geq \delta.$$

**DEFINITION 2.2.** [28] A Banach space  $K$  is said to satisfy Opial's property if for each sequence  $\{x_n\}$  in  $K$  converging weakly to  $x \in K$ , we have

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|,$$

for all  $y \in K$  s.t.  $x \neq y$ .

**DEFINITION 2.3.** [4] Let  $K$  be a Banach space and let  $T : K \rightarrow K$  be a self map. The mapping  $T$  is called contractive like mapping if there exist a constant  $\delta \in [0, 1)$  and a strictly increasing and continuous function  $\xi : [0, \infty) \rightarrow [0, \infty)$  with  $\xi(0) = 0$  such that for all  $x, y \in K$ ,

$$(2.1) \quad \|Tx - Ty\| \leq \delta \|x - y\| + \xi(\|x - Tx\|).$$

**DEFINITION 2.4.** Let  $K$  be a Banach space and  $M$  be any nonempty subset of  $K$ . Let  $T : M \rightarrow M$  be said to be nonexpansive if for each  $x, y \in K$

$$\|Tx - Ty\| \leq \|x - y\|.$$

DEFINITION 2.5. [7] A mapping  $T : K \rightarrow K$  is said to satisfy condition  $A$ , if  $\exists$  a non decreasing function  $f : [0, \infty) \rightarrow [0, \infty)$  with  $f(0) = 0$  and  $f(c) > 0$  for all  $c > 0$  s.t.  $\|x - Tx\| \geq f(d(x_n, F(T)))$ , for all  $x \in K$ , where  $d(x, F(T)) = \inf\{\|x - x^*\| : x^* \in F(T)\}$ .

The following definition is about the rate of convergence due to Berinde [26] which is used to verify that our iteration process (3.1) convergence faster than the other existing iteration process.

DEFINITION 2.6. [26] Let  $\{a_n\}_{n=0}^\infty$  and  $\{b_n\}_{n=0}^\infty$  be two sequence of positive numbers such that converge to  $a$  and  $b$  respectively. Assume that there exists

$$\lim_{n \rightarrow \infty} \frac{|a_n - a|}{|b_n - b|} = l.$$

- (i) If  $l = 0$ , then the sequence  $\{a_n\}$  converges faster than the sequence  $\{b_n\}$ .
- (ii) If  $0 < l < \infty$ , then we say that the sequence  $\{a_n\}$  and  $\{b_n\}$  have the same rate of converges.
- (iii) If  $l = \infty$ , then the sequence  $\{b_n\}$  converges faster than sequence  $\{a_n\}$ .

DEFINITION 2.7. [2] Let  $\{t_n\}$  be any arbitrary sequence in  $K$ . Then an iteration procedure  $x_{n+1} = f(T, x_n)$ , converging to fixed point  $p$ , is said to  $T$ -stable, if for  $\epsilon_n = \|t_{n+1} - f(T, t_n)\|$ ,  $\forall n \in N$ , we have  $\lim_{n \rightarrow \infty} \epsilon_n = 0$  if and only if  $\lim_{n \rightarrow \infty} t_n = p$ .

LEMMA 2.8. [26] Suppose that for two fixed point iteration processes  $\{u_n\}$  and  $\{v_n\}$  both converging to the same fixed point  $x^*$ , the error estimates

$$\begin{aligned} \|u_n - x^*\| &\leq a_n \quad n \geq 1, \\ \|v_n - x^*\| &\leq b_n \quad n \geq 1 \end{aligned}$$

are available where  $\{a_n\}$  and  $\{b_n\}$  are two sequences of positive numbers converging to zero. If  $\{a_n\}$  converges faster than  $\{b_n\}$ , then  $\{u_n\}$  converges faster than  $\{v_n\}$  to  $x^*$ .

LEMMA 2.9. [25] If  $\lambda$  is a real number such that  $0 \leq \lambda < 1$  and  $\{\epsilon_n\}$  is the sequence of positive numbers such that

$$\lim_{n \rightarrow \infty} \epsilon_n = 0,$$

then for an sequence of positive numbers  $v_n$  satisfying

$$v_{n+1} \leq \lambda v_n + \epsilon_n, \quad \text{for } n = 1, 2, \dots,$$

we have

$$\lim_{n \rightarrow \infty} v_n = 0.$$

LEMMA 2.10. [8] Let  $K$  be a uniformly convex Banach space and  $0 < p \leq t_n \leq q < 1 \forall n \in N$ . Let  $\{x_n\}$  and  $\{y_n\}$  be two sequences of  $K$  s.t.  $\limsup_{n \rightarrow \infty} \|x_n\| \leq a$ ,  $\limsup_{n \rightarrow \infty} \|y_n\| \leq a$  and  $\limsup_{n \rightarrow \infty} \|t_n x_n + (1 - t_n) y_n\| = a$  hold for some  $a \geq 0$ . Then  $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$ .

LEMMA 2.11. [9] Let  $K$  be a uniformly convex Banach space and  $M$  be any nonempty closed convex subset of  $K$ . Let  $T$  be a nonexpansive mapping on  $K$ . Then,  $I - T$  is demiclosed at zero.

### 3. Main result

**3.1. New Iteration Scheme.** In this section, we introduce a new iteration scheme. Let  $K$  be uniformly convex Banach space and  $\phi \neq M$  be closed and convex subset of  $K$ . Let  $T : M \rightarrow M$  be any nonlinear mapping and for each  $u_0 \in M$  construct the sequence  $\{u_n\}$

$$(3.1) \quad \begin{cases} u_{n+1} = T((1 - \alpha_n)v_n + \alpha_n T v_n) \\ v_n = T((1 - \beta_n)w_n + \beta_n T w_n) \\ w_n = T((1 - \gamma_n)x_n + \gamma_n T x_n) \\ x_n = T((1 - \zeta_n)y_n + \zeta_n T y_n) \\ y_n = T((1 - \eta_n)u_n + \eta_n T u_n) \end{cases}$$

For all  $n \geq 1$  and  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\zeta_n\}$  and  $\{\eta_n\}$  are sequences in  $(0, 1)$ .

**3.2. Convergence and Stability Results for Contractive like Mapping.** In this section we establish convergence and stability results for new iteration process (3.1).

**THEOREM 3.1.** *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  be uniformly convex Banach space. Let  $T : M \rightarrow M$  with satisfying equation (2.1) and  $x^*$  be a fixed point of  $T$ . Suppose that  $\{u_n\}$  generated by (3.1) and  $\sum_{n=0}^{\infty} \alpha_n = \infty$  or  $\sum_{n=0}^{\infty} \beta_n = \infty$  or  $\sum_{n=0}^{\infty} \gamma_n = \infty$  or  $\sum_{n=0}^{\infty} \zeta_n = \infty$  or  $\sum_{n=0}^{\infty} \eta_n = \infty$ . Then  $\{u_n\}$  converges strongly to a unique fixed point of  $T$ .*

*Proof.* Using iteration (3.1) and definition (2.1), we have

$$(3.2) \quad \begin{aligned} \|y_n - x^*\| &= \|(T((1 - \eta_n)u_n + \eta_n T u_n)) - x^*\| \\ &\leq \delta(1 - \eta_n)\|u_n - x^*\| + \delta\eta_n\|T u_n - x^*\| \\ &\leq \delta(1 - \eta_n)\|u_n - x^*\| + \delta^2\eta_n\|u_n - x^*\| \\ &= \delta(1 - (1 - \delta)\eta_n)\|u_n - x^*\|, \end{aligned}$$

$$(3.3) \quad \begin{aligned} \|x_n - x^*\| &= \|(T((1 - \zeta_n)y_n + \zeta_n T y_n)) - x^*\| \\ &\leq \delta(1 - \zeta_n)\|y_n - x^*\| + \delta\zeta_n\|T y_n - x^*\| \\ &\leq \delta(1 - \zeta_n)\|y_n - x^*\| + \delta^2\zeta_n\|y_n - x^*\| \\ &= \delta(1 - (1 - \delta)\zeta_n)\|y_n - x^*\|, \end{aligned}$$

$$(3.4) \quad \begin{aligned} \|w_n - x^*\| &= \|(T((1 - \gamma_n)x_n + \gamma_n T x_n)) - x^*\| \\ &\leq \delta(1 - \gamma_n)\|x_n - x^*\| + \delta\gamma_n\|T x_n - x^*\| \\ &\leq \delta(1 - \gamma_n)\|x_n - x^*\| + \delta^2\gamma_n\|x_n - x^*\| \\ &= \delta(1 - (1 - \delta)\gamma_n)\|x_n - x^*\|, \end{aligned}$$

$$\begin{aligned}
 \|v_n - x^*\| &= \|(T((1 - \beta_n)w_n + \beta_nTw_n)) - x^*\| \\
 &\leq \delta(1 - \beta_n)\|w_n - x^*\| + \delta\beta_n\|Tw_n - x^*\| \\
 &\leq \delta(1 - \beta_n)\|w_n - x^*\| + \delta^2\beta_n\|w_n - x^*\| \\
 &= \delta(1 - (1 - \delta)\beta_n)\|w_n - x^*\|,
 \end{aligned}
 \tag{3.5}$$

$$\begin{aligned}
 \|u_{n+1} - x^*\| &= \|(T((1 - \alpha_n)v_n + \alpha_nTv_n)) - x^*\| \\
 &\leq \delta(1 - \alpha_n)\|v_n - x^*\| + \delta\alpha_n\|Tv_n - x^*\| \\
 &\leq \delta(1 - \alpha_n)\|v_n - x^*\| + \delta^2\alpha_n\|v_n - x^*\| \\
 &= \delta(1 - (1 - \delta)\alpha_n)\|v_n - x^*\|,
 \end{aligned}
 \tag{3.6}$$

From equation (3.2), (3.3), (3.4), (3.5) and (3.6), we obtain

$$\begin{aligned}
 \|u_{n+1} - x^*\| &\leq \delta^5(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)(1 - (1 - \delta)\eta_n)\|u_n - x^*\| \\
 &\leq \delta^{5+5} \prod_{k=n-1}^n (1 - (1 - \delta)\alpha_k)(1 - (1 - \delta)\beta_k)(1 - (1 - \delta)\gamma_k)(1 - (1 - \delta)\zeta_k)(1 - (1 - \delta)\eta_k)\|u_{n-1} - x^*\| \\
 &\vdots \\
 &\leq \delta^{5(n+1)} \prod_{k=0}^n (1 - (1 - \delta)\alpha_k)(1 - (1 - \delta)\beta_k)(1 - (1 - \delta)\gamma_k)(1 - (1 - \delta)\zeta_k)(1 - (1 - \delta)\eta_k)\|u_0 - x^*\|
 \end{aligned}
 \tag{3.7}$$

Since  $\delta \in [0, 1)$  and  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\zeta_n\}$  and  $\{\eta_n\}$  are sequences in  $(0, 1)$ . Using inequality  $1 - z \leq e^{-z} \forall z \in [0, 1]$ , thus From equation (3.7), we have

$$\|u_{n+1} - x^*\| \leq \frac{\delta^{5(n+1)}\|u_0 - x^*\|}{e^{(1-\delta)\sum_{k=0}^{\infty} \alpha_k + \sum_{k=0}^{\infty} \beta_k + \sum_{k=0}^{\infty} \gamma_k + \sum_{k=0}^{\infty} \zeta_k + \sum_{k=0}^{\infty} \eta_k}}
 \tag{3.8}$$

Taking limits on both sides

$$\lim_{n \rightarrow \infty} \|u_{n+1} - x^*\| \leq \lim_{n \rightarrow \infty} \frac{\delta^{5(n+1)}\|u_0 - x^*\|}{e^{(1-\delta)\sum_{k=0}^{\infty} \alpha_k + \sum_{k=0}^{\infty} \beta_k + \sum_{k=0}^{\infty} \gamma_k + \sum_{k=0}^{\infty} \zeta_k + \sum_{k=0}^{\infty} \eta_k}} \leq 0.$$

$\{u_n\}$  is strongly convergent to  $x^*$ . Next we have to show that  $x^*$  is unique. Let  $x^*, x^{**} \in F(T)$ , such that  $x^* \neq x^{**}$ . Now

$$\|x^* - x^{**}\| = \|Tx^* - Tx^{**}\|
 \tag{3.9}$$

using equation (2.1), we have

$$\begin{aligned}
 \|Tx^* - Tx^{**}\| &\leq \delta\|x^* - x^{**}\| + \xi(\|x^* - Tx^*\|) \\
 &\leq \|x^* - x^{**}\|
 \end{aligned}
 \tag{3.10}$$

From equation (3.9) and (3.10), we have

$$\|x^* - x^{**}\| \leq \|x^* - x^{**}\|.$$

Clearly we have that  $\|x^* - x^{**}\| = \|x^* - x^{**}\|$ . Hence  $x^* = x^{**}$  □

**THEOREM 3.2.** *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space. Let  $T : M \rightarrow M$  with satisfying equation (2.1) and  $x^*$  be a fixed point of  $T$ . Suppose that  $\{u_n\}$  is generated by iteration process (3.1). Then iteration process (3.1) is  $T$ -stable.*

*Proof.* Let  $\{p_n\}$  be an arbitrary sequence in  $K$  and the sequence generated by (3.1) is  $u_{n+1} = f(T, u_n)$  converging to a unique fixed point  $x^*$  and  $\epsilon_n = \|p_{n+1} - f(T, p_n)\|$ . We have to show that  $\lim_{n \rightarrow \infty} \epsilon_n = 0$  iff  $\lim_{n \rightarrow \infty} p_n = x^*$ . Let  $\lim_{n \rightarrow \infty} \epsilon_n = 0$  and

$$\begin{aligned} \|p_{n+1} - x^*\| &= \|p_{n+1} - f(T, p_n) + f(T, p_n) - x^*\| \\ &\leq \|p_{n+1} - f(T, p_n)\| + \|f(T, p_n) - x^*\| \\ &= \epsilon_n + \|T((1 - \alpha_n)q_n + \alpha_n Tq_n) - x^*\| \\ &\leq \epsilon_n + \delta(1 - (1 - \delta)\alpha_n)\|q_n - x^*\| \\ &= \epsilon_n + \delta(1 - (1 - \delta)\alpha_n)\|T((1 - \beta_n)r_n + \beta_n Tr_n) - x^*\| \\ &\leq \epsilon_n + \delta^2(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)\|r_n - x^*\| \\ &= \epsilon_n + \delta^2(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)\|T((1 - \gamma_n)s_n + \gamma_n Ts_n) - x^*\| \\ &\leq \epsilon_n + \delta^3(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)\|s_n - x^*\| \\ &= \epsilon_n + \delta^3(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)\|T((1 - \zeta_n)t_n + \zeta_n Tt_n) - x^*\| \\ &\leq \epsilon_n + \delta^4(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)\|t_n - x^*\| \\ &= \epsilon_n + \delta^4(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)\|T((1 - \eta_n)p_n + \eta_n Tp_n) - x^*\| \\ &\leq \epsilon_n + \delta^5(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)(1 - (1 - \delta)\eta_n)\|p_n - x^*\|, \end{aligned}$$

Since  $\delta \in [0, 1)$  and  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\zeta_n\}$  and  $\{\eta_n\}$  are sequences in  $(0, 1)$ ,  $\delta^5(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)(1 - (1 - \delta)\eta_n) \in (0, 1)$ . Hence by Lemma 2.9, we have  $\lim_{n \rightarrow \infty} \|p_n - x^*\| = 0$ , which gives  $\lim_{n \rightarrow \infty} p_n = x^*$ . On the other hand, suppose that  $\lim_{n \rightarrow \infty} p_n = x^*$ . Then,

$$\begin{aligned} \epsilon_n &= \|p_{n+1} - f(T, p_n)\| \\ &\leq \|p_{n+1} - x^*\| + \|f(T, p_n) - x^*\| \\ &= \|p_{n+1} - x^*\| + \|T((1 - \alpha_n)q_n + \alpha_n Tq_n) - x^*\| \\ &\leq \|p_{n+1} - x^*\| + \delta(1 - (1 - \delta)\alpha_n)\|q_n - x^*\| \\ &= \|p_{n+1} - x^*\| + \delta(1 - (1 - \delta)\alpha_n)\|T((1 - \beta_n)r_n + \beta_n Tr_n) - x^*\| \\ &\leq \|p_{n+1} - x^*\| + \delta^2(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)\|r_n - x^*\| \\ &= \|p_{n+1} - x^*\| + \delta^2(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)\|T((1 - \gamma_n)s_n + \gamma_n Ts_n) - x^*\| \\ &\leq \|p_{n+1} - x^*\| + \delta^3(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)\|s_n - x^*\| \\ &= \|p_{n+1} - x^*\| + \delta^3(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)\|T((1 - \zeta_n)t_n + \zeta_n Tt_n) - x^*\| \\ &\leq \|p_{n+1} - x^*\| + \delta^4(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)\|t_n - x^*\| \\ &= \|p_{n+1} - x^*\| + \delta^4(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)\|T((1 - \eta_n)p_n + \eta_n Tp_n) - x^*\| \\ &\leq \|p_{n+1} - x^*\| + \delta^5(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)(1 - (1 - \delta)\eta_n)\|p_n - x^*\|, \end{aligned}$$

Taking limit both sides, we have

$$\lim_{n \rightarrow \infty} \epsilon_n = 0.$$

Hence iteration process (3.1) is  $T$ -stable.  $\square$

**3.3. Comparison Result.** In this section, we comparing the new iteration scheme (3.1) and iteration scheme (1.2) for contractive mappings due to Berinde [26] :

**THEOREM 3.3.** *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space. Let  $T : M \rightarrow M$  with satisfying equation (2.1) and  $x^*$  be a fixed point of  $T$ . Suppose that  $\{u_n\}$  and  $\{r_n\}$  are sequences defined by iteration process (3.1) and (1.2) and  $\sum_{n=0}^{\infty} \eta_n = \infty$ . Then  $\{u_n\}$  converges faster than  $\{r_n\}$ .*

*Proof.* From equation (3.7) in Theorem 3.1, we have

$$(3.11) \quad \|u_{n+1} - x^*\| \leq \delta^{5(n+1)} \prod_{k=0}^n (1 - (1 - \delta)\alpha_k)(1 - (1 - \delta)\beta_k)(1 - (1 - \delta)\gamma_k)(1 - (1 - \delta)\zeta_k)(1 - (1 - \delta)\eta_k) \|u_0 - x^*\|$$

Now using iteration process (1.2) and equation (2.1), we have

$$(3.12) \quad \begin{aligned} \|s_n - x^*\| &= \|T(1 - \zeta_n)r_n + \zeta_n Tr_n - x^*\| \\ &\leq \delta(1 - \zeta_n)\|r_n - x^*\| + \delta\zeta_n\|Tr_n - x^*\| \\ &= \delta(1 - (1 - \delta)\zeta_n)\|r_n - x^*\|, \end{aligned}$$

$$(3.13) \quad \begin{aligned} \|q_n - x^*\| &= \|T((1 - \gamma_n)s_n + \gamma_n Ts_n) - x^*\| \\ &\leq \delta(1 - \gamma_n)\|s_n - x^*\| + \delta\gamma_n\|Ts_n - x^*\| \\ &= \delta(1 - (1 - \delta)\gamma_n)\|s_n - x^*\| \end{aligned}$$

$$(3.14) \quad \begin{aligned} \|p_n - x^*\| &= \|T((1 - \beta_n)q_n + \beta_n Tq_n) - x^*\| \\ &\leq \delta(1 - \beta_n)\|q_n - x^*\| + \delta\beta_n\|Tq_n - x^*\| \\ &= \delta(1 - (1 - \delta)\beta_n)\|q_n - x^*\| \end{aligned}$$

$$(3.15) \quad \begin{aligned} \|r_{n+1} - x^*\| &= \|T((1 - \alpha_n)p_n + \alpha_n Tp_n) - x^*\| \\ &\leq \delta(1 - \alpha_n)\|p_n - x^*\| + \delta\alpha_n\|Tp_n - x^*\| \\ &= \delta(1 - (1 - \delta)\alpha_n)\|p_n - x^*\| \end{aligned}$$

From equation (3.12), (3.13), (3.14) and (3.15), we have

$$\begin{aligned} \|r_{n+1} - x^*\| &\leq \delta^4(1 - (1 - \delta)\alpha_n)(1 - (1 - \delta)\beta_n)(1 - (1 - \delta)\gamma_n)(1 - (1 - \delta)\zeta_n)\|r_n - x^*\| \\ &\quad \vdots \\ &\leq \delta^{4(n+1)} \prod_{k=0}^{\infty} (1 - (1 - \delta)\alpha_k)(1 - (1 - \delta)\beta_k)(1 - (1 - \delta)\gamma_k)(1 - (1 - \delta)\zeta_k) \|r_0 - x^*\| \end{aligned}$$

Now

$$\begin{aligned} &\lim_{n \rightarrow \infty} \frac{\|u_{n+1} - x^*\|}{\|r_{n+1} - x^*\|} \\ &\leq \lim_{n \rightarrow \infty} \frac{\delta^{5(n+1)} \prod_{k=0}^n (1 - (1 - \delta)\alpha_k)(1 - (1 - \delta)\beta_k)(1 - (1 - \delta)\gamma_k)(1 - (1 - \delta)\zeta_k)(1 - (1 - \delta)\eta_k) \|u_0 - x^*\|}{\delta^{4(n+1)} \prod_{k=0}^{\infty} (1 - (1 - \delta)\alpha_k)(1 - (1 - \delta)\beta_k)(1 - (1 - \delta)\gamma_k)(1 - (1 - \delta)\zeta_k) \|r_0 - x^*\|} \\ &\leq \lim_{n \rightarrow \infty} \frac{\delta^{(n+1)} \prod_{k=0}^{\infty} (1 - (1 - \delta)\zeta_k) \|u_0 - x^*\|}{\|r_0 - x^*\|} \end{aligned}$$

Using inequality  $1 - z \leq e^{-z} \forall z \in [0, 1]$ , we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\|u_{n+1} - x^*\|}{\|r_{n+1} - x^*\|} &\leq \lim_{n \rightarrow \infty} \frac{\|u_0 - x^*\| \delta^{(n+1)}}{\|r_0 - x^*\| e^{(1-\delta) \sum_{n=0}^{\infty} \eta_n}} \\ &\leq 0. \end{aligned}$$

Hence by definition 2.6 and lemma 2.8, iteration process (3.1) converges to  $x^*$  faster than iteration process (1.2). □

### 3.4. Numerical Example.

EXAMPLE 3.4. Let  $K = R$  and  $M = [0, 60]$  and  $T : M \rightarrow M$  be a mapping defined by  $T(x) = \sqrt{(x^2 - 6x + 48)}$ , for all  $x \in M$  for  $x_0 = 100$  and  $\alpha_n = \beta_n = \gamma_n = \zeta_n = \eta_n = 2/3, n = 1, 2, 3, \dots$  From Table 1 we can see that all the iteration procedure

are converging to  $x^* = 8$  clearly, our iteration process requires the least number of iteration as compared to other iteration scheme.

The convergence behaviour of these iteration process are represented in the Figure 1.

| It No. | New          | Hassan       | Ullah        | Thakur       | Abass        | Agrwal       | Noor         | Ishikawa     |
|--------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| 0      | 100.00000000 | 100.00000000 | 100.00000000 | 100.00000000 | 100.00000000 | 100.00000000 | 100.00000000 | 100.00000000 |
| 1      | 76.88489427  | 81.46543216  | 90.69448347  | 94.09902562  | 93.89214394  | 95.95936500  | 95.44593195  | 96.89242398  |
| 2      | 54.45329523  | 63.31482922  | 81.46533656  | 88.22654404  | 87.81490958  | 91.93153627  | 90.90869431  | 93.79238794  |
| 3      | 33.39538742  | 45.79841313  | 72.33045956  | 82.38639575  | 81.77259116  | 87.91763715  | 86.38999096  | 90.70039161  |
| 4      | 15.97752996  | 29.48754556  | 63.31455234  | 76.58322320  | 75.77041826  | 83.91894046  | 81.89178870  | 87.61698448  |
| 5      | 8.55219930   | 15.97752996  | 54.45286163  | 70.82270235  | 69.81483834  | 79.93689555  | 77.41637248  | 84.54277243  |
| 6      | 8.01354766   | 9.08714700   | 45.79773758  | 65.11186097  | 63.91391009  | 75.97316133  | 72.96641528  | 81.47842546  |
| 7      | 8.00030721   | 8.06121826   | 37.43070552  | 59.45952389  | 58.07786208  | 72.02964733  | 68.54506760  | 78.42468678  |
| 8      | 8.00000695   | 8.00298149   | 29.48579681  | 53.87694799  | 52.31990248  | 68.10856483  | 64.15607319  | 75.38238351  |
| 9      | 8.00000016   | 8.00014401   | 22.19559369  | 48.37874621  | 46.65741883  | 64.21249133  | 59.80392033  | 72.35243933  |
| 10     | 8.00000000   | 8.00000695   | 15.97253226  | 42.98426007  | 41.11379562  | 60.34445234  | 55.49404180  | 69.33588953  |
| 11     | 8.00000000   | 8.00000034   | 11.45495317  | 37.71963945  | 35.72122535  | 56.50802617  | 51.23308243  | 66.33389899  |
| 12     | 8.00000000   | 8.00000002   | 9.08200038   | 32.62104378  | 30.52511930  | 52.70747968  | 47.02926101  | 63.34778373  |
| 13     | 8.00000000   | 8.00000000   | 8.26820505   | 27.73959064  | 25.59100874  | 48.94794547  | 42.89286528  | 60.37903703  |
| 14     | 8.00000000   | 8.00000000   | 8.06076331   | 23.14881556  | 21.01481733  | 45.23565581  | 38.83693502  | 57.42936099  |
| 15     | 8.00000000   | 8.00000000   | 8.01344474   | 18.95476278  | 16.93536486  | 41.57825411  | 34.87820942  | 54.50070518  |
| 16     | 8.00000000   | 8.00000000   | 8.00295873   | 15.30471974  | 13.53809028  | 37.98521309  | 31.03843705  | 51.59531397  |
| 17     | 8.00000000   | 8.00000000   | 8.00065033   | 12.37662020  | 11.01303065  | 34.46839920  | 27.34615248  | 48.71578499  |
| 18     | 8.00000000   | 8.00000000   | 8.00014291   | 10.30990262  | 9.42537361   | 31.04283503  | 23.83895687  | 45.86514160  |
| 19     | 8.00000000   | 8.00000000   | 8.00003140   | 9.07881940   | 8.60168214   | 27.72772040  | 20.56604605  | 43.04692319  |
| 20     | 8.00000000   | 8.00000000   | 8.00000690   | 8.46084622   | 8.23715727   | 24.54776342  | 17.58985797  | 40.26529812  |
| 21     | 8.00000000   | 8.00000000   | 8.00000152   | 8.18727724   | 8.09048406   | 21.53480474  | 14.98376119  | 37.52520521  |
| 22     | 8.00000000   | 8.00000000   | 8.00000033   | 8.07435771   | 8.03406046   | 18.72948889  | 12.8201608   | 34.83253115  |
| 23     | 8.00000000   | 8.00000000   | 8.00000007   | 8.02923577   | 8.01275409   | 16.18214452  | 11.14469090  | 32.19433187  |
| 24     | 8.00000000   | 8.00000000   | 8.00000002   | 8.01144958   | 8.00476633   | 13.95082100  | 9.94733639   | 29.61910606  |
| 25     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00447699   | 8.00177989   | 12.09285757  | 9.15623528   | 27.11712591  |
| 26     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00174951   | 8.00066448   | 10.64692213  | 8.66634550   | 24.70082122  |
| 27     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00068351   | 8.00024804   | 9.61028643   | 8.37671638   | 22.38519001  |
| 28     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00026701   | 8.00009259   | 8.92895137   | 8.21052211   | 20.18816228  |
| 29     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00010430   | 8.00003456   | 8.51487210   | 8.11685900   | 18.13075417  |
| 30     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00004074   | 8.00001290   | 8.27788669   | 8.06462079   | 16.23670723  |
| 31     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00001592   | 8.00000482   | 8.14759647   | 8.03565798   | 14.53113577  |
| 32     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00000622   | 8.00000180   | 8.07768589   | 8.01965293   | 13.03764143  |
| 33     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00000243   | 8.00000067   | 8.04068727   | 8.01082464   | 11.7731536   |
| 34     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00000095   | 8.00000025   | 8.02125334   | 8.00595995   | 10.74534560  |
| 35     | 8.00000000   | 8.00000000   | 8.00000000   | 8.00000037   | 8.00000009   | 8.01108640   | 8.00328084   | 9.94319427   |

TABLE 1. Comparison table

**3.5. Convergence Results for Nonexpansive Mapping.** In this section we establish some convergence results for nonexpansive mappings;

LEMMA 3.5. *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space. Let  $T : M \rightarrow M$  be a nonexpansive mapping with  $F(T) \neq \phi$ . Suppose that  $\{u_n\}$  is generated by (3.1). Then  $\lim_{n \rightarrow \infty} \|u_n - x^*\|$  exists  $\forall x^* \in F(T)$ .*






FIGURE 1. Comparison Plot

*Proof.* Let  $x^* \in F(T) \forall n \in N$ . Using iteration process (3.1), we have

$$\begin{aligned}
 \|y_n - x^*\| &= \|T((1 - \eta_n)u_n + \eta_n T u_n) - x^*\| \\
 &\leq (1 - \eta_n)\|u_n - x^*\| + \eta_n \|T u_n - x^*\| \\
 &\leq \|u_n - x^*\|,
 \end{aligned}
 \tag{3.16}$$

$$\begin{aligned}
 \|x_n - x^*\| &= \|T((1 - \zeta_n)y_n + \zeta_n T y_n) - x^*\| \\
 &\leq (1 - \zeta_n)\|y_n - x^*\| + \zeta_n \|T y_n - x^*\| \\
 &\leq \|y_n - x^*\| \\
 &\leq \|u_n - x^*\|,
 \end{aligned}
 \tag{3.17}$$

$$\begin{aligned}
 \|w_n - x^*\| &= \|T((1 - \gamma_n)x_n + \gamma_n T x_n) - x^*\| \\
 &\leq (1 - \gamma_n)\|x_n - x^*\| + \gamma_n \|T x_n - x^*\| \\
 &\leq \|x_n - x^*\| \\
 &\leq \|u_n - x^*\|,
 \end{aligned}
 \tag{3.18}$$

$$\begin{aligned}
 \|v_n - x^*\| &= \|T((1 - \beta_n)w_n + \beta_n T w_n) - x^*\| \\
 &\leq (1 - \beta_n)\|w_n - x^*\| + \beta_n \|T w_n - x^*\| \\
 &\leq \|w_n - x^*\| \\
 &\leq \|u_n - x^*\|,
 \end{aligned}
 \tag{3.19}$$

Thus

$$\begin{aligned}
 \|u_{n+1} - x^*\| &= \|T((1 - \alpha_n)v_n + \alpha_nTv_n) - x^*\| \\
 &\leq (1 - \alpha_n)\|v_n - x^*\| + \alpha_n\|Tv_n - x^*\| \\
 &\leq \|v_n - x^*\| \\
 (3.20) \qquad &\leq \|u_n - x^*\|,
 \end{aligned}$$

Hence  $\lim_{n \rightarrow \infty} \|u_n - x^*\| = \text{exists}$  for all  $x^* \in F(T)$ .  $\square$

LEMMA 3.6. *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space. Let  $T : M \rightarrow M$  be a nonexpansive mapping with  $F(T) \neq \phi$ . Suppose  $\{u_n\}$  is generated by (3.1). Then  $\lim_{n \rightarrow \infty} \|Tu_n - u_n\| = 0$ .*

*Proof.* Let  $x^* \in F(T)$  and let  $x^* \in M$ . Then by Lemma 3.5

$$\lim_{n \rightarrow \infty} \|u_n - x^*\| = \text{exists}.$$

Let  $\lim_{n \rightarrow \infty} \|u_n - x^*\| = a$ .

**Case I**  $a = 0$ , we are done.

**Case II**  $a > 0$ . From equation (3.16) in Lemma 3.5, we have

$$\begin{aligned}
 \|y_n - x^*\| &\leq \|u_n - x^*\| \\
 (3.21) \qquad \limsup_{n \rightarrow \infty} \|y_n - x^*\| &\leq \limsup_{n \rightarrow \infty} \|u_n - x^*\| = a
 \end{aligned}$$

Since  $T$  is nonexpansive mapping then  $\|Ty_n - x^*\| \leq \|y_n - x^*\|$ . It follows that

$$(3.22) \qquad \limsup_{n \rightarrow \infty} \|Ty_n - x^*\| \leq a.$$

$$\begin{aligned}
 (3.23) \qquad \|u_{n+1} - x^*\| &= \|T((1 - \alpha_n)v_n + \alpha_nTv_n) - x^*\| \\
 &\leq \|v_n - x^*\| \\
 &= \|T((1 - \beta_n)w_n + \beta_nTw_n) - x^*\| \\
 &\leq \|w_n - x^*\| \\
 &= \|T((1 - \gamma_n)x_n + \gamma_nTx_n) - x^*\| \\
 &\leq \|x_n - x^*\| \\
 &= \|T((1 - \zeta_n)y_n + \zeta_nTy_n) - x^*\| \\
 &\leq (1 - \zeta_n)\|u_n - x^*\| + \zeta_n\|Ty_n - x^*\| \\
 &= \|u_n - x^*\| - \zeta_n\|u_n - x^*\| + \zeta_n\|y_n - x^*\|,
 \end{aligned}$$

This implies that

$$\begin{aligned} \frac{\|u_{n+1} - x^*\| - \|u_n - x^*\|}{\zeta_n} &\leq \|y_n - x^*\| - \|u_n - x^*\| \\ \|u_{n+1} - x^*\| - \|u_n - x^*\| &\leq \frac{\|u_{n+1} - x^*\| - \|u_n - x^*\|}{\zeta_n} \\ &\leq \|y_n - x^*\| - \|u_n - x^*\|, \\ \|u_{n+1} - x^*\| &\leq \|y_n - x^*\| \\ a &\leq \liminf_{n \rightarrow \infty} \|y_n - x^*\| \\ \lim_{n \rightarrow \infty} \|y_n - x^*\| &= a \\ \lim_{n \rightarrow \infty} \|T((1 - \eta_n)u_n + \eta_n Tu_n) - x^*\| &= a \\ \lim_{n \rightarrow \infty} \|(1 - \eta_n)(u_n - x^*) + \eta(Tu_n - x^*)\| &= a \end{aligned}$$

Hence by Lemma 2.10, we have

$$\lim_{n \rightarrow \infty} \|Tu_n - u_n\| = 0.$$

□

**THEOREM 3.7.** *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space with satisfy opial’s property. Let  $T : M \rightarrow M$  be a nonexpansive mapping with  $F(T) \neq \phi$ . Suppose that  $\{u_n\}$  is generated by (3.1). Then  $\{u_n\}$  converges weakly to a fixed point of  $T$ .*

*Proof.* Let  $x^* \in F(T)$ . By Lemma 3.5, then  $\lim_{n \rightarrow \infty} \|u_n - x^*\|$  exists. We prove that  $\{u_n\}$  has a unique weak subsequential limit in  $F(T)$ . Let  $p$  and  $q$  be weak limits of the subsequences  $\{u_m\}$  and  $\{u_k\}$  of  $\{u_n\}$  respectively. From Lemma 3.6, we have  $\lim_{n \rightarrow \infty} \|u_n - Tu_n\| = 0$  and  $I - T$  is demiclosed with respect to zero by Lemma 2.11, we have that  $Tp = p$ . Similar to prove that  $q \in F(T)$ . Next we have to show that uniqueness. From Lemma 3.5, we have  $\lim_{n \rightarrow \infty} \|u_n - q\|$  exist. Now suppose that  $p \neq q$ , then by opial’s condition,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n - p\| &= \lim_{m \rightarrow \infty} \|u_m - p\| \\ &< \lim_{m \rightarrow \infty} \|u_m - q\| \\ &= \lim_{n \rightarrow \infty} \|u_n - q\| \\ &= \lim_{k \rightarrow \infty} \|u_k - q\| \\ &\leq \lim_{k \rightarrow \infty} \|u_k - p\| \\ &= \lim_{n \rightarrow \infty} \|u_n - p\|. \end{aligned}$$

which is contradiction, so  $p = q$ . Hence  $\{u_n\}$  converges weakly to a fixed point of  $F(T)$ . □

**THEOREM 3.8.** *Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space. Let  $T : M \rightarrow M$  be a nonexpansive mapping with  $F(T) \neq \phi$ . Suppose that  $\{u_n\}$  is generated by (3.1). Then the sequence  $\{u_n\}$  converges to a fixed*

point of  $T$  iff  $\liminf_{n \rightarrow \infty} d(u_n, F(T)) = 0$ , where  $d(u, F(T)) = \inf\{\|u - x^*\| : x^* \in F(T)\}$ .

*Proof.* Let  $\{u_n\}$  converges to  $x^*$ , then  $\lim_{n \rightarrow \infty} d(u_n, x^*) = 0$ . It follows that  $\lim_{n \rightarrow \infty} d(u_n, F(T)) = 0$ . Therefore,  $\liminf_{n \rightarrow \infty} d(u_n, F(T)) = 0$ .

**Conversely:** Suppose that  $\lim_{n \rightarrow \infty} d(u_n, F(T)) = 0$ . It follows from Lemma 3.3, that  $\lim_{n \rightarrow \infty} \|u_n - x^*\|$  exists and that  $\liminf_{n \rightarrow \infty} d(u_n, F(T))$  exists for all  $x^* \in F(T)$ . But by our assumption  $\liminf_{n \rightarrow \infty} d(u_n, F(T)) = 0$ , therefore we have  $\lim_{n \rightarrow \infty} d(u_n, F(T)) = 0$ . We will show that  $\{u_n\}$  is a Cauchy sequence in  $M$ . Since  $\lim_{n \rightarrow \infty} d(u_n, F(T)) = 0$  for given  $\epsilon > 0$ ,  $\exists, n_0 \in N$  s.t.  $\forall n \geq n_0$ ,

$$d(u_n, F(T)) < \frac{\epsilon}{2}.$$

In particular,  $\inf\{\|u_{n_0} - x^*\| : x^* \in F(T)\} < \frac{\epsilon}{2}$ .

Hence  $\exists x_1^* \in F(T)$  s.t.  $\|u_{n_0} - x_1^*\| < \frac{\epsilon}{2}$ .

Now for  $m, n \geq n_0$ ,

$$\begin{aligned} \|u_{n+m} - u_n\| &\leq \|u_{n+m} - x_1^*\| + \|u_n - x_1^*\| \\ &\leq 2\|u_{n_0} - x_1^*\| \\ &< \epsilon. \end{aligned}$$

Hence  $\{u_n\}$  is a Cauchy sequence in  $M$ . Since  $M$  is closed in a uniformly Banach space  $K$ , so that  $\exists$  a point  $x^* \in M$  s.t.  $\lim_{n \rightarrow \infty} u_n = x^*$ . Now  $\lim_{n \rightarrow \infty} d(u_n, F(T)) = 0$ , gives that  $d(x^*, F(T)) = 0$ , since  $F(T)$  is closed,  $x^* \in F(T)$ .  $\square$

**THEOREM 3.9.** Let  $\phi \neq M \subset K$  be closed and convex, where  $K$  is a uniformly convex Banach space. Let  $T : M \rightarrow M$  be a nonexpansive mapping with  $F(T) \neq \phi$ . Suppose that  $\{u_n\}$  is generated by (3.1). Let  $T$  satisfy condition A, then  $\{u_n\}$  converges strongly to a fixed point of  $T$ .

*Proof.* By using Lemma 3.6, we have

$$(3.24) \quad \lim_{n \rightarrow \infty} \|u_n - Tu_n\| = 0.$$

From condition A and equation (3.24), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} f(d(u_n, F(T))) &\leq \lim_{n \rightarrow \infty} \|u_n - Tu_n\| \\ &\Rightarrow \lim_{n \rightarrow \infty} f(d(u_n, F(T))) = 0. \end{aligned}$$

Since  $f : [0, \infty) \rightarrow [0, \infty)$  is a non decreasing function satisfying  $f(0) = 0$  and  $f(c) > 0$  for all  $c \in (0, \infty)$ , therefore, we have

$$\lim_{n \rightarrow \infty} d(u_n, F(T)) = 0.$$

By Theorem 3.8, the sequence  $\{u_n\}$  strongly converges to a fixed point of  $F(T)$ .  $\square$

#### 4. Conclusion

The conclusion of our work as follows:

- We have introduced a new iteration process and proved it to be faster than other existing processes in this paper with the help of numerical examples.

- We have discussed the convergence and stability results for the approximation of fixed points of the contractive-like mapping.
- We have established some convergence results for the approximation of fixed points of a nonexpansive mapping.

## References

- [1] A.G.Sanatee, L.Rathour, V.N.Mishra and V.Dewangan, *Some fixed point theorems in regular modular metric spaces and application to Caratheodory's type anti-periodic boundary value problem*, The Journal of Analysis (2022), 1–14, DOI: <https://doi.org/10.1007/s41478-022-00469-z>.
- [2] A.M.Harder, *Fixed point theory and stability results for fixed point iteration procedures*. PhD thesis, University of Missouri-Rolla, Missouri, MO, USA, 1987.
- [3] B.S.Thakur, D.Thakur and M.Postolache, *A new iterative scheme for numerical reckoning fixed points of suzuki's generalized nonexpansive mappings*, App. Math. Comp. **275** (2016), 147–155.
- [4] C.O.Imoru and M.O.Olantiwo, *On the stability of Picard and Mann iteration process*, Carpath. J. Math. **19** (2) (2003), 155–160.
- [5] D.P.Shukla, S.K.Tiwari and A.Gautam, *Common fixed point theorems for two steps iterative scheme using nonexpansive mappings*, Ultra scientist **22** (3) (2010), 867–870.
- [6] D.P.Shukla and S.K.Tiwari, *Common fixed point theorems for three steps iterative scheme using nonexpansive mappings*, Napier Indian Advanced Research Journal of Science **8** (2012), 5–8.
- [7] H.F.Senter and W.G.Dotson, *Approximating fixed points of nonexpansive mappings*, Proc. Amer. Math. Soc. **44** (2) (1974), 375–380.
- [8] J.Schu, *Weak and strong convergence to fixed points of asymptotically nonexpansive mappings*, Bull. Austral. Math. Soc. **43** (1991), 153–159.
- [9] K.Goebel and W.A.Kirk, *Topic in metric fixed point theory*, Cambridge University Press, 1990.
- [10] K.Ullah and M.Arshad, *Numerical reckoning fixed points for Suzuki generalized nonexpansive mappings via new iteration process*, Filomat **32** (1) (2018), 187–196.
- [11] L.N.Mishra, S.K.Tiwari and V.N.Mishra, *Fixed point theorems for generalized weakly S-contractive mappings in partial metric spaces*, J. Appl. Anal. Comput. **5** (5) (2015), 600–612.
- [12] L.N.Mishra, S.K.Tiwari, V.N.Mishra and I.A.Khan, *Unique fixed point theorems for generalized contractive mappings in partial metric spaces*, J. Funct. Spaces 2015, Art. ID. 960827, 8pp.
- [13] L.N.Mishra, V.Dewangan, V.N.Mishra and H. Amrulloh, *Coupled best proximity point theorems for mixed g-monotone mappings in partially ordered metric spaces*, J. Math. Comput. Sci. **11** (5) (2021), 6168–6192.
- [14] M.Abbas, T.Nazir, *A new faster iteration process applied to constrained minimation and feasibility problems*, Matematicki Vesnik, vol. **66** (2) (2014), 223–234.
- [15] M.A.Noor, *New approximation schemes for general variational inequalities*, J. Math. Anal. Appl. **251** (2000), 217–229.
- [16] N.Sharma, L.N.Mishra, V.N.Mishra and H.Almusawa, *Endpoint approximation of standard three-step multi-valued iteration algorithm for nonexpansive mappings*, Appl. Math. Inf. Sci. **15** (1) (2021), 73–81.
- [17] N.Sharma, L.N.Mishra, V.N.Mishra and S.Pandey, *Solution of Delay Differential equation via  $N_1^v$  iteration algorithm*, European J. Pure Appl. Math. **13** (5) (2020), 1110–1130.
- [18] N.Sharma, L.N.Mishra, S.N.Mishra and V.N.Mishra, *Empirical study of new iterative algorithm for generalized nonexpansive operators*, J Math Comput Sci. **25** (3) (2022), 284–295.
- [19] P.Shahi, L.Rathour and V.N.Mishra, *Expansive Fixed Point Theorems for tri-simulation functions*, J. Eng. Exact Sci. –jCEC, **8** (3) (2022), 14303–01e.
- [20] R.P.Agarwal, D.O'Regan and D.R.Sahu, *Iterative construction of fixed points of nearly asymptotically nonexpansive mappings*, J. Nonlinear Convex Anal. **8** (1) (2007), 61–79.
- [21] S.Hassan, M.De la Sen, P.Agrawal, Q.Ali and A. Hussain, *A new faster iterative scheme for numerical fixed points estimation of suzuki's generalized non-expansive mappings*, Math. Probl. Eng. 2020, Art. ID 3863819, 9pp.
- [22] S.Ishikawa, *Fixed points by new iteration method*, Proc. Amer. Math. Soc. **149** (1974), 147–150.
- [23] S.B.Nadlor, *Multivalued contraction mappings*, Pac. J. Math. **30** (1969), 475–488.

- [24] S.K.Tiwari, L.Rathour and L.N.Mishra, *Existence of fixed point theorems for complex partial b-metric spaces using S-contractive mapping*, J. linear topol. algebr. **11** (3) (2022), 177–188.
- [25] V.Berinde, *On the stability of some fixed point procedure*, Bul. Stiint. Univ. Baia Mare Ser. B Fasc. Mat. Inform. **XVIII** (1) (2002), 7–14.
- [26] V.Berinde, *Picard iteration converges faster than Mann iteration for a class of quasicontractive operators*, fixed Point Theory Appl. **2** (2004), 97–105.
- [27] W.R.Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. **4** (1953), 506–510.
- [28] Z.Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.

**Omprakash Sahu**

Department of Mathematics,  
Babu Pandhri Rao Kridatt Govt.College Silouti,Dhamtari,  
Raipur, India  
*E-mail*: om2261995@yahoo.com

**Amitabh Banerjee**

Department of Mathematics,  
Govt. J. Y. Chhattishgarh College Raipur, India,  
Raipur, India  
*E-mail*: amitabh\_61@yahoo.com

**Niyati Gurudwan**

Department of Mathematics,  
Govt. J. Y. Chhattishgarh College Raipur, India,  
Raipur, India  
*E-mail*: niyati.kuhu@gmail.com