

## APPLICATION OF GEGENBAUER POLYNOMIALS TO CERTAIN CLASSES OF BI-UNIVALENT FUNCTIONS OF ORDER $\nu + i\varsigma$

OMAR ALNAJAR\*, ALA AMOURAH, AND MASLINA DARUS

ABSTRACT. In this paper, a new class of bi-univalent functions that are described by Gegenbauer polynomials is presented. We obtain the estimates of the Taylor-Maclaurin coefficients  $|m_2|$  and  $|m_3|$  for each function in this class of bi-univalent functions. In addition, the Fekete-Szegő problems function new are also studied.

### 1. preliminaries

Legendre presented polynomials in his study in 1784 [6]. They are used extensively in analysis to solve boundary value problems like the Laplace equation and the Schrodinger equation by use of polynomial interpolation and approximation. They also have many further applications.

Legendre polynomials have several applications in physics, and are of them is the investigation of spherical harmonics and the construction of gravitational potentials.

Later in the 19th century, a scientist named Leopold Gegenbauer [9], who was of German and Czech ancestry, suggested a fresh set of orthogonal polynomials. is a nice example of a polynomial that has orthogonal qualities in its features. These orthogonal polynomials have a weight function that is more general and are specified on the interval  $[-1, 1]$ . The Fekete-Szegő inequality is a well-known inequality in mathematics that has applications to holomorphic one-to-one functions, complex analysis, and univalent (or holomorphic) functions. The inequality bears the names of the two mathematicians who independently proved it in the early 20th century, Michael Fekete and George-Szegő [21]. In 1967, Lewin [22] discovered that  $|m_2| \leq 1.51$  while researching the bi-univalent function class  $\Sigma$ .

Many areas of mathematics, like complex analysis and geometry, use a class of analytical functions called "bi-univalent functions," which are defined in the complex plane. The bi-univalent functions are a kind of extension of the univalent functions.

---

Received November 3, 2023. Revised December 18, 2023. Accepted March 5, 2024.

2010 Mathematics Subject Classification: 30C45, 30C50.

Key words and phrases: bi-univalent functions, Holomorphic functions, Gegenbauer polynomials, Fekete and Szegő problem.

\* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2024.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

They are responsible for mapping the unit disk of the complex plane to a certain area.

Continued study of bi-univalent functions has yielded several significant insights and advances in complex analysis and geometry [5, 20, 27]. In general

Let  $\mathcal{A}$  be the class of functions  $f$  of the form

$$(1) \quad f(\mathfrak{Z}) = \mathfrak{Z} + m_2\mathfrak{Z}^2 \dots = \mathfrak{Z} + \sum_{k=2}^{\infty} m_k\mathfrak{Z}^k, \quad (\mathfrak{Z} \in \mathfrak{h}).$$

which are analytical on the disk  $\mathfrak{h} = \{\mathfrak{Z} \in \mathbb{C} : |\mathfrak{Z}| < 1\}$ . We also call  $\mathcal{S}$  the subclass of  $\mathcal{A}$  made up of functions of Eq. (1) that are also univalent in  $\mathfrak{h}$ .

Differential subordination of analytical functions gives the field of geometric function theory many powerful tools that can be very helpful. Porwal and Darus presented a paper on the bi-univalent [25]. Additionally, see [10], which carries out analytical operations on the disk denoted by  $\mathfrak{h} = \{\mathfrak{Z} \in \mathbb{C} : |\mathfrak{Z}| < 1\}$ .  $\mathcal{S}$  is also the name given to the subclass of  $\mathcal{A}$  that is composed of functions of Eq (1) that are also univalent in  $\mathfrak{h}$ .

Every mathematical function  $f \in \mathcal{S}$  has an inverse, denoted by  $f^{-1}$ .

$$f^{-1}(f(\mathfrak{Z})) = \mathfrak{Z} \quad (\mathfrak{Z} \in \mathfrak{h})$$

and

$$\mathfrak{N} = f(f^{-1}(\mathfrak{N})) \quad (|\mathfrak{N}| < r_0(f); r_0(f) \geq 0.25)$$

where

$$(2) \quad g(\mathfrak{N}) = f^{-1}(\mathfrak{N}) = \mathfrak{N} - m_2\mathfrak{N}^2 + (-m_3 + 2m_2^2)\mathfrak{N}^3 - (m_4 + 5m_2^3 - 5m_3m_2)\mathfrak{N}^4 + \dots$$

A function is said to be bi-univalent in  $\mathfrak{h}$  if both  $f(\mathfrak{Z})$  and  $f^{-1}(\mathfrak{Z})$  are univalent in  $\mathfrak{h}$ . This is because both of these functions evaluate to the same value.

Let  $\Sigma$  stand for the class of bi-univalent in  $\mathfrak{h}$  given by (1). The following are some examples from the class  $\Sigma$ .

$$\frac{\mathfrak{Z}}{1 - \mathfrak{Z}}, \quad \log \frac{1}{1 - \mathfrak{Z}}.$$

On the other hand, the well-known Koebe function is not included in  $\Sigma$ . The following are some additional examples of standard functions found in  $\mathfrak{h}$ :

$$\frac{2\mathfrak{Z} - \mathfrak{Z}^2}{2} \text{ and } \frac{\mathfrak{Z}}{1 - \mathfrak{Z}^2},$$

do not belong to  $\Sigma$  either. See reference [17], for interesting subclasses of functions in class  $\Sigma$ . Following Srivastava et al. [16], foundational work, several new subclasses of the bi-univalent function class  $\Sigma$  were established, was a catalyst for the creation of several unique subclasses inside the bi-univalent function class and estimations for

the initial two coefficients,  $|m_2|$  and  $|m_3|$  in the Taylor-Maclaurin series expansion (1), which were obtained in [8, 12–15, 17–19, 26, 28].

Amourah et al. [4] The following Gegenbauer polynomial generating function was the subject of a study that was carried out in the year 2020.

$$(3) \quad H_\alpha(\gamma, \mathfrak{S}) = \frac{1}{(1 - 2\gamma\mathfrak{S} + \mathfrak{S}^2)^\alpha},$$

where  $\gamma \in [-1, 1]$  and  $\mathfrak{S} \in \mathfrak{h}$  play a role. When  $\gamma$  is held constant, the function  $H_\alpha$  is analytic in  $\mathfrak{h}$ , making its expansion through the Taylor series straightforward, as demonstrated by the following example.

$$(4) \quad H_\alpha(\gamma, \mathfrak{S}) = \sum_{k=0}^{\infty} C_k^\alpha(\gamma)\mathfrak{S}^k,$$

where  $C_k^\alpha(\gamma)$  stands for a polynomial of degree  $k$  that belongs to the Gegenbauer family.

When  $\alpha = 0$ , it is very clear that  $H_\alpha$  does not achieve anything. Because of this, the generating function of the Gegenbauer polynomial is defined as follows:

$$(5) \quad H_0(\gamma, \mathfrak{S}) = 1 - \log(1 - 2\gamma\mathfrak{S} + \mathfrak{S}^2) = \sum_{k=0}^{\infty} C_k^0(\gamma)\mathfrak{S}^k,$$

for  $\alpha = 0$ . Furthermore, it should be emphasized that, as stated in [23], it is preferred for the normalization to be greater than  $-0.5$ . The following recurrence relations are also applicable to the definition of Gegenbauer polynomials.

$$(6) \quad C_k^\alpha(\gamma) = \frac{1}{k} [2\gamma(k + \alpha - 1)C_{k-1}^\alpha(\gamma) - (k + 2\alpha - 2)C_{k-2}^\alpha(\gamma)],$$

keeping initial estimates in mind

$$(7) \quad C_0^\alpha(\gamma) = 1, C_1^\alpha(\gamma) = 2\alpha\gamma \text{ and } C_2^\alpha(\gamma) = 2\alpha(1 + \alpha)\gamma^2 - \alpha.$$

Chebyshev polynomials and Legendre polynomials are special cases of Gegenbauer polynomials  $C_k^\alpha(\gamma)$ . When  $\alpha = 1$ , and when  $\alpha = 1/2$ , Chebyshev polynomials and Legendre polynomials are special cases of Gegenbauer polynomials  $C_k^\alpha(\gamma)$ .

In recent years, a significant number of researchers have focused their attention on researching connections that exist between univalent functions, orthogonal polynomials, and bi-univalent functions, see references [1–3, 7, 8, 11, 24].

Before determining bounds for the  $|m_2|$  and  $|m_3|$  Taylor-Maclaurin coefficients, we first define and then construct a new subclass that is related to Gegenbauer polynomials. The Fekete-Szego functional difficulties are also addressed for functions that belong to this new class. which, part, are formed by Amourah et al. [4]

## 2. Definition and examples

In this section, the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, \beta, \nu, \varsigma, \gamma)$  is defined as follows:

DEFINITION 2.1. For  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$  and  $0 \leq \beta \leq 1$ , a function  $f \in \Sigma$  is said to be in the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, \beta, \nu, \varsigma, \gamma)$  if it satisfies the subordinations:

$$\left( \frac{\mathfrak{I}f'(\mathfrak{I})}{f(\mathfrak{I})} \right)^{\nu+i\varsigma} \left[ (1-\beta) \frac{\mathfrak{I}f'(\mathfrak{I})}{f(\mathfrak{I})} + \beta \left( 1 + \frac{\mathfrak{I}f''(\mathfrak{I})}{f'(\mathfrak{I})} \right) \right]^{\sigma} \prec H_{\alpha}(\gamma, \mathfrak{I})$$

and

$$\left( \frac{\aleph g'(\aleph)}{g(\aleph)} \right)^{\nu+i\varsigma} \left[ (1-\beta) \frac{\aleph g'(\aleph)}{g(\aleph)} + \beta \left( 1 + \frac{\aleph g''(\aleph)}{g'(\aleph)} \right) \right]^{\sigma} \prec H_{\alpha}(\gamma, \aleph).$$

The function  $g(\aleph) = f^{-1}(\aleph)$  is defined by (2) where  $\gamma \in (\frac{1}{2}, 1]$  and  $H_{\alpha}$  is the generating function of the Gegenbauer polynomials given by (3).

Special cases:

i) Assume that  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$  and  $\beta = 0$ , a function  $f \in \Sigma$  is said to be in the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, 0, \nu, \varsigma, \gamma)$  if it satisfies the subordinations:

$$\left( \frac{\mathfrak{I}f'(\mathfrak{I})}{f(\mathfrak{I})} \right)^{\nu+i\varsigma} \left[ \frac{\mathfrak{I}f'(\mathfrak{I})}{f(\mathfrak{I})} \right]^{\sigma} \prec H_{\alpha}(\gamma, \mathfrak{I})$$

and

$$\left( \frac{\aleph g'(\aleph)}{g(\aleph)} \right)^{\nu+i\varsigma} \left[ \frac{\aleph g'(\aleph)}{g(\aleph)} \right]^{\sigma} \prec H_{\alpha}(\gamma, \aleph),$$

where  $\gamma \in (\frac{1}{2}, 1]$ .

ii) Assume that For  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$  and  $\beta = 1$ , a function  $f \in \Sigma$  is said to be in the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, 1, \nu, \varsigma, \gamma)$  if it satisfies the subordinations:

$$\left( \frac{\mathfrak{I}f'(\mathfrak{I})}{f(\mathfrak{I})} \right)^{\nu+i\varsigma} \left[ 1 + \frac{\mathfrak{I}f''(\mathfrak{I})}{f'(\mathfrak{I})} \right]^{\sigma} \prec H_{\alpha}(\gamma, \mathfrak{I})$$

and

$$\left( \frac{\aleph g'(\aleph)}{g(\aleph)} \right)^{\nu+i\varsigma} \left[ 1 + \frac{\aleph g''(\aleph)}{g'(\aleph)} \right]^{\sigma} \prec H_{\alpha}(\gamma, \aleph),$$

where  $\gamma \in (\frac{1}{2}, 1]$ .

## 3. Coefficient bounds of the class $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, \beta, \nu, \varsigma, \gamma)$

Here we gave the upper bound for  $|m_2|$  and  $|m_3|$  of the class  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, \beta, \nu, \varsigma, \gamma)$  provided in Definition 2.1 as follows.

THEOREM 3.1. For  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$  and  $0 \leq \beta \leq 1$ , let  $f \in \mathcal{A}$  be in the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, \beta, \nu, \varsigma, \gamma)$ . Then

$$|m_2| \leq \frac{2|\alpha\gamma| \sqrt{|\alpha\gamma|}}{\sqrt{\left| 2 \left[ \begin{array}{c} \alpha((\nu+i\varsigma)(\nu+i\varsigma+1) + \Upsilon(\sigma, \beta, \nu, \varsigma)) \\ - (1+\alpha)(\nu+i\varsigma+\sigma(\beta+1))^2 \end{array} \right] \gamma^2 + (\nu+i\varsigma+\sigma(\beta+1))^2 \right|}},$$

and

$$|m_3| \leq \frac{2|\alpha\gamma|}{2\sqrt{(\nu + \sigma(2\beta + 1))^2 + \varsigma^2}} + \frac{4\alpha^2\gamma^2}{[(\nu + \sigma(\beta + 1))^2 + \varsigma^2]},$$

where

$$(8) \quad \Upsilon(\sigma, \beta, \nu, \varsigma) = \sigma(\beta + 1) [2(\nu + i\varsigma + 1) + (\sigma - 1)(\beta + 1)].$$

*Proof.* Let  $f \in \mathfrak{G}_\Sigma^\alpha(\sigma, \beta, \nu, \varsigma, \gamma)$ . Then there are two holomorphic functions  $\phi, \varphi$  given by

$$(9) \quad \phi(\mathfrak{S}) = c_1\mathfrak{S} + c_2\mathfrak{S}^2 + c_3\mathfrak{S}^3 + \dots \quad (\mathfrak{S} \in \mathfrak{h})$$

and

$$(10) \quad \varphi(\aleph) = d_1\aleph + d_2\aleph^2 + d_3\aleph^3 + \dots \quad (\aleph \in \mathfrak{h}),$$

with  $\phi(0) = \varphi(0) = 0, |\phi(\mathfrak{S})| < 1, |\varphi(\aleph)| < 1, \mathfrak{S}, \aleph \in \mathfrak{h}$  such that

$$(11) \quad \left(\frac{\mathfrak{S}f'(\mathfrak{S})}{f(\mathfrak{S})}\right)^{\nu+i\varsigma} \left[ (1-\beta)\frac{\mathfrak{S}f'(\mathfrak{S})}{f(\mathfrak{S})} + \beta \left(1 + \frac{\mathfrak{S}f''(\mathfrak{S})}{f'(\mathfrak{S})}\right) \right]^\sigma = H_\alpha(\gamma, \phi(\mathfrak{S}))$$

and

$$(12) \quad \left(\frac{\aleph g'(\aleph)}{g(\aleph)}\right)^{\nu+i\varsigma} \left[ (1-\beta)\frac{\aleph g'(\aleph)}{g(\aleph)} + \beta \left(1 + \frac{\aleph g''(\aleph)}{g'(\aleph)}\right) \right]^\sigma = H_\alpha(\gamma, \varphi(\aleph)).$$

Combining (9), (10) and (11) yields

$$(13) \quad \left(\frac{\mathfrak{S}f'(\mathfrak{S})}{f(\mathfrak{S})}\right)^{\nu+i\varsigma} \left[ (1-\beta)\frac{\mathfrak{S}f'(\mathfrak{S})}{f(\mathfrak{S})} + \beta \left(1 + \frac{\mathfrak{S}f''(\mathfrak{S})}{f'(\mathfrak{S})}\right) \right]^\sigma = 1 + C_1^\alpha(\gamma)c_1\mathfrak{S} + [C_1^\alpha(\gamma)c_2 + C_2^\alpha(\gamma)c_1^2]\mathfrak{S}^2 + \dots$$

and

$$(14) \quad \left(\frac{\aleph g'(\aleph)}{g(\aleph)}\right)^{\nu+i\varsigma} \left[ (1-\beta)\frac{\aleph g'(\aleph)}{g(\aleph)} + \beta \left(1 + \frac{\aleph g''(\aleph)}{g'(\aleph)}\right) \right]^\sigma = 1 + C_1^\alpha(\gamma)d_1\aleph + [C_1^\alpha(\gamma)d_2 + C_2^\alpha(\gamma)d_1^2]\aleph^2 + \dots$$

It is quite well-known that if  $|\phi(\mathfrak{S})| < 1$  and  $|\varphi(\aleph)| < 1, \mathfrak{S}, \aleph \in \sigma$ , then

$$(15) \quad |c_i| \leq 1 \quad \text{and} \quad |d_i| \leq 1 \quad \text{for all } i \in \mathbb{N}.$$

Comparing the corresponding coefficients in (13), after simplifying, we have

$$(16) \quad (\nu + i\varsigma + \sigma(\beta + 1)) m_2 = C_1^\alpha(\gamma)c_1,$$

$$(17) \quad \begin{aligned} & 2(\nu + i\varsigma + \sigma(2\beta + 1)) m_3 \\ & + \frac{1}{2} \left[ (\nu + i\varsigma)(\nu + i\varsigma - 1) + \sigma(\beta + 1) \left( \frac{2(\nu + i\varsigma)}{+(\sigma - 1)(\beta + 1)} \right) - 2(\nu + i\varsigma + \sigma(3\beta + 1)) \right] m_2^2 \\ & = C_1^\alpha(\gamma)c_2 + C_2^\alpha(\gamma)c_1^2, \end{aligned}$$

Comparing the corresponding coefficients in (14), after simplifying, we have

$$(18) \quad -(\nu + i\varsigma + \sigma(\beta + 1)) m_2 = C_1^\alpha(\gamma)d_1,$$

and

$$\begin{aligned}
 & 2(\nu + i\varsigma + \sigma(2\beta + 1))(2m_2^2 - m_3) \\
 & + \frac{1}{2} \left[ (\nu + i\varsigma)(\nu + i\varsigma - 1) + \sigma(\beta + 1) \left( \frac{2(\nu + i\varsigma)}{+(\sigma - 1)(\beta + 1)} \right) - 2(\nu + i\varsigma + \sigma(3\beta + 1)) \right] m_2^2 \\
 (19) \quad & = C_1^\alpha(\gamma)d_2 + C_2^\alpha(\gamma)d_1^2.
 \end{aligned}$$

It follows from (16) and (18) that

$$(20) \quad c_1 = -d_1$$

and

$$(21) \quad 2(\nu + i\varsigma + \sigma(\beta + 1))^2 m_2^2 = [C_1^\alpha(\gamma)]^2 (c_1^2 + d_1^2).$$

If we add (17) to (19), we find that

$$\begin{aligned}
 (22) \quad & [(\nu + i\varsigma)(\nu + i\varsigma + 1) + \sigma(\beta + 1)(2(\nu + i\varsigma + 1) + (\sigma - 1)(\beta + 1))] m_2^2 \\
 & = C_1^\alpha(\gamma)(c_2 + d_2) + C_2^\alpha(\gamma)(c_1^2 + d_1^2).
 \end{aligned}$$

Substituting the value of  $c_1^2 + d_1^2$  from (21) in the right hand side of (22), we deduce that

$$(23) \quad m_2^2 = \frac{[C_1^\alpha(\gamma)]^3 (c_2 + d_2)}{[C_1^\alpha(\gamma)]^2 ((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, \beta, \nu, \varsigma)) - 2C_2^\alpha(\gamma)(\nu + i\varsigma + \sigma(\beta + 1))^2}.$$

where  $\Upsilon(\sigma, \beta, \nu, \varsigma)$  is given by (8).

Further computations using (7), (15) and (23), we obtain

$$|m_2| \leq \frac{2|\alpha\gamma| \sqrt{|\alpha\gamma|}}{\sqrt{\left| 2 \left[ \begin{array}{c} \alpha((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, \beta, \nu, \varsigma)) \\ -(1 + \alpha)(\nu + i\varsigma + \sigma(\beta + 1))^2 \end{array} \right] \gamma^2 + (\nu + i\varsigma + \sigma(\beta + 1))^2 \right|}}.$$

Next, if we subtract (19) from (17), we can easily see that

$$(24) \quad 4(\nu + i\varsigma + \sigma(2\beta + 1))(m_3 - m_2^2) = C_1^\alpha(\gamma)(c_2 - d_2) + C_2^\alpha(\gamma)(c_1^2 - d_1^2).$$

In view of (20) and (21), we get from (24)

$$m_3 = \frac{C_1^\alpha(\gamma)(c_2 - d_2)}{4(\nu + i\varsigma + \sigma(2\beta + 1))} + \frac{[C_1^\alpha(\gamma)]^2 (c_1^2 + d_1^2)}{2(\nu + i\varsigma + \sigma(\beta + 1))^2}.$$

Thus applying (7), we obtain

$$|m_3| \leq \frac{2|\alpha\gamma|}{2\sqrt{(\nu + \sigma(2\beta + 1))^2 + \varsigma^2}} + \frac{4\alpha^2\gamma^2}{[(\nu + \sigma(\beta + 1))^2 + \varsigma^2]}.$$

Theorem's proof is now complete. □

**4. Fekete–Szegő inequality for the class  $\mathfrak{G}_\Sigma^\alpha(\sigma, \beta, \nu, \varsigma, \gamma)$**

In this section, we prove the following Fekete–Szegő inequality for functions  $f$  in the class  $\mathfrak{G}_\Sigma^\alpha(\sigma, \beta, \nu, \varsigma, \gamma)$ , using the values of  $m_2^2$  and  $m_3$ .

**THEOREM 4.1.** *For  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$ ,  $0 \leq \beta \leq 1$  and  $\eta \in \mathbb{R}$ , let  $f \in \mathcal{A}$  be in the family  $\mathfrak{G}_\Sigma^\alpha(\sigma, \beta, \nu, \varsigma, \gamma)$ . Then*

$$|m_3 - \eta m_2^2| \leq \begin{cases} \frac{|\alpha\gamma|}{\sqrt{(\nu + \sigma(2\beta + 1))^2 + \varsigma^2}}, & |\eta - 1| \leq |\Omega(\alpha, \nu, \varsigma, \beta)| \\ \frac{8\alpha^2|\gamma|^3|1-\eta|}{|2\alpha\gamma^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, \beta, \nu, \varsigma)) - (2(1 + \alpha)\gamma^2 - 1)(\nu + i\varsigma + \sigma(\beta + 1))^2|}, & |\eta - 1| \geq |\Omega(\alpha, \nu, \varsigma, \beta)| \end{cases}$$

where

$$\Omega(\alpha, \nu, \varsigma, \beta) = \frac{2\alpha\gamma^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, \beta, \nu, \varsigma)) - (2(1 + \alpha)\gamma^2 - 1)(\nu + i\varsigma + \sigma(\beta + 1))^2}{8\alpha\gamma^2(\nu + i\varsigma + \sigma(2\beta + 1))}$$

*Proof.* It follows from (23) and (24) that

$$\begin{aligned} & m_3 - \eta m_2^2 \\ &= \frac{C_1^\alpha(\gamma)(c_2 - d_2)}{4(\nu + i\varsigma + \sigma(2\beta + 1))} + (1 - \eta)m_2^2 \\ &= \frac{C_1^\alpha(\gamma)(c_2 - d_2)}{4(\nu + i\varsigma + \sigma(2\beta + 1))} + \frac{[C_1^\alpha(\gamma)]^3(c_2 + d_2)(1 - \eta)}{[C_1^\alpha(\gamma)]^2 \left( \begin{matrix} (\nu + i\varsigma)(\nu + i\varsigma + 1) \\ + \Upsilon(\sigma, \beta, \nu, \varsigma) \end{matrix} \right) - 2C_2^\alpha(\gamma)(\nu + i\varsigma + \sigma(\beta + 1))^2} \\ &= C_1^\alpha(\gamma) \left[ \left( \psi(\mu, r) + \frac{1}{4(\nu + i\varsigma + \sigma(2\beta + 1))} \right) c_2 + \left( \psi(\mu, r) - \frac{1}{4(\nu + i\varsigma + \sigma(2\beta + 1))} \right) d_2 \right], \end{aligned}$$

where

$$\psi(\eta, \gamma) = \frac{[C_1^\alpha(\gamma)]^2(1 - \eta)}{[C_1^\alpha(\gamma)]^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, \beta, \nu, \varsigma)) - 2C_2^\alpha(\gamma)(\nu + i\varsigma + \sigma(\beta + 1))^2}.$$

According to (7), we find that

$$|m_3 - \eta m_2^2| \leq \begin{cases} \frac{2|C_1^\alpha(\gamma)|}{4(\nu + i\varsigma + \sigma(2\beta + 1))} & 0 \leq |\psi(\eta, \gamma)| \leq \frac{1}{4(\nu + i\varsigma + \sigma(2\beta + 1))}, \\ 2|C_1^\alpha(\gamma)||\psi(\eta, \gamma)| & |\psi(\eta, \gamma)| \geq \frac{1}{4(\nu + i\varsigma + \sigma(2\beta + 1))}. \end{cases}$$

This completes the proof of Theorem 4.1. □

**5. Corollaries and consequences**

Here we find two corollaries where  $\beta = 0$  and  $\beta = 1$ , respectively.

COROLLARY 5.1. For  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$  and  $\beta = 0$ , let  $f \in \mathcal{A}$  be in the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\gamma, 0, \nu, \varsigma, \gamma)$ . Then

$$|m_2| \leq \frac{2|\alpha\gamma| \sqrt{|\alpha\gamma|}}{\sqrt{\left| 2 \left[ \begin{array}{c} \alpha((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, 0, \nu, \varsigma)) \\ - (1 + \alpha)(\nu + i\varsigma + \sigma)^2 \end{array} \right] \gamma^2 + (\nu + i\varsigma + \sigma)^2 \right|}},$$

$$|m_3| \leq \frac{2|\alpha\gamma|}{2\sqrt{(\nu + \sigma)^2 + \varsigma^2}} + \frac{4\alpha^2\gamma^2}{[(\nu + \sigma)^2 + \varsigma^2]},$$

and

$$|m_3 - \eta m_2^2| \leq \begin{cases} \frac{|\alpha\gamma|}{\sqrt{(\nu + \sigma)^2 + \varsigma^2}}, & |\eta - 1| \leq |\Omega(\alpha, \nu, \varsigma, 0)| \\ \frac{8\alpha^2|\gamma|^3|1-\eta|}{|2\alpha\gamma^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + \sigma[2(\nu + i\varsigma + 1) + (\sigma - 1)]) - (-1 + (2 + 2\alpha)\gamma^2)(\nu + i\varsigma + 2\sigma)^2|}, & |\eta - 1| \geq |\Omega(\alpha, \nu, \varsigma, 0)| \end{cases}$$

where

$$\Omega(\alpha, \nu, \varsigma, 0) = \frac{2\alpha\gamma^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, 0, \nu, \varsigma)) - (-1 + (2 + 2\alpha)\gamma^2)(\nu + i\varsigma + \sigma)^2}{8\alpha\gamma^2(\nu + i\varsigma + \sigma)}.$$

COROLLARY 5.2. For  $\nu > 0$ ,  $\varsigma \in \mathbb{R}$ ,  $\nu + i\varsigma \neq 0$ ,  $0 \leq \sigma \leq 1$  and  $\beta = 1$ , let  $f \in \mathcal{A}$  be in the family  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, 1, \nu, \varsigma, \gamma)$ . Then

$$|m_2| \leq \frac{2|\alpha\gamma| \sqrt{|\alpha\gamma|}}{\sqrt{\left| 2 \left[ \begin{array}{c} \alpha((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, 1, \nu, \varsigma)) \\ - (1 + \alpha)(\nu + i\varsigma + 2\sigma)^2 \end{array} \right] \gamma^2 + (\nu + i\varsigma + 2\sigma)^2 \right|}},$$

$$|m_3| \leq \frac{2|\alpha\gamma|}{2\sqrt{(\nu + 3\sigma)^2 + \varsigma^2}} + \frac{4\alpha^2\gamma^2}{[(\nu + 2\sigma)^2 + \varsigma^2]},$$

and

$$|m_3 - \eta m_2^2| \leq \begin{cases} \frac{|\alpha\gamma|}{\sqrt{(\nu + 3\sigma)^2 + \varsigma^2}}, & |\eta - 1| \leq |\Omega(\alpha, \nu, \varsigma, 1)| \\ \frac{8\alpha^2|\gamma|^3|1-\eta|}{|2\alpha\gamma^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + 2\sigma[2(\nu + i\varsigma + 1) + 2(\sigma - 1)]) - (-1 + (2 + 2\alpha)\gamma^2)(\nu + i\varsigma + 2\sigma)^2|}, & |\eta - 1| \geq |\Omega(\alpha, \nu, \varsigma, 1)| \end{cases}$$

where

$$\Omega(\alpha, \nu, \varsigma, 1) = \frac{2\alpha\gamma^2((\nu + i\varsigma)(\nu + i\varsigma + 1) + \Upsilon(\sigma, 1, \nu, \varsigma)) - (-1 + (2 + 2\alpha)\gamma^2)(\nu + i\varsigma + 2\sigma)^2}{8\alpha\gamma^2(\nu + i\varsigma + 3\sigma)}.$$



REMARK 5.3. The findings of this study would lead to a number of other novel findings for the classes  $\mathfrak{G}_{\Sigma}^1(\sigma, \beta, \nu, \varsigma, \gamma)$  for Chebyshev polynomials and  $\mathfrak{G}_{\Sigma}^{0,5}(\sigma, \beta, \nu, \varsigma, \gamma)$  for Legendre polynomials.

**Concluding Remark:** The problems with the coefficients that are associated with each new subclass  $\mathfrak{G}_{\Sigma}^{\alpha}(\sigma, \beta, \nu, \varsigma, \gamma)$ , of the class of bi-univalent functions on the open unit disk  $\mathfrak{h}$ , have been presented in this work and investigated. These problems can be found in the class of bi-univalent functions. The definitions for this bi-univalent function class can be found in reference 2.1. The estimates of the Taylor-Maclaurin coefficients  $|m_2|$  and  $|m_3|$ , as well as the Fekete-Szegö functional problems for functions belonging to this new subclass, have been obtained. The estimations presented here are applicable to functions that fall into each of the bi-univalent function classes. The number of factors that were evaluated for our original results was reduced, and as a consequence, a number of new outcomes have been demonstrated to follow. The problem of obtaining estimates on the boundaries of the value  $|m_n|$  for  $n \geq 4; n \in \mathbb{N}$ , for the classes stated in this paragraph still exists.

## 6. Statements and Declarations

**6.1. Ethical Approval.** Not applicable.

**6.2. Competing Interests.** The authors declare no conflicts of interest.

**6.3. Authors Contributions.** Conceptualization, O.A. and A.A.; methodology, M.D. and O.A.; software, M.D. and O.A.; validation, A.A. and O.A.; formal analysis, O.A. and M.D., investigation, O.A. and A.A.; resources, O.A. and A.A.; data curation, O.A. and A.A., writing—original draft preparation, O.A. and A.A.; writing—review and editing, M.D. and A.A.; visualization, O.A. and M.D.; supervision, M.D. and A.A.; project administration, O.A. and A.A.; funding acquisition, A.A.

All authors have read and agreed to the published version of the manuscript.

**6.4. Availability of Data and Materials.** A statement on how any datasets used can be accessed.

## References

- [1] A. Amourah, B. A. Frasin, M. Ahmad and F. Yousef, *Exploiting the Pascal Distribution Series and Gegenbauer Polynomials to Construct and Study a New Subclass of Analytic Bi-Univalent Functions*, *Symmetry*. **14** (1) (2022), 147.  
<https://dx.doi.org/10.3390/sym14010147>
- [2] A. Amourah, B. A. Frasin, G. Murugusundaramoorthy and T. Al-Hawary, *Bi-Bazilevič functions of order  $\vartheta + i\delta$  associated with  $(p, q)$ -Lucas polynomials*, *AIMS Mathematics*. **6** (5) (2021), 4296–4305.
- [3] A. Amourah, T. Al-Hawary and B. A. Frasin, *Application of Chebyshev polynomials to certain class of bi-Bazilevič functions of order  $\alpha + i\beta$* , *Afr. Mat.* (2021), 1–8.
- [4] A. Amourah, B.A. Frasin and T. Abdeljawad, *Fekete-Szegö inequality for analytic and bi-univalent functions subordinate to Gegenbauer polynomials*, *J. Funct. Spaces*. (2021), 1–7.

- [5] A. Amourah, O. Alnajjar, M. Darus, A. Shdouh and O. Ogilat, *Estimates for the Coefficients of Subclasses Defined by the Bell Distribution of Bi-Univalent Functions Subordinate to Gegenbauer Polynomials*, Mathematics. **11** (8) (2023), 1799.  
<https://dx.doi.org/10.3390/math11081799>
- [6] A. Legendre, *Recherches sur laattraction des sphéroides homogènes*, Mémoires présentes par divers savants a laAcadémie des Sciences de laInstitut de France, Paris. **10** (1785), 411–434.
- [7] B. A. Frasin and M. K. Aouf, *New subclasses of bi-univalent functions*, Appl. Math. Lett. **24** (2011), 1569–1573.  
<https://dx.doi.org/10.1016/j.aml.2011.03.048>
- [8] B.A. Frasin, S.R. Swamy and J. Nirmala, *Some special families of holomorphic and Al-Oboudi type bi-univalent functions related to  $k$ -Fibonacci numbers involving modified Sigmoid activation function*, Afrika Matematika. **32** (2021), 631–643.
- [9] E. Kobald, J. A. Gmeiner and O. Stolz, *Leopold Gegenbauer*, (1904), 3–10.
- [10] F. Yousef, S. Alroud and M. Illafe, *A comprehensive subclass of bi-univalent functions associated with Chebyshev polynomials of the second kind*, Boletín de la Sociedad Matemática Mexicana. (2019), 1–11.
- [11] F. Yousef, S. Alroud and M. Illafe, *New subclasses of analytic and bi-univalent functions endowed with coefficient estimate problems*, Anal. Math. Physics. **11** (2021), 1–12.
- [12] F. Yousef, T. Al-Hawary and G. Murugusundaramoorthy, *Fekete-Szegö functional problems for some subclasses of bi-univalent functions defined by Frasin differential operator*, Afrika Matematika. **30** (2019), 495–503.
- [13] G. Murugusundaramoorthy, *Subclasses of starlike and convex functions involving Poisson distribution series*, Afr. Mat. **28** (2017), 1357–1366.
- [14] G. Murugusundaramoorthy, N. Magesh and V. Prameela, *Coefficient bounds for certain subclasses of bi-univalent function*, Abst. Appl. Anal. (2013).  
<http://dx.doi.org/10.1155/2013/573017>
- [15] H. M. Srivastava, Ş. Altinkaya and S. Yalçın, *Certain subclasses of bi-univalent functions associated with the Horadam polynomials. Iranian Journal of Science and Technology*, Transactions. A. Science. **43** (4) (2019), 1873–1879.
- [16] H. M. Srivastava, A. K. Mishra and P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Applied mathematics letters. **23** (10) (2010), 1188–1192.  
<http://dx.doi.org/10.1016/j.aml.2010.05.009>
- [17] H. M. Srivastava, G. Murugusundaramoorthy and N. Magesh, 2013. *Certain subclasses of bi-univalent functions associated with the Hohlov operator* Global J. Math. Anal. **1** (2) (2013), 67-73.
- [18] H. M. Srivastava and D. Bansal, *Coefficient estimates for a class of analytic and bi-univalent functions*, Journal of the Egyptian Mathematical Society. **23** (2) (2015), 242-246.  
<http://dx.doi.org/10.1016/j.joems.2014.04.002>
- [19] H. O. Guney, G. Murugusundaramoorthy and Sokol, *Subclasses of bi-univalent functions related to shell-like curves connected with Fibonacci numbers*, Acta Universitatis Sapientiae, Mathematica. **10** (2018), 70-84.  
<https://dx.doi.org/10.2478/ausm-2018-0006>
- [20] H. U .Rehman, M. Darus and J. Salah, *Generalizing Certain Analytic Functions Correlative to the  $n$ -th Coefficient of Certain Class of Bi-Univalent Functions*, Journal of Mathematics. (2021), 1–14.  
<https://dx.doi.org/10.1155/2021/6621315>
- [21] M. Fekete and G. Szegö, *Eine Bemerkung Äber ungerade schlichte Funktionen*, J. Lond. Math. Soc. **1** (2) (1933), 85–89.
- [22] M. Lewin, *On a Coefficient problem for bi-univalent functions*, Proc Amer Math Soc. **18** (1967), 63–68.
- [23] N. M. Atakishiyev, M. Rahman and S . K. Suslov, *On classical orthogonal polynomials*, Constructive Approximation. **11** (1995), 181–226.
- [24] O. Alnajjar, A. Amourah and M. Darus, *The Characteristics of Inclusion Pertaining to Univalent Functions Associated with Bell Distribution Functions*, Int. J. Open Problems Complex Analysis. **15** (2) (2023), 46–61.

- [25] S. Porwal and M. Darus, *On a new subclass of bi-univalent functions*, Journal of the Egyptian Mathematical Society. **21** (3) (2013), 190-193.  
<http://dx.doi.org/10.1016/j.joems.2013.02.007>
- [26] S. Altinkaya and S. Yalcin, *On the  $(p, q)$ -Lucas polynomial coefficient bounds of the bi-univalent function class*, Boletin de la Sociedad Matematica Mexicana. **25** (3) (2018), 567–575.
- [27] V. Kumar, N. E. Cho, V. Ravichandran and H. M. Srivastava, *Sharp coefficient bounds for starlike functions associated with the Bell numbers*, Mathematica Slovaca. **69** (5) (2019), 1053–1064.  
<http://dx.doi.org/10.1515/ms-2017-0289>
- [28] Z. Peng, G. Murugusundaramoorthy and T. Janani, *Coefficient estimate of bi-univalent functions of complex order associated with the Hohlov operator*, J. Complex Analysis. (2014).  
<http://dx.doi.org/10.1155/2014/693908>

**Omar Alnajjar**

Department of Mathematical Sciences, Faculty of Science and Technology,  
Universiti Kebangsaan Malaysia, Bangi 43600, Malaysia  
*E-mail*: P117246@siswa.ukm.edu.my

**Ala Amourah**

Mathematics Education Program, Faculty of Education and Arts, Sohar University,  
Sohar 3111, Oman.  
*E-mail*: AAmourah@su.edu.om

**Maslina Darus**

Department of Mathematical Sciences, Faculty of Science and Technology,  
Universiti Kebangsaan Malaysia, Bangi 43600, Malaysia  
*E-mail*: maslina@ukm.edu.my