

WEAK EXTENDED ORDER ALGEBRAS HAVING ADJOINT TRIPLES

YONG CHAN KIM AND JUNG MI KO*

ABSTRACT. We study the properties of weak extended order algebras having adjoint pairs (triples) or Galois pairs. In particular, we investigate the various laws on weak extended order algebras.

1. Introduction

Wille [9] introduced the structures on lattices which are important mathematical tools for data analysis and knowledge processing. MV-algebra was introduced by Chang [2] to provide algebraic models for many valued propositional logic. Recently, it is developed in many directions (BL-algebra, residuated algebra) [1,3,4,7,9]. Recently, Guido et al. [5] introduced extended order algebras as the generalization of residuated algebras. Kim and Ko [6] introduced the properties of weak extended order algebras. In particular, we investigate the properties of commutative and associative extended-order algebras.

In this paper, we study the properties of weak extended order algebras having adjoint pairs (triples) or Galois pairs. In particular, we investigate the various laws on weak extended order algebras.

Received May 14, 2013. Revised June 25, 2013. Accepted September 13, 2013.

2010 Mathematics Subject Classification: 06A06, 06A15, 06B30, 54F05, 68U35.

Key words and phrases: (Commutative, associative) weak extended order algebras, adjoint pairs (triples), Galois pairs.

This work was supported by Research Institute of Natural Science of Gangneung-Wonju National University.

*Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2013.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

2. Preliminaries

DEFINITION 2.1. [5,6] Let (L, \wedge, \vee) be a lattice. A triple (L, \Rightarrow, \top) is called a *weak extended order algebra* (shortly, w-eo algebra) iff it satisfies the following properties:

- (O1) $a \Rightarrow \top = \top$ (upper bounded condition)
- (O2) $a \Rightarrow a = \top$ (reflexive condition)
- (O3) if $a \Rightarrow b = \top$ and $b \Rightarrow a = \top$, then $a = b$
- (O4) if $a \Rightarrow b = \top$ and $b \Rightarrow c = \top$, then $a \Rightarrow c = \top$.

A triple (L, \Rightarrow, \top) is called a *right w-eo algebra* if it satisfies (O1), (O2), (O3) and

- (O5) if $a \Rightarrow b = \top$, then $(c \Rightarrow a) \Rightarrow (c \Rightarrow b) = \top$.

A triple (L, \Rightarrow, \top) is called a *left w-eo algebra* if it satisfies (O1), (O2), (O3) and

- (O6) if $a \Rightarrow b = \top$, then $(b \Rightarrow c) \Rightarrow (a \Rightarrow c) = \top$.

A w-eo algebra is called a *right distributive w-eo algebra* if

- (O7) $a \Rightarrow \bigwedge_i b_i = \bigwedge_i (a \Rightarrow b_i)$.

A w-eo algebra is called a *left distributive w-eo algebra* if

- (O8) $\bigvee_i a_i \Rightarrow b = \bigwedge_i (a_i \Rightarrow b)$.

(1) A w-eo algebra has an *adjoint pair* (\Rightarrow, \odot) if there exists a binary operation \odot such that

$$a \odot b \leq c \text{ iff } b \leq a \Rightarrow c.$$

(2) A w-eo algebra has a *Galois pair* $(\Rightarrow, \rightarrow)$ if there exists a binary operation \rightarrow such that

$$b \leq a \Rightarrow c \text{ iff } a \leq b \rightarrow c.$$

(3) A w-eo algebra has *symmetrical* if it has a Galois pair $(\Rightarrow, \rightarrow)$ and (L, \rightarrow, \top) is a w-eo algebra.

(4) A w-eo algebra has an *adjoint triple* $(\Rightarrow, \odot, \rightarrow)$ if there exists binary operation \odot and \rightarrow such that $a \odot b \leq c$ iff $b \leq a \Rightarrow c$ iff $a \leq b \rightarrow c$.

(5) A w-eo algebra is called a *w-ceo algebra* if L is complete.

THEOREM 2.2. [6] (1) If (L, \Rightarrow, \top) is a right w-eo algebra, then it is a w-eo algebra.

(2) If (L, \Rightarrow, \top) is a left w-eo algebra, then it is a w-eo algebra.

THEOREM 2.3. [5] Let (L, \Rightarrow, \top) be a right-distributive w-ceo algebra and \odot be defined by

$$a \odot x = \bigwedge \{y \in L \mid x \leq a \Rightarrow y\}.$$

Then \odot and \Rightarrow form an adjoint pair, i.e.

$$x \odot y \leq z \text{ iff } y \leq x \Rightarrow z.$$

THEOREM 2.4. [5] Let (L, \Rightarrow, \top) be a left-distributive w-ceo algebra and \rightarrow be defined by

$$g_a(y) = y \rightarrow a = \bigvee \{x \in L \mid y \leq x \Rightarrow a\}.$$

Then $(\Rightarrow, \rightarrow)$ forms an adjoint pair, i.e.

$$y \leq x \Rightarrow a \text{ iff } x \leq y \rightarrow a.$$

THEOREM 2.5. [6] Let (L, \Rightarrow, \top) be a w-eo algebra having an adjoint pair (\Rightarrow, \odot) and \Rightarrow be defined by

$$a \Rightarrow b = \top \text{ iff } a \leq b$$

For each $a, b, c, a_i, b_i \in L$, the following properties hold.

- (1) $a \odot b \leq a$ and $a \odot (a \Rightarrow b) \leq b \leq a \Rightarrow a \odot b$;
- (2) $a \odot \top = a$;
- (3) $a \odot \perp = \perp \odot a = \perp$;
- (4) If $b \leq c$, then $a \odot b \leq a \odot c$.
- (5) (L, \Rightarrow, \top) be a right w-eo algebra.
- (6) If (L, \Rightarrow, \top) is a complete lattice, (L, \Rightarrow, \top) is a right distributive w-eo algebra and $a \odot (\bigvee_{i \in \Gamma} b_i) = \bigvee_{i \in \Gamma} (a \odot b_i)$.
- (7) (L, \Rightarrow, \top) is a left w-eo algebra iff $a \odot c \leq b \odot c$ for $a \leq b$.
- (8) If (L, \Rightarrow, \top) is a left w-eo algebra, then (L, \Rightarrow, \top) be a right w-eo algebra.
- (9) If (L, \Rightarrow, \top) is a left w-eo algebra and $a \odot (b \odot c) = (a \odot b) \odot c$, then

$$(a \Rightarrow b) \Rightarrow ((c \Rightarrow a) \Rightarrow (c \Rightarrow b)) = \top.$$

- (10) If (L, \Rightarrow, \top) is a complete and left w-eo algebra, then (L, \Rightarrow, \top) is a left distributive w-eo algebra and $(\bigvee_{i \in \Gamma} a_i) \odot b = \bigvee_{i \in \Gamma} (a_i \odot b)$.

- (11) If $\top \Rightarrow a = a$, then $\top \odot a = a$, $a = \bigwedge_b (b \Rightarrow (b \odot a))$ and $a = \bigvee_b (b \odot (b \Rightarrow a))$.

- (12) If (L, \Rightarrow, \top) is a left w-eo algebra and $\top \Rightarrow a = a$, then $a \odot b \leq b$ and $a \Rightarrow (b \Rightarrow a) = \top$.

THEOREM 2.6. [6] Let (L, \Rightarrow, \top) be a w-eo algebra having a Galois pair $(\Rightarrow, \rightarrow)$. For each $a, b, c \in L$, the following properties hold.

- (1) $a \leq (a \rightarrow b) \Rightarrow b$ and $a \leq (a \Rightarrow b) \rightarrow b$.

(2) $\top \rightarrow a = a$. Furthermore, if $(L, \Rightarrow, \rightarrow, \top)$ is a symmetrical w-eo algebra, then $\top \Rightarrow a = a$.

(3) If $\top \Rightarrow a = a$, then $a \rightarrow b = \top$ iff $a \leq b$.

(4) (L, \Rightarrow, \top) is a left w-eo algebra. If $\top \Rightarrow a = a$, (L, \rightarrow, \top) is a left w-eo algebra.

(5) (L, \Rightarrow, \top) is a right w-eo algebra iff (L, \rightarrow, \top) is a right w-eo algebra.

(6) $a = \bigwedge_b((a \Rightarrow b) \rightarrow b)$. If $\top \Rightarrow a = a$, then $a = \bigwedge_b((a \rightarrow b) \Rightarrow b)$.

(7) If (L, \Rightarrow, \top) is a complete lattice, $\bigvee_i a_i \Rightarrow b = \bigwedge_i(a_i \Rightarrow b)$ and $\bigvee_i a_i \rightarrow b = \bigwedge_i(a_i \rightarrow b)$.

THEOREM 2.7. [6] Let (L, \Rightarrow, \top) be a w-eo algebra having adjoint triple with \odot and \rightarrow . Then the following properties hold.

(1) (L, \Rightarrow, \top) is an eo algebra;

(2) If $\top \Rightarrow a = a$, then (L, \rightarrow, \top) is an eo algebra;

(3) If $a \leq b$, then $a \odot c \leq b \odot c$ and $c \odot a \leq c \odot b$.

(4) If $\top \Rightarrow a = a$, then $\top \odot a = a$.

DEFINITION 2.8. [5] A w-eo algebra (L, \Rightarrow, \top) is commutative iff it satisfies

$$a \Rightarrow (b \Rightarrow c) = \top \text{ iff } b \Rightarrow (a \Rightarrow c) = \top$$

THEOREM 2.9. [6] Let (L, \Rightarrow, \top) be a w-eo algebra having adjoint triple with \odot and \rightarrow . Then the following statements are equivalent:

(1) (L, \Rightarrow, \top) is commutative;

(2) (L, \odot, \top) is commutative;

(3) (L, \rightarrow, \top) is commutative eo algebra with $\Rightarrow = \rightarrow$.

(4) $a \rightarrow b \leq a \Rightarrow b$ for all $a, b \in L$.

(5) $a \Rightarrow b \leq a \rightarrow b$ for all $a, b \in L$.

(6) $a \leq (a \Rightarrow b) \Rightarrow b$ for all $a, b \in L$.

(7) $a \leq (a \rightarrow b) \rightarrow b$ for all $a, b \in L$.

(8) $(a \Rightarrow b) \odot a \leq b$ for all $a, b \in L$.

(9) $a \odot (a \rightarrow b) \leq b$ for all $a, b \in L$.

(10) $b \leq a \Rightarrow b \odot a$ for all $a, b \in L$.

(11) $a \leq b \rightarrow b \odot a$ for all $a, b \in L$.

3. Weak extended order algebras having adjoint triples

THEOREM 3.1. *Let (L, \Rightarrow, \top) be a w-*eo* algebra having adjoint triple with \odot and \rightarrow . Then the following statements are equivalent:*

- (1) $a \Rightarrow (b \Rightarrow c) = b \Rightarrow (a \Rightarrow c)$ for all $a, b, c \in L$.
- (2) $b \rightarrow c \leq (a \Rightarrow b) \rightarrow (a \Rightarrow c)$ for all $a, b, c \in L$.
- (3) $a \odot (b \odot c) = b \odot (a \odot c)$ for all $a, b, c \in L$.
- (4) $b \rightarrow c \leq (a \odot b) \rightarrow (a \odot c)$ for all $a, b, c \in L$.
- (5) $b \rightarrow (a \Rightarrow c) = (a \odot b) \rightarrow c$ for all $a, b, c \in L$.

Proof. (1) \Leftrightarrow (2) Since $a \Rightarrow (b \Rightarrow c) \leq b \Rightarrow (a \Rightarrow c)$, then $b \leq (a \Rightarrow (b \Rightarrow c)) \rightarrow (a \Rightarrow c)$. Put $b = b \rightarrow c$. Since $b \leq (b \rightarrow c) \Rightarrow c$, then

$$\begin{aligned} b \rightarrow c &\leq (a \Rightarrow ((b \rightarrow c) \Rightarrow c)) \rightarrow (a \Rightarrow c) \\ &\leq (a \Rightarrow b) \rightarrow (a \Rightarrow c). \end{aligned}$$

Conversely, since $b \leq (b \Rightarrow c) \rightarrow c \leq (a \Rightarrow (b \Rightarrow c)) \rightarrow (a \Rightarrow c)$, then $a \Rightarrow (b \Rightarrow c) \leq b \Rightarrow (a \Rightarrow c)$.

(1) \Leftrightarrow (3) Since $a \Rightarrow (b \Rightarrow a \odot (b \odot c)) = b \Rightarrow (a \Rightarrow a \odot (b \odot c)) \geq b \Rightarrow (b \odot c) \geq c$, then

$$\begin{aligned} a \Rightarrow (b \Rightarrow a \odot (b \odot c)) \geq c &\text{ iff } b \Rightarrow a \odot (b \odot c) \geq a \odot c \\ &\text{ iff } a \odot (b \odot c) \geq b \odot (a \odot c). \end{aligned}$$

Similarly, $a \odot (b \odot c) \leq b \odot (a \odot c)$.

Conversely, since $a \odot (b \odot (a \Rightarrow (b \Rightarrow c))) = b \odot (a \odot (a \Rightarrow (b \Rightarrow c))) \leq b \odot (b \Rightarrow c) \leq c$, we have

$$\begin{aligned} a \odot (b \odot (a \Rightarrow (b \Rightarrow c))) \leq c &\text{ iff } b \odot (a \Rightarrow (b \Rightarrow c)) \leq a \Rightarrow c \\ &\text{ iff } a \Rightarrow (b \Rightarrow c) \leq b \Rightarrow (a \Rightarrow c) \end{aligned}$$

Similarly, $a \Rightarrow (b \Rightarrow c) \geq b \Rightarrow (a \Rightarrow c)$.

(3) \Leftrightarrow (4)

$$\begin{aligned} (b \rightarrow c) \odot (a \odot b) &= a \odot ((b \rightarrow c) \odot b) \leq a \odot c \\ \text{iff } b \rightarrow c &\leq (a \odot b) \rightarrow (a \odot c). \end{aligned}$$

$$\begin{aligned} b \leq c \rightarrow b \odot c &\leq (a \odot c) \rightarrow a \odot (b \odot c) \\ \text{iff } b \odot (a \odot c) &\leq a \odot (b \odot c). \end{aligned}$$

(3) \Leftrightarrow (5) By (4), put $c = a \Rightarrow c$.

$$b \rightarrow (a \Rightarrow c) \leq (a \odot b) \rightarrow a \odot (a \Rightarrow c) \leq a \odot b \rightarrow c.$$

$$\begin{aligned} & (a \odot b \rightarrow c) \odot (a \odot b) \leq c \\ & \text{iff } a \odot ((a \odot b \rightarrow c) \odot b) \leq c \\ & \text{iff } (a \odot b \rightarrow c) \odot b \leq a \Rightarrow c \\ & \text{iff } a \odot b \rightarrow c \leq b \rightarrow (a \Rightarrow c). \end{aligned}$$

Hence $a \odot b \rightarrow c \leq b \rightarrow (a \Rightarrow c)$.

Conversely, since $b \odot c \rightarrow b \odot (a \odot c) = c \rightarrow (b \Rightarrow b \odot (a \odot c)) \geq c \rightarrow (a \odot c) \geq a$, we have

$$b \odot c \rightarrow b \odot (a \odot c) \geq a \text{ iff } a \odot (b \odot c) \leq b \odot (a \odot c).$$

□

THEOREM 3.2. *Let (L, \Rightarrow, \top) be a w-eo algebra having adjoint triple with \odot and \rightarrow . Then the following statements are equivalent:*

- (1) $a \rightarrow (b \rightarrow c) = b \rightarrow (a \rightarrow c)$ for all $a, b, c \in L$.
- (2) $b \Rightarrow c \leq (a \rightarrow b) \Rightarrow (a \rightarrow c)$ for all $a, b, c \in L$.
- (3) $(a \odot b) \odot c = (a \odot c) \odot b$ for all $a, b, c \in L$.
- (4) $b \Rightarrow c \leq (b \odot a) \Rightarrow (c \odot a)$ for all $a, b, c \in L$.
- (5) $b \Rightarrow (a \rightarrow c) = (b \odot a) \Rightarrow c$ for all $a, b, c \in L$.

Proof. (1) \Leftrightarrow (2) Since $a \rightarrow (b \rightarrow c) \leq b \rightarrow (a \rightarrow c)$, then $b \leq (a \rightarrow (b \rightarrow c)) \Rightarrow (a \rightarrow c)$. Put $b = b \Rightarrow c$. Since $b \leq (b \Rightarrow c) \rightarrow c$, then

$$\begin{aligned} b \Rightarrow c & \leq (a \rightarrow ((b \Rightarrow c) \rightarrow c)) \Rightarrow (a \rightarrow c) \\ & \leq (a \rightarrow b) \Rightarrow (a \rightarrow c) \end{aligned}$$

Since $b \leq (b \rightarrow c) \Rightarrow c \leq (a \rightarrow (b \rightarrow c)) \Rightarrow (a \rightarrow c)$, then $a \rightarrow (b \rightarrow c) \leq b \rightarrow (a \rightarrow c)$.

(1) \Leftrightarrow (3) Since $c \rightarrow (b \rightarrow (a \odot b) \odot c) = b \rightarrow (c \rightarrow (a \odot b) \odot c) \geq b \rightarrow (a \odot b) \geq a$, we have

$$\begin{aligned} c \rightarrow (b \rightarrow (a \odot b) \odot c) \geq a & \text{ iff } b \rightarrow (a \odot b) \odot c \geq a \odot c \\ & \text{ iff } (a \odot b) \odot c \geq (a \odot c) \odot b. \end{aligned}$$

Conversely, since $((a \rightarrow (b \rightarrow c)) \odot b) \odot a = ((a \rightarrow (b \rightarrow c)) \odot a) \odot b \leq (b \rightarrow c) \odot b \leq c$, we have

$$\begin{aligned} ((a \rightarrow (b \rightarrow c)) \odot b) \odot a \leq c & \text{ iff } (a \rightarrow (b \rightarrow c)) \odot b \leq a \rightarrow c \\ & \text{ iff } a \rightarrow (b \rightarrow c) \leq b \rightarrow (a \rightarrow c). \end{aligned}$$

(3) \Leftrightarrow (4)

$$(b \odot a) \odot (b \Rightarrow c) = (b \odot (b \Rightarrow c)) \odot a \leq c \odot a$$

$$\text{iff } b \Rightarrow c \leq (b \odot a) \Rightarrow (c \odot a)$$

Conversely, since $c \leq a \Rightarrow a \odot c \leq (a \odot b) \Rightarrow (a \odot c) \odot b$, we have

$$c \leq (a \odot b) \Rightarrow (a \odot c) \odot b \text{ iff } (a \odot b) \odot c \leq (a \odot c) \odot b.$$

(3) \Leftrightarrow (5) By (4), put $c = a \rightarrow c$.

$$b \Rightarrow (a \rightarrow c) \leq (b \odot a) \Rightarrow (a \rightarrow c) \odot a \leq b \odot a \Rightarrow c.$$

Since $(b \odot (b \odot a \Rightarrow c)) \odot (b \Rightarrow b \odot a) = (b \odot (b \Rightarrow b \odot a)) \odot (b \odot a \rightarrow c) \leq (b \odot a) \odot (b \odot a \Rightarrow c) \leq c$, then $b \odot (b \odot a \Rightarrow c) \leq (b \Rightarrow b \odot a) \rightarrow c \leq a \rightarrow c$. So, $b \odot a \Rightarrow c \leq b \Rightarrow (a \rightarrow c)$. Hence $b \odot a \Rightarrow c = b \Rightarrow (a \rightarrow c)$.

Conversely, since $a \odot b \Rightarrow (a \odot c) \odot b = a \Rightarrow (b \rightarrow (a \odot c) \odot b) \geq a \Rightarrow a \odot c \geq c$, we have

$$a \odot b \Rightarrow (a \odot c) \odot b \geq c \text{ iff } (a \odot b) \odot c \leq (a \odot c) \odot b.$$

□

DEFINITION 3.3. [5] Let (L, \Rightarrow, \top) be a w-eo algebra having an adjoint pair (\Rightarrow, \odot) . A w-eo algebra (L, \Rightarrow, \top) is associative iff it satisfies

$$a \Rightarrow (b \Rightarrow c) = (b \odot a) \Rightarrow c.$$

THEOREM 3.4. Let (L, \Rightarrow, \top) be a w-eo algebra having an adjoint pair (\Rightarrow, \odot) . Then the following statements are equivalent:

- (1) (L, \Rightarrow, \top) is associative;
- (2) $(a \odot b) \odot c = a \odot (b \odot c)$ for all $a, b, c \in L$.

Proof. (1) \Rightarrow (2)

$$(a \odot b) \odot c \leq a \odot (b \odot c)$$

$$\text{iff } c \leq (a \odot b) \Rightarrow (a \odot (b \odot c)) = b \Rightarrow (a \Rightarrow (a \odot (b \odot c)))$$

$$\text{iff } b \odot c \leq a \Rightarrow a \odot (b \odot c)$$

$$\text{iff } a \odot (b \odot c) \leq a \odot (b \odot c).$$

$$\begin{aligned}
& a \odot (b \odot c) \leq (a \odot b) \odot c \\
& \text{iff } b \odot c \leq a \Rightarrow ((a \odot b) \odot c) \\
& \text{iff } c \leq b \Rightarrow (a \Rightarrow ((a \odot b) \odot c)) = a \odot b \Rightarrow ((a \odot b) \odot c) \\
& \text{iff } (a \odot b) \odot c \leq (a \odot b) \odot c.
\end{aligned}$$

(2) \Rightarrow (1) Since $(b \odot a) \odot (a \Rightarrow (b \Rightarrow c)) = b \odot (a \odot (a \Rightarrow (b \Rightarrow c))) \leq b \odot (b \Rightarrow c) \leq c$, we have $a \Rightarrow (b \Rightarrow c) \leq (b \odot a) \Rightarrow c$.

$$\begin{aligned}
& (b \odot a) \Rightarrow c \leq a \Rightarrow (b \Rightarrow c) \\
& \text{iff } a \odot ((b \odot a) \Rightarrow c) \leq b \Rightarrow c \\
& \text{iff } (b \odot a) \odot ((b \odot a) \Rightarrow c) \leq c.
\end{aligned}$$

□

THEOREM 3.5. *Let (L, \Rightarrow, \top) be a right w-eo algebra having a Galois pair $(\Rightarrow, \rightarrow)$. Then the following statements are equivalent:*

- (1) $a \Rightarrow (b \rightarrow c) = b \rightarrow (a \Rightarrow c)$.
- (2) $b \rightarrow a \leq (a \rightarrow c) \Rightarrow (b \rightarrow c)$ and $b \Rightarrow a \leq (a \Rightarrow c) \rightarrow (b \Rightarrow c)$.

Proof. (1) \Rightarrow (2) Since $(a \rightarrow c) \Rightarrow c \geq a$, we have

$$(a \rightarrow c) \Rightarrow (b \rightarrow c) = b \rightarrow ((a \rightarrow c) \Rightarrow c) \geq b \rightarrow a.$$

Since $(a \Rightarrow c) \rightarrow c \geq a$, we have

$$(a \Rightarrow c) \rightarrow (b \Rightarrow c) = b \Rightarrow ((a \Rightarrow c) \rightarrow c) \geq b \Rightarrow a.$$

(2) \Rightarrow (1) Since $a \leq (a \Rightarrow c) \rightarrow c$,

$$\begin{aligned}
a \Rightarrow (b \rightarrow c) & \geq ((a \Rightarrow c) \rightarrow c) \Rightarrow (b \rightarrow c) \\
& \geq b \rightarrow (a \Rightarrow c).
\end{aligned}$$

Since $b \leq (b \rightarrow c) \Rightarrow c$,

$$\begin{aligned}
b \rightarrow (a \Rightarrow c) & \geq ((b \rightarrow c) \Rightarrow c) \rightarrow (a \Rightarrow c) \\
& \geq a \Rightarrow (b \rightarrow c).
\end{aligned}$$

□

THEOREM 3.6. *Let (L, \Rightarrow, \top) be a right w-eo algebra having adjoint triple with \odot and \rightarrow . Then the following statements are equivalent:*

- (1) $a \Rightarrow (b \rightarrow c) = b \rightarrow (a \Rightarrow c)$.
- (2) (L, \odot, \top) is associative.
- (3) $a \rightarrow (b \rightarrow c) = (a \odot b) \rightarrow c$.

Proof. (1) \Rightarrow (2)

$$\begin{aligned} & (a \odot b) \odot c \leq a \odot (b \odot c) \\ & \text{iff } a \odot b \leq c \rightarrow (a \odot (b \odot c)) \\ & \text{iff } b \leq a \Rightarrow (c \rightarrow (a \odot (b \odot c))) = c \rightarrow (a \Rightarrow (a \odot (b \odot c))) \\ & \text{iff } b \odot c \leq a \Rightarrow (a \odot (b \odot c)) \\ & \text{iff } a \odot (b \odot c) \leq a \odot (b \odot c) \end{aligned}$$

$$\begin{aligned} & a \odot (b \odot c) \leq (a \odot b) \odot c \\ & \text{iff } b \odot c \leq a \Rightarrow (a \odot b) \odot c \\ & \text{iff } b \leq c \rightarrow (a \Rightarrow ((a \odot b) \odot c)) = a \Rightarrow (c \rightarrow ((a \odot b) \odot c)) \\ & \text{iff } a \odot b \leq c \rightarrow (a \odot b) \odot c \\ & \text{iff } (a \odot b) \odot c \leq (a \odot b) \odot c \end{aligned}$$

(2) \Rightarrow (1) Since $a \odot ((a \Rightarrow (b \rightarrow c)) \odot b) = (a \odot (a \Rightarrow (b \rightarrow c))) \odot b \leq (b \rightarrow c) \odot b \leq c$, then $(a \Rightarrow (b \rightarrow c)) \odot b \leq a \Rightarrow c$. So, $a \Rightarrow (b \rightarrow c) \leq b \rightarrow (a \Rightarrow c)$.

Since $(a \odot (b \rightarrow (a \Rightarrow c))) \odot b = a \odot ((b \rightarrow (a \Rightarrow c)) \odot b) \leq a \odot (a \Rightarrow c) \leq c$, then $a \odot (b \rightarrow (a \Rightarrow c)) \leq b \rightarrow c$. So, $b \rightarrow (a \Rightarrow c) \leq a \Rightarrow (b \rightarrow c)$.

(2) \Leftrightarrow (3) Since $(a \rightarrow (b \rightarrow c)) \odot (a \odot b) = ((a \rightarrow (b \rightarrow c)) \odot a) \odot b \leq c$, then $a \rightarrow (b \rightarrow c) \leq (a \odot b) \rightarrow c$.

Since $((a \odot b) \rightarrow c) \odot a \odot b = ((a \odot b) \rightarrow c) \odot (a \odot b) \leq c$, then $(a \odot b) \rightarrow c \leq a \rightarrow (b \rightarrow c)$.

Conversely, it follows from;

$$\begin{aligned} & (a \odot b) \odot c \leq (a \odot b) \odot c \\ & \text{iff } a \odot b \leq c \rightarrow (a \odot b) \odot c \\ & \text{iff } a \leq b \rightarrow (c \rightarrow ((a \odot b) \odot c)) = (b \odot c) \rightarrow (a \odot b) \odot c \\ & \text{iff } a \odot (b \odot c) \leq (a \odot b) \odot c, \end{aligned}$$

$$\begin{aligned} & a \odot (b \odot c) \leq a \odot (b \odot c) \\ & \text{iff } a \leq (b \odot c) \rightarrow a \odot (b \odot c) \\ & \text{iff } a \leq b \rightarrow (c \rightarrow a \odot (b \odot c)) \\ & \text{iff } a \odot b \leq c \rightarrow a \odot (b \odot c) \\ & \text{iff } (a \odot b) \odot c \leq (a \odot b) \odot c. \end{aligned}$$

□

THEOREM 3.7. *Let (L, \Rightarrow, \top) be a w-eo algebra having adjoint triple with \odot and \rightarrow satisfying $\top \Rightarrow a = a$ for all $a \in L$. Then the following statements are equivalent:*

- (1) $a \Rightarrow (b \Rightarrow c) = b \Rightarrow (a \Rightarrow c)$ for all $a, b, c \in L$.
- (2) $a \rightarrow (b \rightarrow c) = b \rightarrow (a \rightarrow c)$ for all $a, b, c \in L$.
- (3) $a \odot b = b \odot a$ and $(a \odot b) \odot c = a \odot (b \odot c)$ for all $a, b, c \in L$.

Proof. (1) \Leftrightarrow (2). Since $a \Rightarrow (b \Rightarrow c) = \top$ iff $b \Rightarrow (a \Rightarrow c) = \top$, then $\Rightarrow = \rightarrow$.

Conversely, since (L, \rightarrow, \top) be an eo algebra having adjoint triple with \odot^* and \Rightarrow with $a \odot^* b = b \odot a$ from Theorem 2.7 (2), then (L, \rightarrow, \top) is commutative. Hence $\Rightarrow = \rightarrow$.

(1) \Leftrightarrow (3). By Theorem 2.9, $a \odot b = b \odot a$. By Theorems 3.1 and 3.2, $(a \odot b) \odot c = (a \odot c) \odot b = b \odot (a \odot c) = a \odot (b \odot c)$.

Conversely, by Theorem 3.1, it follows from $a \odot (b \odot c) = b \odot (a \odot c)$. \square

THEOREM 3.8. *Let (L, \Rightarrow, \top) be a w-eo algebra having $a \Rightarrow (b \Rightarrow c) = b \Rightarrow (a \Rightarrow c)$ for all $a, b, c \in L$ and $(a \Rightarrow \perp) \Rightarrow \perp = a$. Then $a \Rightarrow b = (b \Rightarrow \perp) \Rightarrow (a \Rightarrow \perp)$.*

Proof. It follows from $a \Rightarrow b = a \Rightarrow ((b \Rightarrow \perp) \Rightarrow \perp) = (b \Rightarrow \perp) \Rightarrow (a \Rightarrow \perp)$. \square

EXAMPLE 3.9. (1) Let $([0, 1], \Rightarrow)$ be a unit interval defined as

$$a \Rightarrow b = \begin{cases} 1, & \text{if } a \leq b, \\ (\frac{1}{2} - a) \vee b, & \text{otherwise.} \end{cases}$$

We easily show that $([0, 1], \Rightarrow, 1)$ is an eo algebra having $\bigvee_i x_i \Rightarrow y = \bigwedge_i (x_i \Rightarrow y)$ and $x \Rightarrow \bigwedge_i y_i = \bigwedge_i (x \Rightarrow y_i)$. Put $f_b(x) = x \Rightarrow b$. Define $g_b(y) = y \rightarrow b = \bigvee\{x \in L \mid y \leq f_b(x)\}$. Then $y \leq f_b(x)$ iff $x \leq g_b(y)$. We obtain:

$$a \rightarrow b = \begin{cases} 1, & \text{if } a \leq b, \\ (\frac{1}{2} - a) \vee b, & \text{otherwise.} \end{cases}$$

Put $f_b(x) = b \Rightarrow x$. Define $g_b(y) = b \odot y = \bigwedge\{x \in L \mid y \leq f_b(x)\}$. Then $g_b(y) \leq x$ iff $y \leq f_b(x)$. We obtain:

$$a \odot b = \begin{cases} a \wedge b, & \text{if } a + b \geq \frac{1}{2}, \\ 0, & \text{otherwise.} \end{cases}$$

Since $a \Rightarrow (b \Rightarrow c) = b \Rightarrow (a \Rightarrow c)$ for all $a, b, c \in L$, we have $a \odot b = b \odot a$ and $(a \odot b) \odot c = a \odot (b \odot c)$ for all $a, b, c \in L$. Since $(0.7 \Rightarrow \perp) \Rightarrow \perp = \top$, we have

$$0.7 = 0.8 \Rightarrow 0.7 \neq (0.7 \Rightarrow \perp) \Rightarrow (0.8 \Rightarrow \perp) = \perp \Rightarrow \perp = \perp.$$

EXAMPLE 3.10. Let $K = \{(x, y) \in \mathbb{R}^2 \mid x > 0\}$ be a set and we define an operation $\otimes : K \times K \rightarrow K$ as follows:

$$(x_1, y_1) \otimes (x_2, y_2) = (x_1 x_2, x_1 y_2 + y_1).$$

Then (K, \otimes) is a group with $e = (1, 0)$, $(x, y)^{-1} = (\frac{1}{x}, -\frac{y}{x})$.

We have a positive cone $P = \{(a, b) \in \mathbb{R}^2 \mid a = 1, b \geq 0, \text{ or } a > 1\}$ because $P \cap P^{-1} = \{(1, 0)\}$, $P \otimes P \subset P$, $(a, b)^{-1} \otimes P \otimes (a, b) = P$ and $P \cup P^{-1} = K$. For $(x_1, y_1), (x_2, y_2) \in K$, we define

$$\begin{aligned} (x_1, y_1) &\leq (x_2, y_2) \\ \Leftrightarrow (x_1, y_1)^{-1} \otimes (x_2, y_2) &\in P, (x_2, y_2) \otimes (x_1, y_1)^{-1} \in P \\ \Leftrightarrow x_1 < x_2 \text{ or } x_1 = x_2, y_1 &\leq y_2. \end{aligned}$$

Then (K, \leq, \otimes) is a lattice-group. (ref. [1])

The structure $(L, \odot, \Rightarrow, \rightarrow, (\frac{1}{2}, 1), (1, 0))$ is a generalized residuated lattice with strong negation where $\perp = (\frac{1}{2}, 1)$ is the least element and $\top = (1, 0)$ is the greatest element from the following statements:

$$\begin{aligned} (x_1, y_1) \odot (x_2, y_2) &= (x_1, y_1) \otimes (x_2, y_2) \vee (\frac{1}{2}, 1) \\ &= (x_1 x_2, x_1 y_2 + y_1) \vee (\frac{1}{2}, 1), \\ (x_1, y_1) \Rightarrow (x_2, y_2) &= ((x_1, y_1)^{-1} \otimes (x_2, y_2)) \wedge (1, 0) \\ &= (\frac{x_2}{x_1}, \frac{y_2 - y_1}{x_1}) \wedge (1, 0), \\ (x_1, y_1) \rightarrow (x_2, y_2) &= ((x_2, y_2) \otimes (x_1, y_1)^{-1}) \wedge (1, 0) \\ &= (\frac{x_2}{x_1}, -\frac{x_2 y_1}{x_1} + y_2) \wedge (1, 0). \end{aligned}$$

The structure $(L, \odot, \Rightarrow, \rightarrow, (\frac{1}{2}, 1), (1, 0))$ is a w-eo algebra having an adjoint triple as follows:

$$\begin{aligned} (x_1, y_1) \odot (x_2, y_2) \leq (x_3, y_3) &\text{ iff } (x_2, y_2) \leq (x_1, y_1) \Rightarrow (x_3, y_3) \\ &\text{ iff } (x_1, y_1) \leq (x_2, y_2) \rightarrow (x_3, y_3) \end{aligned}$$

Since

$$\begin{aligned}(x_1, y_1) \Rightarrow ((x_2, y_2) \Rightarrow (x_3, y_3)) &= \left(\frac{x_3}{x_1 x_2}, \frac{y_3 - x_2 y_1 - y_2}{x_1 x_2} \right) \wedge (1, 0) \\ &= ((x_2, y_2) \odot (x_1, y_1)) \Rightarrow (x_3, y_3),\end{aligned}$$

$(L, \odot, \Rightarrow, \rightarrow, (\frac{1}{2}, 1), (1, 0))$ is associative. Since $(1, 0) \Rightarrow (x, y) = (x, y)$ and

$$\begin{aligned}\left(\frac{4}{5}, 1\right) \Rightarrow \left(\left(\frac{5}{6}, 3\right) \Rightarrow \left(\frac{2}{3}, -1\right)\right) &= \left(\frac{4}{5}, 1\right) \Rightarrow \left(\frac{4}{5}, -\frac{24}{5}\right) = \left(1, -\frac{29}{4}\right), \\ \left(\frac{5}{6}, 3\right) \Rightarrow \left(\left(\frac{4}{5}, 1\right) \Rightarrow \left(\frac{2}{3}, -1\right)\right) &= \left(\frac{5}{6}, 3\right) \Rightarrow \left(\frac{5}{6}, -\frac{5}{2}\right) = \left(1, -\frac{33}{5}\right), \\ \left(\frac{4}{5}, 1\right) \Rightarrow \left(\left(\frac{5}{6}, 3\right) \Rightarrow \left(\frac{2}{3}, -1\right)\right) &\neq \left(\frac{5}{6}, 3\right) \Rightarrow \left(\left(\frac{4}{5}, 1\right) \Rightarrow \left(\frac{2}{3}, -1\right)\right),\end{aligned}$$

by Theorems 3.1, 3.2 and 3.7,

$$\begin{aligned}(x_1, y_1) \rightarrow ((x_2, y_2) \rightarrow (x_3, y_3)) &\neq (x_2, y_2) \rightarrow ((x_1, y_1) \rightarrow (x_3, y_3)) \\ (x_1, y_1) \odot ((x_2, y_2) \odot (x_3, y_3)) &\neq (x_2, y_2) \odot ((x_1, y_1) \odot (x_3, y_3)) \\ ((x_1, y_1) \odot (x_2, y_2)) \odot (x_3, y_3) &\neq ((x_1, y_1) \odot (x_3, y_3)) \odot (x_2, y_2) \\ (x_1, y_1) \odot (x_2, y_2) &\neq (x_2, y_2) \odot (x_1, y_1).\end{aligned}$$

References

- [1] G. Birkhoff, *Lattice Theory*, Amer. Math. Soc. Colloq. Publ., **25**, New York (1968)
- [2] C.C. Chang, *Algebraic analysis of many valued logics*, Trans. of A.M.S., **88** (2) (1958), 467–490
- [3] G. Georgescu, A. Popescu, *Non-commutative fuzzy Galois connections*, Soft Comput, **7** (2003), 458–467
- [4] G. Georgescu, A. Popescu, *Non-commutative fuzzy structures and pairs of weak negations*, Fuzzy Sets and Systems, **143** (2004), 129–155
- [5] C. Guido, P. Toto, *Extended ordered algebras*, J. Appl. Log., **6** (2008) 609–626
- [6] Y.C. Kim, J.M. Ko, *Some properties of extended order algebras*, Korean J. Math., **21** (1) (2013) 63–73
- [7] U. Höhle, E. P. Klement, *Non-classical logic and their applications to fuzzy subsets*, Kluwer Academic Publisher, Boston (1995)
- [8] E. Turunen, *Mathematics Behind Fuzzy Logic*, A Springer-Verlag Co. (1999)
- [9] R. Wille, *Restructuring lattice theory; an approach based on hierarchies of concept*, in: 1. Rival(Ed.), *Ordered Sets*, Reidel, Dordrecht, Boston (1982)

Department of Mathematics
Gangneung-Wonju National University
Gangneung 210-702, Korea
E-mail: yck@gwnu.ac.kr

Department of Mathematics
Gangneung-Wonju National University
Gangneung 210-702, Korea
E-mail: jmko@gwnu.ac.kr