

EINSTEIN'S CONNECTION IN 3-DIMENSIONAL *ES*-MANIFOLD

IN HO HWANG

ABSTRACT. The manifold $*g-ESX_n$ is a generalized n -dimensional Riemannian manifold on which the differential geometric structure is imposed by the unified field tensor $*g^{\lambda\nu}$ through the *ES*-connection which is both Einstein and semi-symmetric. The purpose of the present paper is to prove a necessary and sufficient condition for a unique Einstein's connection to exist in 3-dimensional $*g-ESX_3$ and to display a surveyable tensorial representation of 3-dimensional Einstein's connection in terms of the unified field tensor, employing the powerful recurrence relations in the first class.

1. Preliminaries

This paper is a direct continuation of our previous paper [1], which will be denoted by I in the present paper. All considerations in this paper are based on the results and symbolism of I . Whenever necessary, they will be quoted in the present paper. In this section, we introduce a brief collection of basic concepts, notations, and results of I , which are frequently used in the present paper ([2],[3],[4]).

Received April 13, 2015. Revised June 11, 2015. Accepted June 15, 2015.

2010 Mathematics Subject Classification: 83E50, 83C05, 58A05.

Key words and phrases: *ES*-manifold, Einstein's connection.

This research was supported by Incheon National University Research Grant, 2013-2014.

© The Kangwon-Kyungki Mathematical Society, 2015.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

(a) n -dimensional *g -unified field theory

Let X_n be an n -dimensional generalized Riemannian manifold referred to a real coordinate system x^ν , which obeys the coordinate transformations $x^\nu \rightarrow x^{\nu'}$ for which

$$(1.1) \quad \det\left(\frac{\partial x'}{\partial x}\right) \neq 0$$

In $n - g - UFT$ the manifold X_n is endowed with a real nonsymmetric tensor $g_{\lambda\mu}$, which may be decomposed into its symmetric part $h_{\lambda\mu}$ and skew-symmetric part $k_{\lambda\mu}$:

$$(1.2) \quad g_{\lambda\mu} = h_{\lambda\mu} + k_{\lambda\mu}$$

where

$$(1.3) \quad \mathfrak{g} = \det(g_{\lambda\mu}) \neq 0, \quad \mathfrak{h} = \det(h_{\lambda\mu}) \neq 0, \quad \mathfrak{k} = \det(k_{\lambda\mu})$$

In $n - {}^*g - UFT$ the algebraic structure on X_n is imposed by the basic real tensor ${}^*g^{\lambda\nu}$ defined by

$$(1.4) \quad g_{\lambda\mu} {}^*g^{\lambda\nu} = g_{\mu\lambda} {}^*g^{\nu\lambda} = \delta_\mu^\nu$$

It may be also decomposed into its symmetric part ${}^*h^{\lambda\nu}$ and skew-symmetric part ${}^*k^{\lambda\nu}$:

$$(1.5) \quad {}^*g^{\lambda\nu} = {}^*h^{\lambda\nu} + {}^*k^{\lambda\nu}$$

Since $\det({}^*h^{\lambda\nu}) \neq 0$, we may define a unique tensor ${}^*h_{\lambda\mu}$ by

$$(1.6) \quad {}^*h_{\lambda\mu} {}^*h^{\lambda\nu} = \delta_\mu^\nu$$

In $n - {}^*g - UFT$ we use both ${}^*h^{\lambda\nu}$ and ${}^*h_{\lambda\mu}$ as tensors for raising and/or lowering indices of all tensors in X_n in the usual manner. We then have

$$(1.7) \quad {}^*k_{\lambda\mu} = {}^*k^{\rho\sigma} {}^*h_{\lambda\rho} {}^*h_{\mu\sigma}, \quad {}^*g_{\lambda\mu} = {}^*g^{\rho\sigma} {}^*h_{\lambda\rho} {}^*h_{\mu\sigma}$$

so that

$$(1.8) \quad {}^*g_{\lambda\mu} = {}^*h_{\lambda\mu} + {}^*k_{\lambda\mu}$$

The differential geometric structure on X_n is imposed by the tensor ${}^*g^{\lambda\nu}$ by means of a connection $\Gamma_{\lambda\ \mu}^\nu$ defined by a system of equations

$$(1.9) \quad D_\omega {}^*g^{\lambda\nu} = -2S_{\omega\alpha}{}^\nu {}^*g^{\lambda\alpha}$$

where D_ω denotes the symbol of the covariant derivative with respect to $\Gamma_{\lambda^\nu \mu}$ and $S_{\lambda\mu}{}^\nu$ is the torsion tensor of $\Gamma_{\lambda^\nu \mu}$. Under certain conditions the system (1.9) admits a unique solutions $\Gamma_{\lambda^\nu \mu}$.

It has been shown in [5] that if the system (1.9) admits $\Gamma_{\lambda^\nu \mu}$, it must be of the form

$$(1.10) \quad \Gamma_{\lambda^\nu \mu} = * \left\{ \begin{array}{c} \nu \\ \lambda\mu \end{array} \right\} + U^\nu{}_{\lambda\mu} + S_{\lambda\mu}{}^\nu.$$

where

$$(1.11) \quad U_{\nu\lambda\mu} = S_{(\lambda\mu)\nu}^{100} + 2 S_{\nu(\lambda\mu)}^{(10)0}$$

(b) Some notations and results

The following quantities are frequently used in our further considerations:

$$(1.12) \quad *g = \det(*g_{\lambda\mu}), \quad *h = \det(*h_{\lambda\mu}), \quad *k = \det(*k_{\lambda\mu})$$

$$(1.13) \quad *g = \frac{*g}{*h}, \quad *k = \frac{*k}{*h}.$$

$$(1.14) \quad K_p = *k_{[\alpha_1}{}^{\alpha_1} *k_{\alpha_2}{}^{\alpha_2} \dots *k_{\alpha_p]}{}^{\alpha_p}, \quad (p = 0, 1, 2, \dots).$$

$$(1.15) \quad {}^{(0)}*k_\lambda{}^\nu = \delta_\lambda^\nu, \quad {}^{(p)}*k_\lambda{}^\nu = *k_\lambda{}^\alpha {}^{(p-1)}*k_\alpha{}^\nu \quad (p = 1, 2, \dots).$$

$$(1.16) \quad K_{\omega\mu\nu} = \nabla_\nu *k_{\omega\mu} + \nabla_\omega *k_{\nu\mu} + \nabla_\mu *k_{\omega\nu}$$

where ∇_ω is the symbolic vector of the covariant derivative with respect to the christoffel symbols $* \left\{ \begin{array}{c} \nu \\ \lambda\mu \end{array} \right\}$ defined by $*h_{\lambda\mu}$ in the usual way.

In X_n it was proved in [5] that

$$(1.17) \quad K_0 = 1, \quad K_n = *k \text{ if } n \text{ is even, and } K_n = 0 \text{ if } n \text{ is odd.}$$

$$(1.18) \quad *g = 1 + K_2 + \dots + K_{n-\sigma}.$$

$$(1.19) \quad \sum_{s=0}^{n-\sigma} K_s {}^{(n-s)}*k_\lambda{}^\nu = 0 \quad (p = 0, 1, 2, \dots).$$

We also use the following useful abbreviations, denoting an arbitrary tensor $T_{\omega\mu\nu}$ skew-symmetric in the first two indices by T :

$$(1.20) \quad T = T_{\omega\mu\lambda}^{pqr} = T_{\alpha\beta\gamma}^{(p)*} k_{\omega}^{\alpha(q)*} k_{\mu}^{\beta(r)*} k_{\lambda}^{\gamma}$$

and for an arbitrary tensor T^{\dots} for $p = 1, 2, 3, \dots$:

$$(1.21) \quad {}^{(p)}T^{\dots} = {}^{(p-1)*} k_{\alpha}^{\nu} T^{\alpha\dots}$$

On the other hand, it has shown in [6] that the tensor $S_{\lambda\mu}{}^{\nu}$ satisfies

$$(1.22) \quad S = B - 3 \overset{(110)}{S}$$

where

$$(1.23) \quad 2B_{\omega\mu\nu} = K_{\omega\mu\nu} + 3K_{\alpha[\mu\beta}{}^* k_{\omega]}^{\alpha*} k_{\nu}^{\beta}$$

In our subsequent chapter, we start with the relation (1.22) to solve the system (1.9). Furthermore, for the first class, the nonholonomic solution of (1.22) may be given by

$$(1.24) \quad M S_{xyz} = B_{xyz}$$

or equivalently

$$(1.25) \quad 4M S_{xyz} = (2 + \underset{z}{M} \underset{x}{M} + \underset{z}{M} \underset{y}{M}) K_{xyz} + \underset{z}{M} (\underset{x}{M} + \underset{z}{M}) K_{zxy} + \underset{z}{M} (\underset{y}{M} + \underset{z}{M}) K_{yzx}$$

where

$$(1.26) \quad M = 1 + \underset{xyz}{M} + \underset{x}{M} \underset{y}{M} + \underset{y}{M} \underset{z}{M} + \underset{z}{M} \underset{x}{M}$$

Therefore, in virtue of (1.24), we see that a necessary and sufficient condition for the system (1.9) to have a unique solution in the first class is

$$(1.27) \quad \underset{xyz}{M} \neq 0 \text{ for all } x, y, z$$

(c) n -dimensional ES manifold $n - {}^*g$ -UFT

In this subsection, we display an useful representation of the ES connection in $n - {}^*g$ -UFT.

DEFINITION 1.1. A connection $\Gamma_{\lambda}^{\nu}{}_{\mu}$ is said to be *semi-symmetric* if its torsion tensor $S_{\lambda\mu}{}^{\nu}$ is of the form

$$(1.28) \quad S_{\lambda\mu}{}^{\nu} = 2\delta_{[\lambda}^{\nu} X_{\mu]}.$$

for an arbitrary non-null vector X_{μ} .

A connection which is both semi-symmetric and Einstein is called an *ES* connection. An n -dimensional generalized Riemannian manifold X_n , on which the differential geometric structure is imposed by $*g^{\lambda\nu}$ by means of an *ES* connection, is called an n -dimensional $*g$ -*ES* manifold. We denote this manifold by $*g$ -*ESX* $_n$ in our further considerations.

In $*g$ -*ESX* $_n$, the following theorems were proved in I.

THEOREM 1.2. *The main recurrence relation in the first class is*

$$(1.29) \quad {}^{(p+3)*}k_{\lambda}{}^{\nu} = -K_2 {}^{(p+1)*}k_{\lambda}{}^{\nu}, \quad (p = 0, 1, 2, \dots)$$

THEOREM 1.3. *The basic scalars M satisfy*

$$(1.30) \quad MM(M + M) = 0, \quad (x \neq y)$$

$$(1.31) \quad MM(MM - K_2) = 0, \quad (x \neq y)$$

THEOREM 1.4. (*Recurrence relations in the first class*) *If $T_{\omega\mu\nu}$ is a tensor skew-symmetric in the first two indices, then the following recurrence relations hold in the first class of 3- $*g$ -*ESX* $_3$:*

$$(1.32) \quad {}^{(12)r}T = 0, \quad {}^{22r}T = K_2 {}^{11r}T$$

$$(1.33) \quad T{}^{\nu[\omega\mu]}{}_{\nu[\omega\mu]} = 0, \quad T{}^{\nu[\omega\mu]}{}_{\nu[\omega\mu]} = K_2 T{}^{\nu[\omega\mu]}{}_{\nu[\omega\mu]}$$

2. Einstein's connection $\Gamma_{\lambda}^{\nu}{}_{\mu}$ in the first class

In this section, we shall derive surveyable tensorial representations of $S_{\lambda\mu}{}^{\nu}$ and hence $\Gamma_{\lambda}^{\nu}{}_{\mu}$ in terms of $*g^{\lambda\nu}$, employing the recurrence relations.

In the following theorem, we shall prove two relations in X_n . These relations will be used in our subsequent theorem when we are concerned with the solution of (1.9).

THEOREM 2.1. *We have*

$$(2.1) \quad \begin{matrix} (pq)r & (pq)r & (p'q')r & (p'q)r' & (pq)r' \\ B & = S & + S & + S & + S \end{matrix}$$

$$(2.2) \quad \begin{matrix} (pq)r & & (pq)r & & r''(pq) \\ 2 B_{\omega\mu\nu} & & = K_{\omega\mu\nu} & + & K_{\nu[\omega\mu]} \\ & + \frac{1}{2} (& (pq')r' & & (p'q)r' & & r'p'q & & r'q'p \\ & & K_{\omega\mu\nu} & + & K_{\omega\mu\nu} & + & K_{\nu[\omega\mu]} & + & K_{\nu[\omega\mu]}) \end{matrix}$$

where

$$(2.3) \quad p' = p + 1, \quad q' = q + 1, \quad r' = r + 1, \quad r'' = r + 2$$

Proof. In virtue of (1.22) and (1.20), the first relation (2.1) is obtained as in the following way:

$$(2.4) \quad \begin{matrix} (pq)r & & (pq)r \\ B & = & B_{\omega\mu\nu} = \frac{1}{2} B_{\omega\beta\gamma} ((p)^* k_{\omega}^{\alpha} (q)^* k_{\mu}^{\beta} + (q)^* k_{\omega}^{\alpha} (p)^* k_{\mu}^{\beta}) (r)^* k_{\nu}^{\gamma} \\ & = & \frac{1}{2} (S_{\alpha\beta\gamma} + S_{\epsilon\eta\gamma}^* k_{\alpha}^{\epsilon} k_{\beta}^{\eta} + S_{\epsilon\beta\eta}^* k_{\alpha}^{\epsilon} k_{\gamma}^{\eta} + S_{\alpha\epsilon\eta}^* k_{\beta}^{\epsilon} k_{\gamma}^{\eta}) \\ & & \times ((p)^* k_{\omega}^{\alpha} (q)^* k_{\mu}^{\beta} + (q)^* k_{\omega}^{\alpha} (p)^* k_{\mu}^{\beta}) (r)^* k_{\nu}^{\gamma} \end{matrix}$$

After a lengthy calculation, we note that the right-hand side of the above equation is equal to that of (2.1). Similarly, we verify (2.2) using (1.20) and (1.23). \square

THEOREM 2.2. *A necessary and sufficient condition for the system (1.9) to admit a unique solution $\Gamma_{\lambda}^{\nu\mu}$ is that*

$$(2.5) \quad 1 - (K_2)^2 \neq 0$$

Proof. Since M defined by (1.26), is symmetric in x, y, z and satisfies

$$(2.6) \quad \begin{matrix} M & & M & & M \\ 33x & = & 1, & 311 & = & M & = & 1 - K_2, & 123 & = & M & = & M & = & 1 + K_2 \\ & & & 322 & & & & & 112 & & 122 \end{matrix}$$

we have the condition (2.5) in virtue of (1.27). \square

THEOREM 2.3. *The system of equations (1.22) is reduced to a system of the following 5 equations:*

$$(2.7) \quad \begin{cases} B = S + 2 \overset{(10)1}{S} + \overset{110}{S} \\ \overset{(10)1}{B} = \overset{(10)1}{S} + \overset{(20)2}{S} + \overset{112}{S} \\ \overset{110}{B} = (1 + K_2) \overset{110}{S} \\ \overset{(20)2}{B} = (K_2)^2 \overset{(10)1}{S} + \overset{(20)2}{S} - K_2 \overset{112}{S} \\ \overset{112}{B} = (1 + K_2) \overset{112}{S} \end{cases}$$

Proof. This assertion follows from (2.1) using (1.29), (1.32) and (1.33). \square

THEOREM 2.4. The tensor $\overset{(pq)r}{B}_{\omega\mu\nu}$ are given as linear combinations of $\overset{(pq)r}{K}_{\omega\mu\nu}$, as follows:

$$(2.8) \quad \begin{cases} 2 \overset{(10)1}{B}_{\omega\mu\nu} = \overset{(10)1}{K}_{\omega\mu\nu} + \frac{1}{2}(\overset{112}{K}_{\omega\mu\nu} + \overset{(20)2}{K}_{\omega\mu\nu} + \overset{211}{K}_{\nu[\omega\mu]} - K_2 \overset{101}{K}_{\nu[\omega\mu]}) \\ 2 \overset{110}{B}_{\omega\mu\nu} = \overset{110}{K}_{\omega\mu\nu} \\ 2 \overset{(20)2}{B}_{\omega\mu\nu} = \overset{(20)2}{K}_{\omega\mu\nu} + \frac{1}{2}[(K_2)^2 \overset{(10)1}{K}_{\omega\mu\nu} - K_2 \overset{112}{K}_{\nu[\omega\mu]} - K_2 \overset{202}{K}_{\nu[\omega\mu]}] \\ 2 \overset{112}{B}_{\omega\mu\nu} = \overset{112}{K}_{\omega\mu\nu} \end{cases}$$

Proof. These relations are obtained from (2.2) in virtue of (1.29), (1.32) and (1.33). \square

THEOREM 2.5. If the condition (2.5) is satisfied, the unique solution of (1.22) is given by

$$(2.9) \quad [1 - (K_2)^2](S - B) = -2 \overset{(10)1}{B} + (K_2 - 1) \overset{110}{B} + 2 \overset{(20)2}{B} + 2 \overset{112}{B}$$

or equivalently

$$\begin{aligned}
 (2.10) \quad & [1 - (K_2)^2](2S_{\omega\mu\nu} - K_{\omega\mu\nu} - \overset{110}{K}_{\nu[\omega\mu]} - \overset{200}{K}_{\nu[\omega\mu]}) = \\
 & - \overset{(10)1}{K}_{\omega\mu\nu} + \overset{112}{K}_{\omega\mu\nu} + \overset{(20)2}{K}_{\omega\mu\nu} - \overset{211}{K}_{\nu[\omega\mu]} \\
 & + (K_2 - 1)\overset{110}{K}_{\omega\mu\nu} + K_2(\overset{101}{K}_{\nu[\omega\mu]} - \overset{112}{K}_{\nu[\omega\mu]} - \overset{202}{K}_{\nu[\omega\mu]})
 \end{aligned}$$

Proof. (2.9) is the solution of (2.7), while (2.10) is obtained by substituting (2.8) into (2.9) and making use of recurrence relations. \square

THEOREM 2.6. *The tensor $U^\nu_{\lambda\mu}$ is given by*

$$\begin{aligned}
 (2.11) \quad & [1 - (K_2)^2](U_{\nu\lambda\mu} - \overset{[10]0}{B}_{\lambda\mu\nu} - 2 \overset{(10)0}{B}_{\nu(\lambda\mu)}) = \\
 & - K_2(\overset{[10]2}{B}_{\lambda\mu\nu} + 2 \overset{(10)2}{B}_{\nu(\lambda\mu)}) + (K_2 - 1) \overset{[21]0}{B}_{\lambda\mu\nu} \\
 & + \overset{[02]1}{B}_{\lambda\mu\nu} + \overset{[21]2}{B}_{\lambda\mu\nu} - 2 \overset{(20)1}{B}_{\nu(\lambda\mu)} - 2 \overset{111}{B}_{\nu(\lambda\mu)}
 \end{aligned}$$

or equivalently

$$\begin{aligned}
 (2.12) \quad & [1 - (K_2)^2](2U_{\nu\lambda\mu} + \overset{[01]0}{K}_{\lambda\mu\nu} - 2 \overset{(10)0}{K}_{\nu(\lambda\mu)}) = (K_2 - 1) \overset{[21]0}{K}_{\lambda\mu\nu} \\
 & + \overset{[02]1}{K}_{\lambda\mu\nu} + K_2 \overset{[01]2}{K}_{\lambda\mu\nu} - 2(K_2 \overset{(10)2}{K}_{\nu(\lambda\mu)} - \overset{(20)1}{K}_{\nu(\lambda\mu)} - \overset{111}{K}_{\nu(\lambda\mu)})
 \end{aligned}$$

Proof. The representations (2.11), (2.12) are direct consequences of substituting (2.9), (2.10) into (1.11). \square

Now that we have obtained the tensor $S_{\lambda\mu}{}^\nu$ and $U^\nu_{\lambda\mu}$ in terms of ${}^*g^{\lambda\nu}$, it is possible for us to determine $\Gamma_{\lambda}{}^\nu{}_\mu$ by only substituting for S and U into (1.10).

References

- [1] Hwang, I.H., *On the algebra of 3-dimensional ES-manifold*, Korean J. Math. **22** (1) (2014), 207–216.
- [2] Datta, D.k., *Some theorems on symmetric recurrent tensors of the second order*, Tensor (N.S.) **15** (1964), 1105–1136.
- [3] Einstein, A., *The meaning of relativity*, Princeton University Press, 1950.

- [4] Mishra, R.S., *n-dimensional considerations of unified field theory of relativity*, Tensor **9** (1959), 217–225.
- [5] Chung, K.T., *Einstein's connection in terms of $*g^{\lambda\nu}$* , Nuovo cimento Soc. Ital. Fis. B, **27** (1963), (X), 1297–1324
- [6] Hlavatý, V., *Geometry of Einstein's unified field theory*, Noordhoop Ltd., 1957

In Ho Hwang
Department of Mathematics
Incheon National University
Incheon 406-772, Korea
E-mail: ho818@incheon.ac.kr