

## ON LEFT DERIVATIONS OF *BCH*-ALGEBRAS

KYUNG HO KIM AND YONG HOON LEE\*

ABSTRACT. In this paper, we introduce the notion of left derivations of *BCH* algebras and investigate some properties of left derivations in a *BCH*-algebra. Moreover, we introduce the notions of fixed set and kernel set of derivations in a *BCH*-algebra and obtained some interesting properties in medial *BCH*-algebras. Also, we discuss the relations between ideals in a medial *BCH*-algebras.

### 1. Introduction

In 1966, Imai and Iseki introduced two classes of abstract algebras, *BCK*-algebra and *BCI*-algebras [6]. It is known that the class of *BCI*-algebras is a generalization of the class of *BCK*-algebras. In 1983, Hu and Li [3] introduced the notion of a *BCH*-algebra, which is a generalization of the notions of *BCK*-algebras and *BCI*-algebras. They have studied a few properties of these algebras. In this paper, we introduce the notion of left derivations of *BCH* algebras and investigate some properties of left derivations in a *BCH*-algebra. Moreover, we introduce the notions of fixed set and kernel set of derivations in a *BCH*-algebra and obtained some interesting properties in medial *BCH*-algebras. Also, we discuss the relations between ideals in a medial *BCH*-algebras.

---

Received November 4, 2016. Revised April 24, 2017. Accepted April 26, 2017.

2010 Mathematics Subject Classification: 03G25, 06B10, 06D99, 06B35, 06B99.

Key words and phrases: *BCH*-algebra, derivation, left derivation, isotone, fixed set, medial.

\* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2017.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

## 2. Preliminary

By a *BCH-algebra*, we mean an algebra  $(X, *, 0)$  with a single binary operation “ $*$ ” that satisfies the following identities, for any  $x, y, z \in X$ ,

$$(BCH1) \quad x * x = 0,$$

$$(BCH2) \quad x \leq y \text{ and } y \leq x \text{ imply } x = y, \text{ where } x \leq y \text{ if and only if } x * y = 0.$$

$$(BCH3) \quad (x * y) * z = (x * z) * y.$$

In a *BCH-algebra*  $X$ , the following identities are true, for all  $x, y \in X$ ,

$$(BCH4) \quad (x * (x * y)) * y = 0,$$

$$(BCH5) \quad x * 0 = 0 \text{ implies } x = 0,$$

$$(BCH6) \quad 0 * (x * y) = (0 * x) * (0 * y),$$

$$(BCH7) \quad x * 0 = x,$$

$$(BCH8) \quad (x * y) * x = 0 * y,$$

$$(BCH9) \quad x * y = 0 \text{ implies } 0 * x = 0 * y,$$

$$(BCH10) \quad x * (x * y) \leq y.$$

DEFINITION 2.1. Let  $X$  be a *BCH-algebra*. we define a partial order  $\leq$  by putting  $x \leq y$  if and only if  $x * y = 0$ , for every  $x, y \in X$ .

In a *BCH-algebra*  $X$ , the following identity hold:

$$(BCH11) \quad x \leq y \text{ implies } x * z \leq y * z \text{ but } x \leq y \text{ implies } z * y \leq z * x \text{ does not hold.}$$

DEFINITION 2.2. Let  $I$  be a nonempty subset of a *BCH-algebra*  $X$ . Then  $I$  is called an *ideal* of  $X$  if it satisfies:

$$(1) \quad 0 \in I,$$

$$(2) \quad x * y \in I \text{ and } y \in I \text{ imply } x \in I.$$

DEFINITION 2.3. A *BCH-algebra* is said to be *medial* if it satisfies

$$(x * y) * (z * w) = (x * z) * (y * w)$$

for all  $x, y, z, w \in X$ .

In a medial *BCH-algebra*  $X$ , the following identity hold:

$$(BCH12) \quad x * (x * y) = y, \text{ for all } x, y \in X.$$

DEFINITION 2.4. Let  $X$  be a *BCH-algebra*. Then the set  $X_+ = \{x \in X \mid 0 * x = 0\}$  is called a *BCH-part* of  $X$ .

DEFINITION 2.5. A *BCH-algebra*  $X$  is said to be *commutative* if for all  $x, y \in X$ ,  $y * (y * x) = x * (x * y)$ , i.e.,  $x \wedge y = y \wedge x$ . For a *BCH-algebra*  $X$ , we denote  $x \wedge y = y * (y * x)$ , for all  $x, y \in X$ .

DEFINITION 2.6. Let  $X$  be a *BCH*-algebra. A map  $d : X \rightarrow X$  is a *left-right derivation* (briefly, *(l, r)-derivation*) of  $X$  if it satisfies the identity

$$d(x * y) = (d(x) * y) \wedge (x * d(y))$$

for all  $x, y \in X$ . If  $d$  satisfies the identity

$$d(x * y) = (x * d(y)) \wedge (d(x) * y)$$

for all  $x, y \in X$ , then  $d$  is a *right-left derivation* (briefly, *(r, l)-derivation*) of  $X$ . Moreover, if  $d$  is both an *(l, r)* and *(r, l)*-derivation of  $X$ , then  $d$  is a *derivation* of  $X$ .

### 3. Left derivations of *BCH*-algebras

In what follows, let  $X$  denote a *BCH*-algebra unless otherwise specified.

DEFINITION 3.1. Let  $X$  be a *BCH*-algebra. By a *left derivation* of  $X$ , we mean a self-map  $d$  satisfying

$$d(x * y) = (x * d(y)) \wedge (y * d(x))$$

for all  $x, y \in X$ .

EXAMPLE 3.2. Let  $X = \{0, 1, 2\}$  be a *BCH*-algebra with Cayley table as follows:

$*$	0	1	2
0	0	0	2
1	1	0	2
2	2	2	0

Define a self-map  $d : X \rightarrow X$  by

$$d(x) = \begin{cases} 2 & \text{if } x = 0, 1 \\ 0 & \text{if } x = 2 \end{cases}$$

Then it is easy to check that  $d$  is a left derivation of a *BCH*-algebra  $X$ .

EXAMPLE 3.3. Let  $X = \{0, 1, 2, 3\}$  be a *BCH*-algebra with Cayley table as follows:

*	0	1	2	3
0	0	1	2	3
1	1	0	3	2
2	2	3	0	1
3	3	2	1	0

Define a self-map  $d : X \rightarrow X$  by

$$d(x) = \begin{cases} 0 & \text{if } x = 3 \\ 1 & \text{if } x = 2 \\ 2 & \text{if } x = 1 \\ 3 & \text{if } x = 0 \end{cases}$$

Then it is easy to check that  $d$  is a left derivation of a *BCH*-algebra  $X$ .

**DEFINITION 3.4.** A self-map  $d$  of a *BCH*-algebra  $X$  is said to be *regular* if  $d(0) = 0$ .

**PROPOSITION 3.5.** Let  $d$  be a left derivation of  $X$ . If  $0 * x = 0$ , for every  $x \in X$ , then  $d$  is regular.

*Proof.* Let  $0 * x = 0$ , for all  $x \in X$ . Then we have

$$\begin{aligned} d(0) &= d(0 * x) = (0 * d(x)) \wedge (x * d(0)) \\ &= 0 \wedge (x * d(0)) = (x * d(0)) * ((x * d(0)) * 0) \\ &= (x * d(0)) * ((x * d(0)) * 0) = 0 \end{aligned}$$

Hence  $d$  is regular. □

**PROPOSITION 3.6.** Let  $d$  be a left derivation of  $X$ . If there exists  $a \in X$  such that  $a * d(x) = 0$ , for all  $x \in X$ , then  $d$  is regular.

*Proof.* Let  $a * d(x) = 0$  for all  $x \in X$ . Then

$$\begin{aligned} 0 &= a * d(a * x) = a * ((a * d(x)) \wedge (x * d(a))) \\ &= a * (0 \wedge (x * d(a))) \\ &= a * ((x * d(a)) * ((x * d(a)) * 0)) \\ &= a * 0 \\ &= a. \end{aligned}$$

Hence we have

$$\begin{aligned}
 d(0) &= d(a) \\
 &= d(a * 0) \\
 &= (a * d(0)) \wedge (0 * d(a)) \\
 &= 0 \wedge (0 * d(a)) \\
 &= 0.
 \end{aligned}$$

Hence  $d$  is regular.  $\square$

**PROPOSITION 3.7.** *Let  $d$  be a regular left derivation of  $X$ . Then, for all  $x \in X$ ,*

- (1)  $x \leq d(x)$ ,
- (2)  $x \leq d(d(x))$ .

*Proof.* (1) Let  $d$  be a regular left derivation of  $X$ . Then we have

$$\begin{aligned}
 0 &= d(x * x) = (x * d(x)) \wedge (x * d(x)) \\
 &= (x * d(x)) * ((x * d(x)) * (x * d(x))) = (x * d(x)) * 0 \\
 &= x * d(x),
 \end{aligned}$$

which implies  $x \leq d(x)$ , for all  $x \in X$ .

- (2) From (1),  $x \leq d(x) \leq d(d(x))$ , for all  $x \in X$ .  $\square$

**THEOREM 3.8.** *Let  $d$  be a left derivation of a medial *BCH*-algebra  $X$ . Then  $d(x * y) = x * d(y)$ , for all  $x, y \in X$ .*

*Proof.* Let  $x, y \in X$ . Then we have

$$d(x * y) = (x * d(y)) \wedge (y * d(x)) = (y * d(x)) * ((y * d(x)) * (x * d(y))) = x * d(y).$$

$\square$

**PROPOSITION 3.9.** *Let  $X$  be a *BCH*-algebra. Then*

$$d_n(d_{n-1}(\dots(d_2(d_1(x)))) \leq x$$

for  $n \in \mathbb{N}$ , where  $d_1, d_2, \dots, d_n$  are regular left derivations of  $X$ .

*Proof.* For  $n = 1$ ,

$$d_1(x) = d_1(x * 0) = (x * d_1(0)) \wedge (0 * d_1(x)) = x \wedge (0 * d_1(x)) \leq x,$$

by (BCH10). Hence we have  $d_1(x) \leq x$ .  $\square$

Let  $n \in \mathbb{N}$  and  $d_n(d_{n-1}(\dots(d_2(d_1(x)))) \leq x$ . For simplicity, let

$$D_n = d_n(d_{n-1}(\dots(d_2(d_1(x))))).$$

Then

$$\begin{aligned} d_{n+1}(D_n) &= d_{n+1}(D_n * 0) = (D_n * d_{n+1}(0)) \wedge (0 * d_{n+1}(D_n)) \\ &= (0 * d_{n+1}(D_n)) * ((0 * d_{n+1}(D_n)) * D_n) \leq D_n. \end{aligned}$$

Hence  $d_{n+1}(D_n) \leq D_n$ , that is,  $D_{n+1} \leq D_n \leq x$  by assumption, which implies  $D_{n+1} \leq x$ .

**THEOREM 3.10.** *Let  $X$  be a medial BCH-algebra and let  $d$  be a left derivation of  $X$ . Then  $d$  is regular if and only if  $d(x) = x$ , for all  $x \in X$ .*

*Proof.* Let  $d$  be regular. Then we have  $d(0) = 0$ . Hence,

$$\begin{aligned} d(x) &= d(x * 0) \\ &= (x * d(0)) \wedge (0 * d(x)) \\ &= (x * 0) \wedge (0 * d(x)) \\ &= (0 * d(x)) * ((0 * d(x)) * x) \\ &= x. \end{aligned}$$

Hence  $d(x) = x$  for  $x \in X$ . Conversely, assume that  $d(x) = x$  for all  $x \in X$ . Then it is clear that  $d(0) = 0$ . This implies that  $d$  is regular.  $\square$

**PROPOSITION 3.11.** *Let  $d$  be a left derivation of  $X$ . Then we have*

- (1)  $x * d(x) = y * d(y)$ ,
- (2)  $d(x * y) \leq x * d(y)$ ,
- (3)  $d(d(x) * x) = 0$ , for every  $x, y \in X$ .

*Proof.* (1) Since  $x * x = 0$  for all  $x \in X$ , we have  $d(0) = d(x * x) = (x * d(x)) \wedge (x * d(x)) = x * d(x)$ . Similarly,  $d(0) = y * d(y)$ , which implies  $x * d(x) = y * d(y)$ .

(2) Let  $d$  be a left derivation of a BCH-algebra of  $X$  and  $x, y \in X$ . Then

$$\begin{aligned} d(x * y) &= (x * d(y)) \wedge (y * d(x)) \\ &= (y * d(x)) * ((y * d(x)) * (x * (d(y)))) \\ &\leq x * d(y). \end{aligned}$$

(3) Let  $d$  be a left derivation of a BCH-algebra of  $X$  and  $x \in X$ . Then

$$d(d(x) * x) = (d(x) * d(x)) \wedge (x * d(d(x))) = 0 \wedge (x * d(d(x))) = 0.$$

□

**PROPOSITION 3.12.** *Let  $d$  be a regular left derivation of  $X$ . Then  $d : X \rightarrow X$  is an identity map if it satisfies  $d(x) * y = x * d(y)$ , for all  $x, y \in X$ .*

*Proof.* Since  $d$  is regular, we have  $d(0) = 0$ . Let  $x * d(y) = d(x) * y$  for all  $x, y \in X$ . Then  $d(x) = d(x) * 0 = x * d(0) = x * 0 = x$ . Thus  $d$  is an identity map. □

**PROPOSITION 3.13.** *Let  $X$  be a medial *BCH*-algebra  $X$  and let  $d$  be a regular left derivation of  $X$ . Define  $d^2(x) = d(d(x))$ , for all  $x \in X$ . Then  $d^2 = d$ .*

*Proof.* Let  $X$  be a medial *BCH*-algebra  $X$  and let  $d$  be a regular left derivation of  $X$ . Then we have for all  $x \in X$ ,

$$\begin{aligned} d^2(x) &= d(d(x)) = d(d(x * 0)) \\ &= (d(x) * d(0)) \wedge (d(0) * d(d(x))) \\ &= (d(x) * 0) \wedge (0 * d(d(x))) \\ &= d(x) \wedge (0 * d(d(x))) \\ &= (0 * d(d(x))) * [(0 * d(d(x))) * d(x)] = d(x). \end{aligned}$$

□

**DEFINITION 3.14.** Let  $X$  be a *BCH*-algebra. A self-map  $d$  on  $X$  is said to be *isotone* if  $x \leq y$  implies  $d(x) \leq d(y)$ , for  $x, y \in X$ .

**PROPOSITION 3.15.** *Let  $X$  be a medial commutative *BCH*-algebra and let  $d$  be a regular left derivation of  $X$ . Then the following properties are equivalent:*

- (1)  $d$  is an isotone derivation of  $X$ ,
- (2)  $x \leq y$  implies  $d(x * y) = d(x) * y$ .

*Proof.* (1)  $\Rightarrow$  (2). Let  $x, y \in X$  such that  $x \leq y$ . Then we have  $d(x * y) = d(x * y) = d(0) = 0$ . Since  $d$  is isotone, we get  $d(x) * d(y) = 0$ . Thus

$$\begin{aligned} d(x * y) &= 0 = d(x) * d(y) \\ &= d(x) * (d(y) \wedge y) \\ &= d(x) * (y * (y * d(y))) \\ &= d(x) * (d(y) * (d(y) * y)) \\ &= d(x) * y. \end{aligned}$$

(2)  $\Rightarrow$  (1). Let  $x, y \in X$  such that  $x \leq y$ . Thus

$$\begin{aligned} d(x) * d(y) &= d(x) * (d(y) \wedge y) \\ &= d(x) * (y * (y * d(y))) \\ &= d(x) * (d(y) * (d(y) * y)) \\ &= d(x) * y = d(x * y) \\ &= d(0) = 0. \end{aligned}$$

Hence  $d(x) \leq d(y)$ , which implies that  $d$  is an isotone derivation of  $X$ .  $\square$

**PROPOSITION 3.16.** *Let  $X$  be a medial commutative BCH-algebra and let  $d$  be a regular left derivation of  $X$ . Then the following properties are equivalent:*

- (1)  $d(x * y) = d(x) * y$ ,
- (2)  $d(x * y) = d(x) * d(y)$ , for all  $x, y \in X$ .

*Proof.* (1)  $\Rightarrow$  (2). Let  $X$  be a medial commutative BCH-algebra and let  $d$  be a regular left derivation of  $X$ . Then we have

$$\begin{aligned} d(x * y) &= d(x) * y = d(x) * (y \wedge d(y)) = d(x) * ((dy) * (d(y) * y)) \\ &= d(x) * (y * (y * d(y))) = d(x) * d(y). \end{aligned}$$

Hence  $d(x * y) = d(x) * d(y)$ .

(2)  $\Rightarrow$  (1). Let  $X$  be a medial BCH-algebra and let  $d$  be a regular left derivation of  $X$ . Then we have

$$\begin{aligned} d(x * y) &= d(x) * d(y) \\ &= d(x) * (d(y) \wedge y) \\ &= d(x) * (y \wedge d(y)) \\ &= d(x) * y, \end{aligned}$$

which implies  $d(x * y) = d(x) * y$ .  $\square$

From the Proposition 3.15 and 3.16, we have the following theorem.

**THEOREM 3.17.** *Let  $X$  be a medial commutative BCH-algebra and let  $d$  be a regular left derivation of  $X$ . Then the following properties are equivalent:*

- (1)  $d$  is an isotone derivation of  $X$ ,
- (2)  $x \leq y$  implies  $d(x * y) = d(x) * y$ ,
- (3)  $d(x * y) = d(x) * d(y)$ , for all  $x, y \in X$ .

**THEOREM 3.18.** *Let  $X$  be a medial commutative *BCH*-algebra and let  $d$  be a regular left derivation of  $X$ . Then the following properties are equivalent:*

- (1)  $d$  is an isotone derivation of  $X$ ,
- (2)  $d(x * y) = d(x) * d(y)$ ,
- (3)  $d(x \wedge y) = d(x) \wedge d(y)$ , for all  $x, y \in X$ .

*Proof.* (1)  $\Rightarrow$  (2). It follows from Theorem 3.17.

(2)  $\Rightarrow$  (3). Let  $d(x * y) = d(x) * d(y)$ , for all  $x, y \in X$ . Then we have

$$\begin{aligned} d(x \wedge y) &= d(y * (y * x)) \\ &= d(y) * d(y * x) \\ &= d(y) * (d(y) * d(x)) \\ &= d(x) \wedge d(y). \end{aligned}$$

(3)  $\Rightarrow$  (1). Let  $d(x \wedge y) = d(x) \wedge d(y)$  and  $x \leq y$ . Then  $x * y = 0$ , for all  $x, y \in X$ . Hence we have

$$\begin{aligned} d(x) &= d(x * 0) \\ &= d(x * (x * y)) \\ &= d(y \wedge x) \\ &= d(x) * (d(x) * d(y)) \\ &\leq d(y). \end{aligned}$$

Hence  $d(x) \leq d(y)$ , which implies that  $d$  is an isotone derivation of  $X$ .  $\square$

**DEFINITION 3.19.** An ideal  $I$  of  $X$  is said to be *d*-invariant if  $d(I) \subset I$ .

**PROPOSITION 3.20.** *Let  $d$  be a left derivation of a medial *BCH*-algebra  $X$ . Then  $d$  is regular if and only if every ideal of  $X$  is *d*-invariant.*

*Proof.* Let  $d$  is regular. Then by Theorem 3.10,  $d(x) = x$  for all  $x \in X$ . Let  $y \in d(A)$ , where  $A$  is an ideal of  $X$ . Then we have  $y = d(x)$  for some  $x \in A$ . Thus

$$\begin{aligned} y * x &= d(x) * x \\ &= x * x \\ &= 0 \in A. \end{aligned}$$

This implies that  $y \in A$  and  $d(A) \subset A$ . Conversely, let every ideal of  $X$  be  $d$ -invariant. Then  $d(\{0\}) \subset \{0\}$ , and so  $d(0) = 0$ , which implies that  $d$  is regular.  $\square$

**THEOREM 3.21.** *In a medial BCH-algebra  $X$ , a self-map  $d$  is a left derivation of  $X$  if and only if it is a  $(r, l)$ -derivation of  $X$ .*

*Proof.* Let  $d$  be a left derivation of a medial BCH-algebra  $X$ . First, we show that  $d$  is a  $(r, l)$ -derivation of  $X$ . Then

$$\begin{aligned} d(x * y) &= x * d(y) \\ &= (d(x) * y) * [(d(x) * y) * (x * d(y))] \\ &= (x * d(y)) \wedge (d(x) * y) \end{aligned}$$

for all  $x, y \in X$ . Conversely, let  $d$  be a  $(r, l)$ -derivation of  $X$ . Then

$$\begin{aligned} d(x * y) &= (x * d(y)) \wedge (d(x) * y) \\ &= (d(x) * y) * [(d(x) * y) * (x * d(y))] \\ &= x * d(y) = (y * d(x)) * [(y * d(x)) * (x * d(y))] \\ &= (x * d(y)) \wedge (y * d(x)). \end{aligned}$$

Hence,  $d$  is a left derivation of  $X$ .  $\square$

Let  $d$  be a left derivation of  $X$ . Define a set  $Fix_d(x)$  by

$$Fix_d(X) = \{x \in X \mid d(x) = x\}.$$

**PROPOSITION 3.22** *Let  $d$  be a left derivation of  $X$ . Then  $Fix_d(X)$  is a subalgebra of  $X$ .*

*Proof.* Let  $x, y \in Fix_d(X)$ . Then  $d(x) = x$  and  $d(y) = y$ , and so we have

$$\begin{aligned} d(x * y) &= (x * d(y)) \wedge (y * d(x)) = (x * y) \wedge (y * x) \\ &= (y * x) * ((y * x) * (x * y)) \\ &= x * y, \end{aligned}$$

which implies  $x * y \in Fix_d(X)$ .  $\square$

**PROPOSITION 3.23.** *Let  $d$  be a left derivation of a medial BCH-algebra  $X$ . If  $x, y \in Fix_d(X)$ , then  $x \wedge y \in Fix_d(X)$ .*

*Proof.* Let  $x, y \in \text{Fix}_d X$ , Then  $d(x) = x$  and  $d(y) = y$ , and so we have

$$\begin{aligned} d(x \wedge y) &= d(y * (y * x)) = (y * d(y * x)) \wedge ((y * x) * d(y)) \\ &= y * d(y * x) = (y * [(y * d(x)) \wedge (x * d(y))]) \\ &= y * (y * d(x)) = y * (y * x) \\ &= x \wedge y, \end{aligned}$$

which implies  $x \wedge y \in \text{Fix}_d(X)$ .  $\square$

**PROPOSITION 3.24.** *Let  $d$  be a left derivation of  $X$ . If  $x \in \text{Fix}_d(X)$ , then we have  $(d \circ d)(x) = x$ .*

*Proof.* Let  $x \in \text{Fix}_d(X)$ . Then we have

$$(d \circ d)(x) = d(d(x)) = d(x) = x.$$

This completes the proof.  $\square$

**PROPOSITION 3.25.** *Let  $d$  be a left derivation of  $X$ . If there exists  $x, y \in X$  such that  $x \leq y$  and  $y \in \text{Fix}_d(X)$ , then  $d$  is regular.*

*Proof.* Let  $x, y$  be such that  $x \leq y$  and  $d(y) = y$ . Then

$$\begin{aligned} d(0) &= d(x * y) \\ &= (x * d(y)) \wedge (y * d(x)) \\ &= (x * y) \wedge (y * d(x)) \\ &= 0 \wedge (y * d(x)) \\ &= (y * d(x)) * (y * d(x)) \\ &= 0. \end{aligned}$$

Hence  $d$  is regular.  $\square$

**PROPOSITION 3.26.** *Let  $d$  be a left derivation of a medial commutative *BCH*-algebra  $X$ . If  $x \leq y$  and  $y \in \text{Fix}_d(X)$ , then  $x \in \text{Fix}_d(X)$ .*

*Proof.* Let  $x, y$  be such that  $x \leq y$  and  $d(y) = y$ . Then

$$\begin{aligned}
d(x) &= d(x \wedge y) \\
&= d(y * (y * x)) \\
&= d(x * (x * y)) \\
&= (x * d(x * y)) \wedge ((x * y) * d(x)) \\
&= (x * ((x * d(y)) \wedge (y * d(x)))) \wedge ((x * y) * d(x)) \\
&= (x * ((x * y)) \wedge (y * d(x))) \wedge (0 * d(x)) \\
&= (x * (0 \wedge (y * d(x)))) \wedge (0 * d(x)) \\
&= x \wedge (0 * d(x)) \\
&= (0 * d(x)) * ((0 * d(x)) * x) \\
&= x.
\end{aligned}$$

Hence  $x \in \text{Fix}_d(X)$ . □

**THEOREM 3.27.** *Let  $X$  be a medial BCH-algebra and let  $d$  be a left derivation of  $X$ . If  $\text{Fix}_d(X) \neq \phi$ , then  $d$  is regular.*

*Proof.* Let  $y \in \text{Fix}_d(X)$ . Then we get  $d(y) = y$  and

$$\begin{aligned}
d(0) &= d(0 \wedge y) \\
&= d(y * (y * 0)) \\
&= (y * d(y * 0)) \wedge ((y * 0) * d(y)) \\
&= (y * d(y)) \wedge (y * d(y)) \\
&= (y * y) \wedge (y * y) \\
&= 0 * 0 = 0.
\end{aligned}$$

Hence  $d$  is regular. □

In the following, we will consider a left derivation  $d$  with  $\text{Fix}_d(X) \neq \phi$ .

**THEOREM 3.28.** *Let  $d$  be a left derivation of a medial BCH-algebra  $X$ . If  $X$  is commutative, then  $\text{Fix}_d(X)$  is an ideal of  $X$ .*

*Proof.* Let  $X$  be a medial BCH-algebra and let  $d$  be a left derivation of  $X$ . Then from Theorem 3.27,  $d$  is regular, and so  $0 \in \text{Fix}_d(X)$ . Let  $x * y \in \text{Fix}_d(X)$  and  $y \in \text{Fix}_d(X)$ . Then we get  $d(x * y) = x * y$  and

$d(y) = y$ . Thus we have

$$\begin{aligned}
d(x) &= d(x \wedge y) = d(y * (y * x)) \\
&= d(x * (x * y)) \\
&= (x * d(x * y)) \wedge ((x * y) * d(x)) \\
&= (x * (x * y)) \wedge ((x * y) * d(x)) \\
&= (y * (y * x)) \wedge ((x * y) * d(x)) \\
&= x \wedge ((x * y) * d(x)) \\
&= ((x * y) * d(x)) * [((x * y) * d(x)) * x] \\
&= x
\end{aligned}$$

□

**THEOREM 3.29.** *Let  $X$  be a medial *BCH*-algebra and let  $d_1$  and  $d_2$  be two isotone regular left derivations on  $X$ . Then  $d_1 = d_2$  if and only if  $Fix_{d_1}(X) = Fix_{d_2}(X)$ .*

*Proof.* It is clear that  $d_1 = d_2$  implies  $Fix_{d_1}(X) = Fix_{d_2}(X)$ . Conversely, let  $Fix_{d_1}(X) = Fix_{d_2}(X)$  and  $x \in X$ . By Proposition 3.13,  $d_1(x) \in Fix_{d_1}(X) = Fix_{d_2}(X)$  and so

$$d_2(d_1(x)) = d_1(x).$$

Similarly, we can have  $d_1(d_2(x)) = d_2(x)$ . Since  $d_1$  and  $d_2$  are isotone, we have  $d_2(d_1(x)) \leq d_2(x) = d_1(d_2(x))$  and so  $d_2(d_1(x)) \leq d_1(d_2(x))$ . Symmetrically, we can also have  $d_1(d_2(x)) \leq d_2(d_1(x))$ , which implies that

$$d_1(d_2(x)) = d_2(d_1(x)).$$

It follows that  $d_1(x) = d_2(d_1(x)) = d_1(d_2(x)) = d_2(x)$ , i.e.,  $d_1 = d_2$ . □

Let  $d$  be a left derivation of  $X$ . Define a *Kerd* by

$$Kerd = \{x \mid d(x) = 0\}$$

for all  $x \in X$ .

**THEOREM 3.30.** *Let  $d$  be a regular left derivation of a medial *BCH*-algebra  $X$ . Then *Kerd* is an ideal of  $X$ .*

*Proof.* Clearly,  $0 \in Kerd$ . Let  $x * y \in Kerd$  and  $y \in Kerd$ . Then we have  $0 = d(x * y) = x * d(y) = x * 0 = x$ , and so  $d(x) = d(0) = 0$ . This implies  $x \in Kerd$ . Hence *Kerd* is an ideal of  $X$ . □

DEFINITION 3.31. Let  $X$  be a  $BCH$ -algebra and let  $I$  be a proper ideal of  $X$ . Then  $I$  is said to be *prime* if  $a \wedge b \in I$  implies  $a \in I$ , or  $b \in I$ , for any  $a, b \in X$ .

THEOREM 3.32. *Let  $X$  be a medial  $BCH$ -algebra. Then for any left derivation  $d$ ,  $Ker d$  is a prime ideal of  $X$ .*

*Proof.* Let  $X$  be a medial  $BCH$ -algebra and let  $d$  be a left derivation of  $X$ . Then  $Ker d$  is an ideal of  $X$  by Theorem 3.30. Let  $x \wedge y \in Ker d$ . Then  $d(x \wedge y) = 0$ . Hence  $d(x) = d(x \wedge y) = 0$ , which implies  $x \in Ker d$ . This show that  $Ker d$  is a prime ideal of  $X$ .  $\square$

THEOREM 3.33. *Let  $d$  be a regular left derivation of a medial  $BCH$ -algebra  $X$ . Then the following are equivalent:*

- (1)  $Ker(d) = \{0\}$ ,
- (2)  $d$  is one-to-one,
- (3)  $d$  is the identity derivation.

*Proof.* ((1) $\implies$ (2)) Suppose that  $Ker(d) = \{0\}$  and  $d(x) = d(y)$ , for any  $x, y \in X$ . Since  $x \leq d(x)$ , it follows from (BCH11) that  $d(x * y) = x * d(y) \leq d(x) * d(y) = 0$ , so that  $d(x * y) = 0$ . Hence  $x * y \in Ker(d)$ , and so  $x * y = 0$ , and  $x \leq y$ . Similarly, we have  $y \leq x$ . This implies  $x = y$ . Therefore  $d$  is one-to-one.

((2) $\implies$ (3)) Suppose that  $d$  is one-to-one. For every  $x \in X$ , we have

$$d(d(x) * x) = d(x) * d(x) = 0 = d(0)$$

and so  $d(x) * x = 0$ , i.e.,  $d(x) \leq x$ . Since  $x \leq d(x)$  for every  $x \in X$ ,  $d(x) = x$ . Hence  $d$  is an identity derivation.

((3) $\implies$ (1)) It is obvious.  $\square$

DEFINITION 3.34. Let  $X$  be a  $BCH$ -algebra and let  $d_1, d_2$  be two self-maps of  $X$ . Define  $d_1 \circ d_2 : X \rightarrow X$  by

$$(d_1 \circ d_2)(x) = d_1(d_2(x))$$

for all  $x \in X$ .

Let  $Der_r(X)$  be the set of all isotone regular derivations of a  $BCH$ -algebra  $X$ . Then  $Der_r(X)$  is a poset with the partial order defined by

$$d_1 \leq d_2 \text{ if and only if } d_1(x) \leq d_2(x) \text{ for all } x \in X,$$

for any  $d_1, d_2 \in Der_r(X)$  and identity map  $id_X$  is the least element in  $Der_r(X)$ .

**THEOREM 3.35.** *Let  $X$  be a medial *BCH*-algebra. Then  $Der_r(X)$  is a semilattice with  $d_1 \vee d_2 = d_1 \circ d_2$  for every  $d_1, d_2 \in Der_r(X)$ .*

*Proof.* Let  $d_1, d_2 \in Der_r(X)$ . Then  $x \leq d_2(x)$  implies  $d_1(x) \leq d_1(d_2(x))$ , for all  $x \in X$ . Also,  $d_2(x) \leq d_1(d_2(x))$ , for all  $x \in X$ . That is,  $d_1 \leq d_1 \circ d_2$  and  $d_2 \leq d_1 \circ d_2$ , hence  $d_1 \circ d_2$  is an upper bound of  $d_1$  and  $d_2$ .

Suppose that  $d$  is an upper bound of  $d_1$  and  $d_2$ . Then  $d_1(x) \leq d(x)$  and  $d_2(x) \leq d(x)$ , for all  $x \in X$ . These imply  $d_1(d_2(x)) \leq d_1(d(x)) \leq d(d(x)) = d(x)$ , for all  $x \in X$ , and  $d_1 \circ d_2 \leq d$ . Hence  $d_1 \circ d_2$  is the least upper bound of  $d_1$  and  $d_2$ .  $\square$

**PROPOSITION 3.36.** *Let  $X$  be a medial *BCH*-algebra and let  $d_1, d_2$  be two left derivations of  $X$ . Then  $d_1 \circ d_2$  is a left derivation of  $X$ .*

*Proof.* Let  $x, y \in X$ . Then we have

$$\begin{aligned} (d_1 \circ d_2)(x * y) &= d_1(d_2(x * y)) \\ &= d_1((x * d_2(y)) \wedge (y * d_2(x))) \\ &= d_1((y * d_2(x)) * ((y * d_2(x)) * (x * d_2(y)))) \\ &= d_1(x * d_2(y)) = x * d_1(d_2(y)) \\ &= (y * d_1(d_2(x))) * [(y * (d_1(d_2(x))) * (x * d_1(d_2(y))))] \\ &= (x * (d_1 \circ d_2)(y)) \wedge (y * (d_1 \circ d_2)(x)). \end{aligned}$$

Hence  $d_1 \circ d_2$  is a left derivation of  $X$ .  $\square$

**DEFINITION 3.37.** Let  $X$  be a *BCH*-algebra and let  $d_1, d_2$  be two self-maps of  $X$ . Define  $d_1 \wedge d_2 : X \rightarrow X$  by

$$(d_1 \wedge d_2)(x) = d_1(x) \wedge d_2(x)$$

for all  $x \in X$ .

**PROPOSITION 3.38.** *Let  $X$  be a medial *BCH*-algebra and let  $d_1$  and  $d_2$  be two left derivations of  $X$ . Then  $d_1 \wedge d_2$  is a left derivation of  $X$ .*

*Proof.* Let  $x, y \in X$ . Then we have

$$\begin{aligned} (d_1 \wedge d_2)(x * y) &= d_1(x * y) \wedge d_2(x * y) \\ &= (x * d_1(y)) \wedge (y * d_2(x)) \\ &= x * d_1(y) \\ &= (x * (d_1 \wedge d_2)(y)) \wedge (y * (d_1 \wedge d_2)(x)). \end{aligned}$$

Hence  $d_1 \wedge d_2$  is a left derivation of  $X$ .  $\square$

$Der(X)$  denote the set of all left derivations of  $X$ .

PROPOSITION 3.39. *Let  $d_1, d_2, d_3 \in Der(X)$ . Then*

$$d_1 \wedge (d_2 \wedge d_3) = (d_1 \wedge d_2) \wedge d_3.$$

*Proof.* Let  $d_1, d_2$  and  $d_3$  be left derivations on  $X$ . Then

$$\begin{aligned} ((d_1 \wedge d_2) \wedge d_3)(x * y) &= (d_1 \wedge d_2)(x * y) \wedge d_3(x * y) \\ &= d_3(x * y) * (d_3(x * y) * (d_1 \wedge d_2)(x * y)) \\ &= (d_1 \wedge d_2)(x * y) \\ &= (x * d_2(y)) * ((x * d_2(y)) * (x * d_1(y))) \\ &= x * d_1(y). \end{aligned}$$

Similarly, we have

$$\begin{aligned} d_1 \wedge (d_2 \wedge d_3)(x * y) &= d_1(x \wedge y) \wedge (d_2 \wedge d_3)(x * y) \\ &= d_1(x * y) \wedge ((d_2(x * y) \wedge d_3(x * y))) \\ &= (x * d_1(y)) \wedge ((x * d_3(y)) * ((x * d_3(y)) * x * d_2(y))) \\ &= x * d_1(y). \end{aligned}$$

This implies that  $(d_1 \wedge (d_2 \wedge d_3))(x * y) = ((d_1 \wedge d_2) \wedge d_3)(x * y)$ . Put  $y = 0$ , we have  $(d_1 \wedge (d_2 \wedge d_3))(x) = ((d_1 \wedge d_2) \wedge d_3)(x)$ . This implies  $d_1 \wedge (d_2 \wedge d_3) = (d_1 \wedge d_2) \wedge d_3$ .  $\square$

THEOREM 3.40. *Let  $X$  be a BCH-algebra. Then  $(Der(X), \wedge)$  is a semigroup.*

## References

- [1] M. A. Chaudhry, *On BCH-algebras*, Math. Japonica. **36** (4) (1991), 665–676.
- [2] M. A. Chaudhry and H. Fakhur-ud-din, *Ideals and Filter in BCH-algebras*, Math. Japonica. **44** (1996) no. 1, 101–112.
- [3] Q. P. Hu and X. Li, *On BCH-algebra*, Math. Sem. Notes Kobe Univ. **11** (2) (1983), 313–320.
- [4] Q. P. Hu and X. Li, *On proper BCH-algebra*, Math. Japon. **30** (4) (1985), 659–661.
- [5] K. Iseki, *An algebra related with a propositional calculus*, Proc. Japan Acad **42** (1966), 26–29.
- [6] Y. Imai and K. Iseki, *On axiom system of propositional calculi XIV*, Proc. Japan Academy. **42** (1966), 19–22.

**Kyung Ho Kim**

Department of Mathematics  
Korea National University of Transportation  
Chungju 380-702, Korea  
*E-mail:* ghkim@ut.ac.kr

**Yong Hoon Lee**

Department of Mathematics  
Dankook University  
Cheonan 330-714, Korea  
*E-mail:* yonghoon@dankook.ac.kr