

ON KANTOROVICH FORM OF GENERALIZED SZÁSZ-TYPE OPERATORS USING CHARLIER POLYNOMIALS

ABDUL WAFI[†], NADEEM RAO, AND DEEPMALA^{*}

ABSTRACT. The aim of this article is to introduce a new form of Kantorovich Szász-type operators involving Charlier polynomials. In this manuscript, we discuss the rate of convergence, better error estimates. Further, we investigate order of approximation in the sense of local approximation results with the help of Ditzian-Totik modulus of smoothness, second order modulus of continuity, Peetre's K-functional and Lipschitz class.

1. Introduction

Bernstein [2] defined the positive linear operators using binomial distribution and proved pointwise and uniform approximation in the space of continuous functions on $[0, 1]$. These operators provide the powerful tool for numerical analysis, computer added geometric design (CAGD) and solutions of differential equations. But these operators are not suitable for discontinuous functions. Later on, Kantorovich [9] generalized the Bernstein operators for integrable functions. Szász [21] introduced

Received January 26, 2017. Revised March 19, 2017. Accepted March 20, 2017.
2010 Mathematics Subject Classification: 41A10, 41A25, 41A36, 41A36.

Key words and phrases: Kantorovich Szász operators, Charlier polynomials, Ditzian-Totik modulus of smoothness, Peetre's K-functional, Lipschitz class.

[†] This work was supported by University Grant Commission (UGC) under the scheme of Basic Scientific Research (BSR) Fellowship.

^{*} Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2017.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

linear positive operators in the sense of exponential growth on non-negative semi axes. Several generalizations of these operators have been studied by different researchers ([1, 6, 7, 13, 18–20, 23–26]). Some interesting results can be seen in ([5, 12, 14–17]). Many operators preserve the constant and linear functions but these operators do not preserve x^2 . King [10] introduced a method in order to preserve x^2 for the Bernstein operators. Recently, Varma and Taşdelen [27] gave a Kantorovich-Szász type operators by means of Charlier polynomials [8] having the generating function of the form

$$(1) \quad e^t \left(1 - \frac{t}{a}\right)^u = \sum_{k=0}^{\infty} C_k^{(a)}(u) \frac{t^k}{k!}, \quad |t| < a,$$

and the explicit representation

$$C_k^{(a)}(u) = \sum_{r=0}^k \binom{k}{r} (-u)_r \left(\frac{1}{a}\right)^r,$$

where $(\alpha)_k$ is the Pochhammer's symbol given by

$$(\alpha)_0 = 1, (\alpha)_k = \alpha(\alpha + 1)\dots(\alpha + k - 1), \quad k \in \mathbb{N}.$$

We note that for $a > 0$ and $u \leq 0$, Charlier polynomials are positive. They [27] defined Kantorovich-Szász-type operators as

$$(2) \quad L_n^*(f; x, a) = ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nx)}{k!} \\ \times \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) dt,$$

where $a > 1$, $n \in \mathbb{N}$ and $x \geq 0$. Motivated by the above development, we define a new sequence of Kantorovich-Szász-type operators which preserves constant and quadratic test functions i.e. $e_0(x)$ and $e_2(x)$ ($e_i(x) = x^i, i = 0, 2$)

$$(3) \quad K_{n,a}^*(f; r_{n,a}^*(x)) = ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \\ \times \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(s) ds,$$

where

$$(4) \quad r_{n,a}^*(x) = \frac{-(4 + \frac{1}{a-1}) + \sqrt{(4 + \frac{1}{a-1})^2 + 4(n^2x^2 - \frac{10}{3})}}{2n}$$

and $r_{n,a}^*(x) \geq 0$ for $x \in \left[\frac{\sqrt{\frac{10}{3}}}{n}, \infty \right)$.

We observe that

(i) For a fixed $x \in \left[\frac{\sqrt{\frac{10}{3}}}{n}, \infty \right)$, $r_{n,a}^*(x) \rightarrow x$ as $n \rightarrow \infty$ on $[0, \infty)$ and

operators (3) reduce to operators (2)

and

(ii) For a fixed x , $r_{n,a}^*(x) = x$ as $a \rightarrow \infty$ and taking $x - \frac{1}{n}$ instead of x , operators (3) reduce to the Classical Kantorovich-Szász operators [22].

In the present paper, we discuss the rate of convergence for continuous functions, first order derivative of the function. Further, we investigate some direct and local approximation results using Ditzian-Totik modulus of smoothness, second order modulus of continuity, Peetre's K-functional and Lipschitz space.

2. Basic Estimates

LEMMA 2.1. *From generating function (1) and differentiation, we have*

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} &= e \left(1 - \frac{1}{a} \right)^{-(a-1)nr_{n,a}^*(x)}, \\ \sum_{k=0}^{\infty} k \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} &= e \left(1 - \frac{1}{a} \right)^{-(a-1)nr_{n,a}^*(x)} (1 + nr_{n,a}^*(x)), \\ \sum_{k=0}^{\infty} k^2 \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} &= e \left(1 - \frac{1}{a} \right)^{-(a-1)nr_{n,a}^*(x)} \\ &\quad \times \left(2 + \left(3 + \frac{1}{a-1} \right) nr_{n,a}^*(x) + n^2 r_{n,a}^{*2}(x) \right). \end{aligned}$$

LEMMA 2.2. Let $e_i(t) = t^i$, $i = 0, 1, 2$ and $r_{n,a}^*(x) \geq 0$. Then for the operators $K_{n,a}^*$ defined by (3), we have

$$\begin{aligned} K_{n,a}^*(e_0; r_{n,a}^*(x)) &= 1, \\ K_{n,a}^*(e_1; r_{n,a}^*(x)) &= \frac{-(1 + \frac{1}{a-1}) + \sqrt{(4 + \frac{1}{a-1})^2 + 4(n^2x^2 - \frac{10}{3})}}{2n}, \\ K_{n,a}^*(e_2; r_{n,a}^*(x)) &= x^2. \end{aligned}$$

Proof. Using Lemma 2.1 and equation (4), we prove Lemma 2.2. \square

LEMMA 2.3. Let $\psi_x^i(t) = (t - x)^i$, $i = 0, 1, 2$ and $r_{n,a}^*(x) \geq 0$, $n \in \mathbb{N}$. Then we have

$$\begin{aligned} K_{n,a}^*(\psi_x^0; r_{n,a}^*(x)) &= 1, \\ K_{n,a}^*(\psi_x^1; r_{n,a}^*(x)) &= -\frac{(1 + \frac{1}{a-1})}{2n} + \frac{\frac{8}{3} + \frac{8}{a-1} + \frac{1}{(a-1)^2}}{2n(\sqrt{(4 + \frac{1}{a-1})^2 + 4(n^2x^2 - \frac{10}{3})} + 2nx)}, \\ K_{n,a}^*(\psi_x^2; r_{n,a}^*(x)) &= \left(1 + \frac{1}{a-1}\right) \cdot \frac{x}{n} - \frac{\frac{8}{3} + \frac{8}{a-1} + \frac{1}{(a-1)^2}}{\sqrt{(4 + \frac{1}{a-1})^2 + 4(n^2x^2 - \frac{10}{3})} + 2nx} \\ &\quad \times \frac{x}{n} + x^2. \end{aligned}$$

Proof. In view of Lemma 2.2 and linearity property, we can easily prove this Lemma. \square

3. Rate of convergence

For $f \in C[0, \infty)$, where $C[0, \infty)$ is the set of all continuous functions on $[0, \infty)$, the modulus of continuity for a uniformly continuous f is

$$\omega(f; \delta) = \sup_{|t-y| \leq \delta} |f(t) - f(y)|, \quad t, y \in [0, \infty).$$

For a uniformly continuous f in $C[0, \infty)$ and $\delta > 0$, one has

$$(5) \quad |f(t) - f(y)| \leq \left(1 + \frac{(t-y)^2}{\delta^2}\right) \omega(f; \delta).$$

THEOREM 3.1. *Let f be uniformly continuous and bounded in $C\left[\frac{\sqrt{\frac{10}{3}}}{n}, \infty\right)$, $n \in \mathbb{N}$. Then for the operators $K_{n,a}^*$, we have*

$$|K_{n,a}^*(f; x) - f(x)| \leq 2\omega\left(f; \delta_{n,a}\right),$$

where $\delta_{n,a} = \sqrt{K_{n,a}^*(\psi_x^2; x)}$, holds uniformly in each compact subset of $\left[\frac{\sqrt{\frac{10}{3}}}{n}, \infty\right)$.

Proof. From (5), we have

$$\begin{aligned} & |K_{n,a}^*(f; x) - f(x)| \\ & \leq ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \\ & \quad \times \int_{\frac{k}{n}}^{\frac{k+1}{n}} |f(t) - f(x)| dt \\ & \leq \left\{ ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \right. \\ & \quad \times \left. \int_{\frac{k}{n}}^{\frac{k+1}{n}} \left(1 + \frac{(t-x)^2}{\delta_{n,a}^2}\right) dt \right\} \omega(f; \delta_{n,a}) \\ & \leq \left\{ 1 + \frac{K_{n,a}^*(\psi_x^2; x)}{\delta_{n,a}^2} \right\} \omega(f; \delta_{n,a}) \\ & = 2\omega(f; \delta_{n,a}), \end{aligned}$$

where $\delta_{n,a} = \sqrt{K_{n,a}^*(\psi_x^2; x)}$, holds uniformly in each compact subset of $[0, \infty)$. □

REMARK 3.2. For the Kantorovich-Szász type operators L_n^* given by (2), we have, for every $f \in C[0, \infty) \cap E$, where

$$E := \{f : [0, \infty) \rightarrow \mathbb{R} : |f(x)| \leq Me^{Ax}, A \in \mathbb{R}, M \in \mathbb{R}^+\},$$

one has

$$|L_n^*(f; x, a) - f(x)| \leq 2\omega(f; \delta),$$

where $\delta = \sqrt{\frac{x}{n} \left(1 + \frac{1}{a-1}\right) + \frac{10}{3n^2}}$ (see [27]). Here, we shall show that our operators $K_{n,a}^*$ has the better approximation than the operators L_n^* . Since

$$\frac{x}{n} \left(1 + \frac{1}{a-1}\right) < \frac{x}{n} \left(1 + \frac{1}{a-1}\right) + \frac{10}{3n^2},$$

which shows that $\delta_{n,a} < \delta$. This implies that our operators (3) converge uniformly more rapidly than operators defined by (2).

THEOREM 3.3. *Let $\omega_1(f; \delta)$ is the modulus of continuity of $f'(x)$. For $f \in C \left[\frac{\sqrt{\frac{10}{3}}}{n}, \infty \right)$ and bounded on $\left[\frac{\sqrt{\frac{10}{3}}}{n}, \infty \right)$, $n \in \mathbb{N}$ with continuous derivative, we have*

$$|K_{n,a}^*(f; x) - f(x)| \leq \omega_1(f'; n^{-\frac{1}{2}}) \sqrt{K_{n,a}^*(\psi_x^2(t); x)} \left\{ 1 + \sqrt{n} \sqrt{K_{n,a}^*(\psi_x^2(t); x)} \right\},$$

where $\delta_n = n^{-\frac{1}{2}}$.

Proof. It is known that

$$\begin{aligned} f(x_1) - f(x_2) &= (x_1 - x_2)f'(\xi) \\ (6) \qquad \qquad &= (x_1 - x_2)f'(x_1) + (x_1 - x_2)[f'(\xi) - f'(x_1)], \end{aligned}$$

for $x_1, x_2 \in \left[\frac{\sqrt{\frac{10}{3}}}{n}, b \right]$, $b > \frac{\sqrt{\frac{10}{3}}}{n}$ and $x_1 < \xi < x_2$. Also, we have (see [11], Theorem 1.6.2, pp. 21)

$$|(x_1 - x_2)[f'(\xi) - f'(x_1)]| \leq |x_1 - x_2|(\lambda + 1)\omega_1(\delta), \quad \lambda = \lambda(x_1, x_2; \delta). \quad (7)$$

Next, we find

$$\begin{aligned}
 |K_{n,a}^*(f; x) - f(x)| &= \left| ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \right. \\
 (8) \quad &\times \left. \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) - f(x) dt \right|.
 \end{aligned}$$

Using (6) and (7), we get

$$\begin{aligned}
 |K_{n,a}^*(f; x) - f(x)| &\leq \left| ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \right. \\
 &\times \left. \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \int_{\frac{k}{n}}^{\frac{k+1}{n}} (x-t) f'(x) dt \right| \\
 &\quad + \omega_1(f; \delta_n) ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \\
 &\times \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \int_{\frac{k}{n}}^{\frac{k+1}{n}} |t-x|(\lambda+1) dt \\
 &\leq \omega_1(f; \delta_n) \left\{ e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \right. \\
 &\times \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \int_{\frac{k}{n}}^{\frac{k+1}{n}} |t-x| dt \\
 &\quad \left. + e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \right.
 \end{aligned}$$

$$\begin{aligned}
& \times \left\{ \sum_{\lambda \geq 1} C_k^{(a)}(- (a-1)nr_{n,a}^*(x)) \int_{\frac{k}{n}}^{\frac{k+1}{n}} |t-x| \lambda(x,t; \delta) dt \right\} \\
& \leq \omega_1(f'; \delta_n) \left\{ e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \int_{\frac{k}{n}}^{\frac{k+1}{n}} |t-x| dt \right. \\
& \quad \left. + \frac{1}{\delta_n} e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(- (a-1)nr_{n,a}^*(x))}{k!} \int_{\frac{k}{n}}^{\frac{k+1}{n}} (t-x)^2 dt \right\} \\
& \leq \omega_1(f'; \delta_n) \left(\sqrt{K_{n,a}^*(\psi_x^2; x)} + \frac{K_{n,a}^*(\psi_x^2; x)}{\delta_n} \right) \\
& = \omega_1(f'; \delta_n) \sqrt{K_{n,a}^*(\psi_x^2; x)} \left\{ 1 + \frac{\sqrt{K_{n,a}^*(\psi_x^2; x)}}{\delta_n} \right\}.
\end{aligned}$$

Taking $\delta_n = n^{-\frac{1}{2}}$, we get

$$|K_{n,a}^*(f; x) - f(x)| \leq \omega_1(f'; n^{-\frac{1}{2}}) \sqrt{K_{n,a}^*(\psi_x^2(t); x)} \left\{ 1 + \sqrt{n} \sqrt{K_{n,a}^*(\psi_x^2(t); x)} \right\}.$$

□

4. Direct Estimate

Ditzian-Totik Modulus of smoothness [4] is defined for continuous and bounded functions as:

$$\begin{aligned}
& \omega_{\varphi^\lambda}^2(f; \delta) \\
& = \sup_{0 < h \leq \delta} \|\Delta_{h\varphi(x)}^2 f(x)\| \\
& = \sup_{0 < h \leq \delta} \sup_{x \pm h\varphi^\lambda(x) \in [0, b], b < \infty} |f(x - h\varphi^\lambda(x)) - 2f(x) + f(x + h\varphi^\lambda(x))|,
\end{aligned}$$

where $\varphi^2(x) = x$ and Peetre's K-functional [4] is given by

$$K_{\varphi^\lambda}(f, \delta^2) = \inf_g \left(\|f - g\|_{C[0,b], b < \infty} + \delta^2 \|\varphi^{2\lambda} g''\|_{C[0,b], b < \infty} \right), \quad g, g' \in AC_{loc}. \quad (9)$$

The K-functional is equivalent to the modulus of smoothness, i.e.,

$$(10) \quad C^{-1}K_{\varphi^\lambda}(f, \delta^2) \leq \omega_{\varphi^\lambda}^2(f, \delta) \leq CK_{\varphi^\lambda}(f, \delta^2) \text{ where } C > 0.$$

THEOREM 4.1. For all the continuous functions f defined on $\left[\frac{\sqrt{\frac{10}{3}}}{n}, b \right], \frac{\sqrt{\frac{10}{3}}}{n}$
 $< b < \infty, n \in \mathbb{N}$ and $a > 1$, we have

$$|K_{n,a}^*(f; x) - f(x)| \leq C\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}) \text{ for the large value of } n,$$

where $0 \leq \lambda \leq 1$ and $\varphi^2(x) = x$.

Proof. Using (9), (10), we have

$$(11) \quad \|f - g\|_{C[0,b], b < \infty} \leq A\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}),$$

$$(12) \quad n^{-1}\varphi(x)^{2-2\lambda} \|\varphi^{2\lambda} g''\|_{C[0,b], b < \infty} \leq B\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}).$$

Next, we can choose $g_n \equiv g_{n,x,\lambda}$ for fixed x and λ such that

$$\begin{aligned} & |K_{n,a}^*(f; x) - f(x)| \\ & \leq |K_{n,a}^*(f - g_n; x) - (f - g_n)(x)| + |K_{n,a}^*(g_n; x) - g_n(x)|, \\ & \leq 2 \|f - g_n\|_{C[0,b], b < \infty} + |K_{n,a}^*(g_n; x) - g_n(x)|. \end{aligned}$$

From (11), we get

$$|K_{n,a}^*(f; x) - f(x)| \leq 2A\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}) + |K_{n,a}^*(g_n; x) - g_n(x)|. \quad (13)$$

Now, the last term can be calculated by using Taylor's formula

$$\begin{aligned}
& |K_{n,a}^*(g_n(t) - g_n(x); x)| \\
& \leq |g'_n(x)K_{n,a}^*((t-x); x)| + \left| K_{n,a}^* \left(\int_t^x (x-u)g''_n(u)du; x \right) \right| \\
& \leq K_{n,a}^* \left(\frac{|x - \frac{k}{n}|}{\varphi^{2\lambda}(x)} \int_{\frac{k}{n}}^x \varphi^{2\lambda}(u)|g''_n(u)du|; x \right) \\
& \leq \| \varphi^{2\lambda} g''_n \|_{C[0,b], b < \infty} \frac{1}{\varphi^{2\lambda}(x)} K_{n,a}^*((t-x)^2; x) \\
& \leq \| \varphi^{2\lambda} g''_n \|_{C[0,b], b < \infty} \frac{1}{\varphi^{2\lambda}(x)} \frac{x n K_{n,a}^*((t-x)^2; x)}{n x} \\
& \leq \| \varphi^{2\lambda} g''_n \|_{C[0,b], b < \infty} \frac{x n^{-1} n K_{n,a}^*((t-x)^2; x)}{\varphi^{2\lambda}(x) x}.
\end{aligned}$$

For sufficiently large value of n , we get

$$\frac{n K_{n,a}^*((t-x)^2; x)}{x} \leq \left(1 + \frac{1}{a-1} \right).$$

Therefore

$$(14) \quad |K_{n,a}^*(g_n(t) - g_n(x); x)| \leq \left(1 + \frac{1}{a-1} \right) B \omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}} \varphi(x)^{1-\lambda}).$$

Using (13) and (14), we get

$$|K_{n,a}^*(f(t) - f(x); x)| \leq M \omega_\lambda^2 \left(f, n^{-\frac{1}{2}} \varphi(x)^{1-\lambda} \right),$$

where $M = \max \left(2A, \left(1 + \frac{1}{a-1} \right) B \right)$. □

Let $C_B[0, \infty)$ denote the space of real valued continuous and bounded functions f on $[0, \infty)$ endowed with the norm

$$\|f\| = \sup_{0 \leq x < \infty} |f(x)|.$$

Then, for any $\delta > 0$, Peeter's K-functional is defined as

$$K_2(f, \delta) = \inf\{\|f - g\| + \delta\|g''\| : g \in C_B^2[0, \infty)\},$$

where $C_B^2[0, \infty) = \{g \in C_B[0, \infty) : g', g'' \in C_B[0, \infty)\}$. By Devore and Lorentz [3], p.177, Theorem 2.4], there exists an absolute constant $C > 0$ such that

$$K_2(f; \delta) \leq C\omega_2(f; \sqrt{\delta}),$$

where $\omega_2(f; \delta)$ is the second order modulus of continuity is defined as

$$\omega_2(f, \sqrt{\delta}) = \sup_{0 < h \leq \sqrt{\delta}} \sup_{x \in [0, \infty)} |f(x + 2h) - 2f(x + h) + f(x)|.$$

THEOREM 4.2. *Let $f \in C_B^2\left[\frac{\sqrt{10}}{n}, \infty\right)$, $n \in \mathbb{N}$. Then for all $x \in \left[\frac{\sqrt{10}}{n}, \infty\right)$, $n \in \mathbb{N}$ there exist a constant $C > 0$ such that*

$$|K_{n,a}^*(f; x) - f(x)| \leq C\omega_2(f; \sqrt{\Pi_{n,a}(x)}) + \omega(f; K_{n,a}^*(\psi_x; x))$$

where $\Pi_{n,a}(x) = K_{n,a}^*(\psi_x^2; x) + (K_{n,a}^*(\psi_x; x))^2$.

Proof. First, we define the auxiliary operators as follows

$$(15) \quad \widehat{K}_{n,a}^*(f; x) = K_{n,a}^*(f; x) + f(x) - f(\Lambda_{n,a}(x)),$$

where $\Lambda_{n,a}(x) = K_{n,a}^*(\psi_x; x) + x$. We find that

$$\widehat{K}_{n,a}^*(1; x) = 1,$$

$$\widehat{K}_{n,a}^*(\psi_x(t); x) = 0,$$

$$(16) \quad |\widehat{K}_{n,a}^*(f; x)| \leq 3\|f\|.$$

Let $g \in C_B^2 \left[\frac{\sqrt{10}}{n}, \infty \right)$, using Taylor's theorem, we have

$$g(t) = g(x) + (t-x)g'(x) + \int_x^t (t-v)g''(v)dv.$$

Now

$$\begin{aligned} \widehat{K}_{n,a}^*(g; x) - g(x) &= g'(x)\widehat{K}_{n,a}^*(t-x; x) + \widehat{K}_{n,a}^*\left(\int_x^t (t-v)g''(v)dv; x\right) \\ &= \widehat{K}_{n,a}^*\left(\int_x^t (t-v)g''(v)dv; x\right) \\ &= K_{n,a}^*\left(\int_x^t (t-v)g''(v)dv; x\right) - \int_x^{\Lambda_{n,a}} (\Lambda_{n,a}-v)g''(v)dv. \end{aligned}$$

Therefore

$$(17) \quad \begin{aligned} &|\widehat{K}_{n,a}^*(g; x) - g(x)| \\ &\leq \left| K_{n,a}^*\left(\int_x^t (t-v)g''(v)dv; x\right) \right| + \left| \int_x^{\Lambda_{n,a}} (\Lambda_{n,a}-v)g''(v)dv \right|. \end{aligned}$$

Since

$$(18) \quad \left| \int_x^t (t-v)g''(v)dv \right| \leq (t-x)^2 \|g''\|,$$

and

$$(19) \quad \left| \int_x^{\Lambda_{n,a}} (\Lambda_{n,a} - v)g''(v)dv \right| \leq (\Lambda_{n,a} - x)^2 \|g''\|.$$

Then from (17), (18) and (19), we have

$$(20) \quad \begin{aligned} |\widehat{K}_{n,a}^*(g; x) - g(x)| &\leq \left\{ K_{n,a}^*((t-x)^2; x) + (\Lambda_{n,a} - x)^2 \right\} \|g''\| \\ &= \Pi_{n,a}(x) \|g''\|. \end{aligned}$$

Next, we have

$$\begin{aligned} |K_{n,a}^*(f; x) - f(x)| &\leq |\widehat{K}_{n,a}^*(f - g; x)| + |(f - g)(x)| + |\widehat{K}_{n,a}^*(g; x) - g(x)| \\ &\quad + |f(\Lambda_{n,a}) - f(x)|, \end{aligned}$$

using (20), we have

$$\begin{aligned} |K_{n,a}^*(f; x) - f(x)| &\leq 4\|f - g\| + |\widehat{K}_{n,a}^*(g; x) - g(x)| + |f(\Lambda_{n,a}) - f(x)| \\ &\leq 4\|f - g\| + \Pi_{n,a}(x) \|g''\| + \omega\left(f; K_{n,a}^*(\psi_x; x)\right). \end{aligned}$$

By the definition of Peetre's K-functional, we find

$$|K_{n,a}^*(f; x) - f(x)| \leq C\omega_2\left(f; \sqrt{\Pi_{n,a}(x)}\right) + \omega\left(f; K_{n,a}^*(\psi_x; x)\right).$$

This completes the proof of Theorem 4.2. □

Now, we discuss a local result in Lipschitz class

$$Lip_M^*(\alpha) = \left\{ f \in C_B[0, \infty) : |f(t) - f(x)| \leq M \frac{|t-x|^\alpha}{(t+x)^{\frac{\alpha}{2}}} : x, t \in (0, \infty) \right\},$$

where M is a constant and $0 < \alpha \leq 1$.

THEOREM 4.3. *Let $f \in Lip_M^*(\alpha)$ and $x \in \left(\frac{\sqrt{10}}{n}, \infty\right)$, $n \in \mathbb{N}$. Then, we have*

$$|K_{n,a}^*(f; x) - f(x)| \leq M \left[\frac{\Theta_{n,a}(x)}{x} \right]^{\frac{\alpha}{2}},$$

where $\Theta_{n,a}(x) = K_{n,a}^*((t-x)^2; x)$.

Proof. Let $\alpha = 1$ and $x \in (0, \infty)$. Then, for $f \in Lip_M^*(1)$, we have

$$\begin{aligned} & |K_{n,a}^*(f; x) - f(x)| \\ & \leq ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \\ & \quad \times \int_{\frac{k}{n}}^{\frac{k+1}{n}} |f(t) - f(x)| dt \\ & \leq Mne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \\ & \quad \times \int_{\frac{k}{n}}^{\frac{k+1}{n}} \frac{|t-x|}{\sqrt{t+x}} dt. \\ & \leq \frac{M}{\sqrt{x}} ne^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \\ & \quad \times \int_{\frac{k}{n}}^{\frac{k+1}{n}} |t-x| dt \\ & \leq \frac{M}{\sqrt{x}} K_{n,a}^*(|t-x|; x) \\ & \leq M \frac{\sqrt{K_{n,a}^*((t-x)^2; x)}}{\sqrt{x}} \\ & = M \left(\frac{\Theta_{n,a}(x)}{x} \right)^{\frac{1}{2}}. \end{aligned}$$

Thus, the assertion holds for $\alpha = 1$. Now, we will prove for $\alpha \in (0, 1)$. From the Hölder Inequality with $p = \frac{1}{\alpha}$, $q = \frac{1}{1-\alpha}$, we have

$$\begin{aligned}
 & |K_{n,a}^*(f; x) - f(x)| \\
 &= \left(e^{-1} \left(1 - \frac{1}{a} \right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \right. \\
 &\times \left. \left(n \int_{\frac{k}{n}}^{\frac{k+1}{n}} |f(t) - f(x)| dt \right)^{\frac{1}{\alpha}} \right)^{\alpha} \\
 &\times \left(e^{-1} \left(1 - \frac{1}{a} \right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \right)^{1-\alpha} \\
 &\leq \left(e^{-1} \left(1 - \frac{1}{a} \right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \right. \\
 &\times \left. \left(n \int_{\frac{k}{n}}^{\frac{k+1}{n}} |f(t) - f(x)| dt \right)^{\frac{1}{\alpha}} \right)^{\alpha}.
 \end{aligned}$$

Since $f \in Lip_M^*$, we obtain

$$\begin{aligned}
 & |K_{n,a}^*(f; x) - f(x)| \\
 &\leq M \left(ne^{-1} \left(1 - \frac{1}{a} \right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \right. \\
 &\times \left. \int_{\frac{k}{n}}^{\frac{k+1}{n}} \frac{|t-x|}{\sqrt{t+x}} dt \right)^{\alpha} \\
 &\leq \frac{M}{x^{\frac{\alpha}{2}}} \left(ne^{-1} \left(1 - \frac{1}{a} \right)^{(a-1)nr_{n,a}^*(x)} \sum_{k=0}^{\infty} \frac{C_k^{(a)}(-(a-1)nr_{n,a}^*(x))}{k!} \right. \\
 &\times \left. \int_{\frac{k}{n}}^{\frac{k+1}{n}} |t-x| dt \right)^{\alpha} \\
 &= \frac{M}{x^{\frac{\alpha}{2}}} (K_{n,a}^*(|t-x|; x))^{\alpha} \\
 &\leq M \left(\frac{\Theta_{n,a}(x)}{x} \right)^{\frac{\alpha}{2}}.
 \end{aligned}$$

This completes the proof of Theorem 4.3. \square

Acknowledgement

The authors are thankful to the referee(s) and editor for their valuable suggestions which improves the paper in the present form. The second author is thankful to University Grant Commission(UGC), New Delhi-India, for financial support under UGC-BSR scheme UGC-BSR-Fellowship-Mathematics/RPS/RO/JMI/2015.

References

- [1] T.Acar, L.N.Mishra and V.N.Mishra, *Simultaneous approximation for generalized Srivastava-Gupta operator*, Journal of Function Spaces, **2015**, Article ID 936308, (2015) 11 pages, doi:10.1155/2015/936308.
- [2] S.N. Bernstein, *Démonstration du théorème de Weierstrass*, fondée sur le calcul des probabilités, Commun. Soc. Math. Kharkow **2** (13) (1912-1913), 1–2.
- [3] R.A. DeVore and G.G. Lorentz, *Constructive Approximation*, Grundlehren der Mathematischen Wissenschaften [Fundamental principles of Mathematical Sciences], (Springer-Verlag, Berlin, 1993).
- [4] Z. Ditzian and V. Totik, *Moduli of smoothness*, Springer Series in Computational Mathematics, 8. Springer-Verlag, New York, 1987.
- [5] R.B. Gandhi, Deepmala and V.N. Mishra, *Local and global results for modified Szász - Mirakjan operators*, Math. Method. Appl. Sci. (2016), DOI:10.1002/mma.4171.
- [6] H.Gonska and I.Rasa, *Asymptotic behaviour of differentiated Bernstein polynomials*, Mat. Vesnik, **61** (2009), 53–60.
- [7] H.Gonska, M.Heilmann and I.Rasa, *Kantorovich operators of order k*, Numer. Funct. Anal. Optimiz. **32** (2011), 717–738.
- [8] M.E.H. Ismail, *Classical and Quantum Orthogonal Polynomials in one Variable*, Cambridge University Press, Cambridge, 2005.
- [9] L.V. Kantorovich, *Sur certains développements suivant les polynômes la forme de S. Bernstein*, I, II, C. R. Acad URSS, (1930) 563–568, 595–600.
- [10] J.P. King, *Positive linear operators which preserves x^2* , Acta Math. Hungar **99** (3) (2003), 203–208.
- [11] G.G. Lorentz, *Mathematical Expositions*, No. 8, Bernstein polynomials, University of Toronto Press, Toronto 1953.
- [12] V.N. Mishra, K. Khatri and L.N. Mishra, *On Simultaneous Approximation for Baskakov-Durrmeyer-Stancu type operators*, Journal of Ultra Scientist of Physical Sciences **24** (3) (2012), 567–577.
- [13] V.N. Mishra, K. Khatri, L.N.Mishra and Deepmala, *Inverse result in simultaneous approximation by Baskakov-Durrmeyer-Stancu operators*, Journal of Inequalities and Applications 2013, 2013:586.doi:10.1186/1029-242X-2013-586.

- [14] V.N. Mishra, H.H. Khan, K. Khatri and L.N. Mishra, *Hypergeometric Representation for Baskakov-Durrmeyer-Stancu Type Operators*, Bulletin of Mathematical Analysis and Applications, **5** (3) (2013), 18–26.
- [15] V.N. Mishra, K. Khatri and L.N. Mishra, *Some approximation properties of q -Baskakov-Beta-Stancu type operators*, Journal of Calculus of Variations, Volume 2013, Article ID 814824, 8 pages. <http://dx.doi.org/10.1155/2013/814824>.
- [16] V.N. Mishra, K. Khatri and L.N. Mishra *Statistical approximation by Kantorovich type Discrete q -Beta operators*, Advances in Difference Equations **2013**, 2013:345, DOI:10.1186/10.1186/1687-1847-2013-345.
- [17] V.N. Mishra, P. Sharma and L.N. Mishra, *On statistical approximation properties of q -Baskakov-Szász-Stancu operators*, Journal of Egyptian Mathematical Society **24** (3) (2016), 396–401. DOI:10.1016/j.joems.2015.07.005.
- [18] V.N. Mishra, R.B.Gandhi and F.Nasierh, *Simultaneous approximation by Szász-Mirakjan-Durrmeyer-type operators*, Bollettino dell'Unione Matematica Italiana **8** (4) (2016), 297–305.
- [19] V.N. Mishra and R.B. Gandhi, *Simultaneous approximation by Szász-Mirakjan-Stancu-Durrmeyer type operators*, Periodica Mathematica Hungarica **74** (1), (2017), 118–127. DOI:10.1007/s10998-016-0145-0.
- [20] R.N. Mohapatra and Z. Walczak, *Remarks on a class of Szász-Mirakjan type operators*, East J. Approx., **15** (2) (2009), 197–206.
- [21] O. Szász, *Generalization of S. Bernstein's polynomials to the infinite interval*, J. Research Nat. Bur. Standards Sci. **45** (3-4) (1950), 239–245.
- [22] V. Totik, *Approximation by Szász-Mirakjan-Kantorovich operators in $L_p(p > 1)$* , Anal. Math. **9** (2) (1983), 147–167.
- [23] A. Wafi and N. Rao, *Stancu-variant of generalized Baskakov operators*, Filomat, (2015) (In Press).
- [24] A. Wafi, N.Rao and D. Rai, *Approximation properties by generalized-Baskakov-Kantorovich-Stancu type operators*, Appl. Math. Inf. Sci. Lett., **4** (3) (2016), 111–118.
- [25] A. Wafi and N. Rao, *Szász-Durrmeyer operators based on Dunkl analogue*, Complex Anal. Oper. Theory, (2017) 1–18. doi:10.1007/s11785-017-0647-7.
- [26] A. Wafi and N. Rao, *A generalization of Szász-type operators which preserves constant and quadratic test functions*, Cogent Mathematics (2016), 3: 1227023.
- [27] S. Varma and F. Taşdelen, *Szász type operators involving Charlier polynomials*, Math. Comput. Modeling, 56 (5-6) (2012) 118–122.

Abdul Wafi

Department of Mathematics
Jamia Millia Islamia
New Delhi 110025, India
E-mail: awafi@jmi.ac.in

Nadeem Rao

Department of Mathematics
Jamia Millia Islamia
New Delhi-110025, India
E-mail: nadeemrao1990@gmail.com

Deepmala

Mathematics Discipline,
PDPM Indian Institute of Information Technology
Design & Manufacturing
Jabalpur, Madhya Pradesh-482005, India
E-mail: dmrai23@gmail.com