

ON SINGULAR INTEGRAL OPERATORS INVOLVING POWER NONLINEARITY

SEVGI ESEN ALMALI, GÜMRAH UYSAL,
VISHNU NARAYAN MISHRA*, AND ÖZGE ÖZALP GÜLLER

ABSTRACT. In the current manuscript, we investigate the pointwise convergence of the singular integral operators involving power nonlinearity given in the following form:

$$T_\lambda(f; x) = \int_a^b \sum_{m=1}^n f^m(t) K_{\lambda, m}(x, t) dt, \quad \lambda \in \Lambda, \quad x \in (a, b),$$

where Λ is an index set consisting of the non-negative real numbers, and $n \geq 1$ is a finite natural number, at μ -generalized Lebesgue points of integrable function $f \in L_1(a, b)$. Here, f^m denotes m -th power of the function f and (a, b) stands for arbitrary bounded interval in \mathbb{R} or \mathbb{R} itself. We also handled the indicated problem under the assumption $f \in L_1(\mathbb{R})$.

1. Introduction

In [17], Taberski presented and proved some theorems about the pointwise approximation properties of functions f in $L_1 \langle -\pi, \pi \rangle$ using a family of convolution type of linear singular integral operators depending

Received June 6, 2017. Revised November 5, 2017. Accepted November 8, 2017.
2010 Mathematics Subject Classification: 41A35, 41A25, 47G10, 47A58.

Key words and phrases: Pointwise convergence, Nonlinear integral operators, μ -generalized Lebesgue point.

* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2017.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

on two parameters of the form:

$$(1.1) \quad L_\lambda(f; x) = \int_{-\pi}^{\pi} f(t) K_\lambda(t-x) dt, \quad x \in \langle -\pi, \pi \rangle, \quad \lambda \in \Omega \subset \mathbb{R}_0^+,$$

where $\langle -\pi, \pi \rangle$ stands for closed, semi-closed or open interval and $K_\lambda(t)$ is the kernel satisfying suitable properties. After this study, the pointwise convergence of the operators of type (1.1) was investigated by Gadjiev [9] and Rydzewska [14] at generalized Lebesgue points and μ -Lebesgue points of functions $f \in L_1 \langle -\pi, \pi \rangle$, respectively. Later on, Karsli and Ibikli [10] generalized some results in [9], [14] and [17], by handling the operators of type (1.1) in the space $L_1 \langle a, b \rangle$. For some studies on linear singular integral operators in miscellaneous settings, the reader may see also [3], [11] and [7].

Musiela [12] investigated the approximation properties of nonlinear integral operators in the following setting:

$$(1.2) \quad T_w f(y) = \int_G K_w(x-y; f(x)) dx, \quad y \in G, \quad w \in \Lambda,$$

where G is a locally compact Abelian group enriched with Haar measure, and Λ is a non-empty index set with desired topology, by reconciling K_w with the Lipschitz condition on its second variable. Then, Musielak [13] advanced his previous sounding analysis [12] by considering the indicated operators in generalized Orlicz spaces. Afterwards, Swiderski and Wachnicki [16] investigated the pointwise convergence of the operators of type (1.2) in accordance with two parameters at Lebesgue points of the functions $f \in L_p(G)$. For some studies related to mentioned studies above, we refer the reader to see [1], [2], [4], [5], [6], [8] and [19].

Let $L_1(a, b)$ be the space of all measurable functions f for which $|f|$ is integrable on the arbitrary bounded intervals (a, b) of \mathbb{R} . Similarly, let $L_1(\mathbb{R})$ be the space of all measurable functions f for which $|f|$ is integrable on \mathbb{R} . The norms on these spaces are given by $\|f\|_{L_1(a,b)} = \int_a^b |f(t)| dt$ and $\|f\|_{L_1(\mathbb{R})} = \int_{\mathbb{R}} |f(t)| dt$ (see, for example, [18]). The current manuscript can be seen as a generalization and a continuation of [1]. The main aim of this paper is to obtain pointwise convergence of singular

integral operators involving power nonlinearity in the following form:

$$(1.3) \quad T_\lambda(f; x) = \int_a^b \sum_{m=1}^n f^m(t) K_{\lambda,m}(x, t) dt, \quad \lambda \in \Lambda, \quad x \in (a, b),$$

where Λ is an index set consisting of the non-negative real numbers with accumulation point λ_0 (or $\lambda_0 = \infty$) and $n \geq 1$ is a finite natural number, to the function $f \in L_1(a, b)$ at its μ -generalized Lebesgue points as λ tends to λ_0 . On the other hand, we also handled the indicated problem under the assumption $f \in L_1(\mathbb{R})$. Here, f^m denotes m -th power of the function and (a, b) denotes arbitrary bounded interval in \mathbb{R} or \mathbb{R} itself. Note that taking $n = 1$ transforms the operators of type (1.3) into linear form. Also, the rate of convergences are discussed. The operators of type (1.3) are the special type of nonlinear integral operators building the bridge between summation type operators and integral type operators. Therefore, doing a research on their approximation properties is remarkable from the theoretical point of view.

The paper is organized as follows: In Section 2, we present preliminary concepts. In Section 3, we prove main results. In Section 4, we deal with the rates of pointwise convergences.

2. Preliminaries

Now, we start by stating the following definition which is the heart of the matter.

DEFINITION 2.1. Let δ_1 be a given fixed positive real number and $\delta_1 > h > 0$. A point $x_0 \in (a, b)$ (or $x_0 \in \mathbb{R}$) is called μ -generalized Lebesgue point of the function $f \in L_1(a, b)$ (or $f \in L_1(\mathbb{R})$), if the following relation:

$$\lim_{h \rightarrow 0} \frac{1}{\mu(h)} \int_0^h |f(t + x_0) - f(x_0)| dt = 0$$

(also, when the integral is taken from $-h$ to 0) holds, where $\mu : \mathbb{R} \rightarrow \mathbb{R}$ is an increasing and absolutely continuous function on $[0, \delta_1]$ with $\mu(0) = 0$.

REMARK 2.2. For the ideas and concepts used in Definition 2.1, we refer the reader to see [9], [10] and [14]. On the other hand, if we take $\mu(t) = t$, then we obtain the well-known definition of Lebesgue point.

Now, in view of Definition 1 in [1], we may express the following definition which can be seen as a generalization of the properties of usual approximate identities, such as well-known Gauss-Weierstrass kernel.

DEFINITION 2.3. (Class A) Let Λ be an index set consisting of the non-negative real numbers with accumulation point λ_0 (or $\lambda_0 = \infty$). Keeping in mind that the natural number n is finite, for $m = 1, \dots, n$, a family $\{K_{\lambda,m}\}_{\lambda \in \Lambda}$ consisting of the globally integrable functions $K_{\lambda,m}(x, t) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}_0^+$, is called Class A, if the following conditions hold there:

- a. For any fixed $x \in \mathbb{R}$ and for $m = 1, \dots, n$,

$$\lim_{\lambda \rightarrow \lambda_0} \left| \int_{\mathbb{R}} K_{\lambda,m}(x, t) dt - C_m \right| = 0,$$

where $C_m > 0$ are certain real numbers.

- b. For any fixed $x \in \mathbb{R}$, for $m = 1, \dots, n$ and for every $\xi > 0$,

$$\lim_{\lambda \rightarrow \lambda_0} \left[\sup_{|x-t| \geq \xi} K_{\lambda,m}(x, t) \right] = 0.$$

- c. For any fixed $x \in \mathbb{R}$, for $m = 1, \dots, n$ and for every $\xi > 0$,

$$\lim_{\lambda \rightarrow \lambda_0} \left[\int_{|x-t| \geq \xi} K_{\lambda,m}(x, t) dt \right] = 0.$$

- d. Let δ_0 satisfying $0 < \delta_0 \leq \delta_1$ be a given fixed real number such that $K_{\lambda,m}(x, t)$ is non-decreasing on $[x - \delta_0, x]$ and non-increasing on $[x, x + \delta_0]$ with respect to t , for any fixed λ and $x \in \mathbb{R}$ and for $m = 1, \dots, n$.

Throughout this paper $K_{\lambda,m}$ belongs to Class A.

3. Main Results

Our first main theorem in which (a, b) denotes arbitrary bounded interval in \mathbb{R} is as follows:

THEOREM 3.1. *If $x_0 \in (a, b)$ is a μ -generalized Lebesgue point of function $f \in L_1(a, b)$ and f is bounded on (a, b) , then*

$$\lim_{\lambda \rightarrow \lambda_0} T_\lambda(f; x_0) = \sum_{m=1}^n C_m f^m(x_0)$$

on any set Z on which the function

$$\sum_{m=1}^n \int_{x_0-\delta}^{x_0+\delta} K_{\lambda,m}(x_0, t) \left| \{\mu(|x_0 - t|)\}'_t \right| dt$$

is bounded as λ tends to λ_0 .

Proof. We start by setting $I_\lambda = \left| T_\lambda(f; x_0) - \sum_{m=1}^n C_m f^m(x_0) \right|$.

Next, we define a function $g^m : \mathbb{R} \rightarrow \mathbb{R}$ ($m = 1, \dots, n$) such that

$$g^m(t) = \begin{cases} f^m(t), & t \in (a, b), \\ 0, & t \in \mathbb{R} \setminus (a, b). \end{cases}$$

By condition (a), we can write

$$\begin{aligned} I_\lambda &\leq \int_a^b \sum_{m=1}^n |f^m(t) - f^m(x_0)| K_{\lambda,m}(x_0, t) dt \\ &\quad + \sum_{m=1}^n |f^m(x_0)| \int_{\mathbb{R} \setminus (a,b)} K_{\lambda,m}(x_0, t) dt \\ &\quad + \sum_{m=1}^n |f^m(x_0)| \left| \int_{\mathbb{R}} K_{\lambda,m}(x_0, t) dt - C_m \right| \\ &= I_1 + I_2 + I_3. \end{aligned}$$

It is sufficient to show that the terms on the right hand side of the last inequality tend to zero as λ tends to λ_0 . By conditions (c) and (a), I_2 and I_3 tend to zero as λ tends to λ_0 , respectively.

Now, we consider I_1 . For a fixed real number δ satisfying $0 < \delta < \delta_0$, we split the integral I_1 into four terms as follows:

$$\begin{aligned}
I_1 &= \left[\int_a^{x_0-\delta} + \int_{x_0-\delta}^{x_0} + \int_{x_0}^{x_0+\delta} + \int_{x_0+\delta}^b \right] \sum_{m=1}^n |f^m(t) - f^m(x_0)| K_{\lambda,m}(x_0, t) dt \\
&= I_{11} + I_{12} + I_{13} + I_{14}.
\end{aligned}$$

Since n is finite, we can write I_{11} in the following form:

$$I_{11} = \sum_{m=1}^n \int_a^{x_0-\delta} |f^m(t) - f^m(x_0)| K_{\lambda,m}(x_0, t) dt.$$

From the condition (d), we have

$$\begin{aligned}
I_{11} &\leq \sum_{m=1}^n K_{\lambda,m}(x_0, x_0 - \delta) \left\{ \int_a^{x_0-\delta} |f^m(t)| dt + \int_a^{x_0-\delta} |f^m(x_0)| dt \right\} \\
&\leq \sum_{m=1}^n K_{\lambda,m}(x_0, x_0 - \delta) \left\{ \|f^m\|_{L_1(a,b)} + |f^m(x_0)| (b - a) \right\}.
\end{aligned}$$

Using same method, we can estimate I_{14} . Using condition (d), we shall write

$$\begin{aligned}
I_{14} &\leq \sum_{m=1}^n K_{\lambda,m}(x_0, x_0 + \delta) \left\{ \int_{x_0+\delta}^b |f^m(t)| dt + \int_{x_0+\delta}^b |f^m(x_0)| dt \right\} \\
&\leq \sum_{m=1}^n K_{\lambda,m}(x_0, x_0 + \delta) \left\{ \|f^m\|_{L_1(a,b)} + |f^m(x_0)| (b - a) \right\}.
\end{aligned}$$

By condition (b) of class A and boundedness of f , $I_{11} \rightarrow 0$ and $I_{14} \rightarrow 0$ as λ tends to λ_0 .

On the other hand, since $x_0 \in (a, b)$ is a μ -generalized Lebesgue point of f , for every $\varepsilon > 0$, there exists a number $\delta > 0$ such that

$$(3.1) \quad \int_{x_0}^{x_0+h} |f(t) - f(x_0)| dt < \varepsilon \mu(h)$$

and

$$(3.2) \quad \int_{x_0-h}^{x_0} |f(t) - f(x_0)| dt < \varepsilon\mu(h)$$

hold for every h satisfying $0 < h \leq \delta$.

Now, we define a new function with respect to relation (3.1) as follows:

$$F(t) = \int_{x_0}^t |f(u) - f(x_0)| du.$$

Then, for $t - x_0 \leq \delta$, we have

$$|F(t)| \leq \varepsilon\mu(t - x_0).$$

To go further, we need the following identity (see, for example, [15]):

$$(q^m - r^m) = (q - r) (q^{m-1} + q^{m-2}r + \dots + r^{m-1}),$$

where $q, r \in \mathbb{R}$. Using this identity and depending on supremum value of f on overall (a, b) , we see that there are finite real numbers $B_m > 0$ such that the inequality

$$|f^m(t) - f^m(x_0)| \leq B_m |f(t) - f(x_0)|,$$

where $m = 1, \dots, n$, holds. Therefore, we can estimate I_{13} as follows:

$$\begin{aligned} |I_{13}| &= \sum_{m=1}^n \left| \int_{x_0}^{x_0+\delta} |f^m(t) - f^m(x_0)| K_{\lambda,m}(x_0, t) dt \right| \\ &\leq \sum_{m=1}^n B_m \int_{x_0}^{x_0+\delta} K_{\lambda,m}(x_0, t) |dF(t)|. \end{aligned}$$

Applying integration by parts leads to the following result:

$$|I_{13}| \leq \sum_{m=1}^n B_m \left\{ F(x_0 + \delta) K_{\lambda,m}(x_0, x_0 + \delta) + \int_{x_0}^{x_0+\delta} F(t) d_t (-K_{\lambda,m}(x_0, t)) \right\}.$$

Since $K_{\lambda,m}$ is non-increasing on $[x_0, x_0 + \delta]$, $-K_{\lambda,m}$ is non-decreasing there. Hence, its differential is positive. Then, we can write

$$|I_{13}| \leq \sum_{m=1}^n B_m \left\{ \varepsilon \mu(\delta) K_{\lambda,m}(x_0, x_0 + \delta) + \varepsilon \int_{x_0}^{x_0+\delta} \mu(t - x_0) d_t (-K_{\lambda,m}(x_0, t)) \right\}.$$

Using integration by parts again, we get the following inequality:

$$|I_{13}| \leq \varepsilon \sum_{m=1}^n B_m \int_{x_0}^{x_0+\delta} K_{\lambda,m}(x_0, t) \left| \{\mu(t - x_0)\}'_t \right| dt.$$

In view of relation (3.2), we obtain the following inequality for the integral I_{12} :

$$|I_{12}| \leq \varepsilon \sum_{m=1}^n B_m \int_{x_0-\delta}^{x_0} K_{\lambda,m}(x_0, t) \left| \{\mu(x_0 - t)\}'_t \right| dt.$$

Setting $B = \max \{B_1, \dots, B_n\}$ and combining corresponding findings for the integrals I_{12} and I_{13} , we get

$$|I_{12}| + |I_{13}| \leq \varepsilon B \sum_{m=1}^n \int_{x_0-\delta}^{x_0+\delta} K_{\lambda,m}(x_0, t) \left| \{\mu(|x_0 - t|)\}'_t \right| dt.$$

Since the right hand side of the above inequality is bounded by the hypothesis, the assertion follows, that is

$$\lim_{\lambda \rightarrow \lambda_0} T_\lambda(f; x_0) = \sum_{m=1}^n C_m f^m(x_0).$$

Thus, the proof is completed. □

In the following theorem, we suppose that $(a, b) = (-\infty, \infty)$.

THEOREM 3.2. *If $x_0 \in \mathbb{R}$ is a μ -generalized Lebesgue point of function $f \in L_1(\mathbb{R})$ and f is bounded on \mathbb{R} , then*

$$\lim_{\lambda \rightarrow \lambda_0} T_\lambda(f; x_0) = \sum_{m=1}^n C_m f^m(x_0)$$

on any set Z on which the function

$$\sum_{m=1}^n \int_{x_0-\delta}^{x_0+\delta} K_{\lambda,m}(x_0, t) \left| \{\mu(|x_0 - t|)\}'_t \right| dt$$

is bounded as λ tends to λ_0 .

Proof. Set $J_\lambda = \left| T_\lambda(f; x_0) - \sum_{m=1}^n C_m f^m(x_0) \right|$. Following the similar strategy as in the previous proof, we have

$$\begin{aligned} J_\lambda \leq & \sum_{m=1}^n K_{\lambda,m}(x_0, x_0 - \delta) \|f^m\|_{L_1(\mathbb{R})} + \sum_{m=1}^n |f^m(x_0)| \int_{-\infty}^{x_0-\delta} K_{\lambda,m}(x_0, t) dt \\ & + \sum_{m=1}^n K_{\lambda,m}(x_0, x_0 + \delta) \|f^m\|_{L_1(\mathbb{R})} + \sum_{m=1}^n |f^m(x_0)| \int_{x_0+\delta}^{\infty} K_{\lambda,m}(x_0, t) dt \\ & + \sum_{m=1}^n |f^m(x_0)| \left| \int_{-\infty}^{\infty} K_{\lambda,m}(x_0, t) dt - C_m \right| \\ & + \varepsilon B^* \sum_{m=1}^n \int_{x_0-\delta}^{x_0+\delta} K_{\lambda,m}(x_0, t) \left| \{\mu(|x_0 - t|)\}'_t \right| dt, \end{aligned}$$

where the number B^* corresponds to number B in previous proof having similar construction process. Now, the result is obvious by the hypotheses. This completes the proof. \square

4. Rate of Convergence

THEOREM 4.1. *Suppose that the hypotheses of Theorem 3.1 are satisfied. Let*

$$\Delta(\lambda, \delta) = \sum_{m=1}^n \int_{x_0-\delta}^{x_0+\delta} K_{\lambda,m}(x_0, t) \left| \{\mu(|x_0 - t|)\}'_t \right| dt,$$

where $0 < \delta < \delta_0$ and $m = 1, \dots, n$, and the following conditions are satisfied:

- (i) $\Delta(\lambda, \delta) \rightarrow 0$ as λ tends to λ_0 , for some $\delta > 0$.
(ii) For every $\xi > 0$,

$$\sup_{|x_0-t| \geq \xi} K_{\lambda,m}(x_0, t) = o(\Delta(\lambda, \delta))$$

as λ tends to λ_0 .

- (iii) For every $\xi > 0$,

$$\int_{|x_0-t| \geq \xi} K_{\lambda,m}(x_0, t) dt = o(\Delta(\lambda, \delta))$$

as λ tends to λ_0 .

- (iv) $\left| \int_{\mathbb{R}} K_{\lambda,m}(x_0, t) dt - C_m \right| = o(\Delta(\lambda, \delta))$
as λ tends to λ_0 .

Then, at each μ -generalized Lebesgue point of function $f \in L_1(a, b)$, we have

$$\left| T_\lambda(f; x_0) - \sum_{m=1}^n C_m f^m(x_0) \right| = o(\Delta(\lambda, \delta)).$$

Proof. The result is obvious by the hypotheses of Theorem 3.1 and Class A conditions. Thus, the proof is omitted. \square

REMARK 4.2. The above result is also valid for the case $f \in L_1(\mathbb{R})$ under the hypotheses of Theorem 3.2.

References

- [1] S. E. Almali, *On approximation properties for non-linear integral operators*, New Trends Math. Sci. **5** (4) (2017), 123–129. (arXiv:1702.04190v1)
- [2] S. E. Almali and A. D. Gadjiev, *On approximation properties of certain multidimensional nonlinear integrals*, J. Nonlinear Sci. Appl. **9** (5) (2016), 3090–3097.
- [3] C. Bardaro, *On approximation properties for some classes of linear operators of convolution type*, Atti Sem. Mat. Fis. Univ. Modena **33** (2) (1984), 329–356.
- [4] C. Bardaro and G. Vinti, *On approximation properties of certain nonconvolution integral operators*, J. Approx. Theory, **62** (3) (1990), 358–371.
- [5] C. Bardaro, J. Musielak and G. Vinti, *Approximation by nonlinear integral operators in some modular function spaces*, Ann. Polon. Math. **63** (2) (1996), 73–182.
- [6] C. Bardaro, J. Musielak and G. Vinti, *Nonlinear Integral Operators and Applications*, de Gruyter Series in Nonlinear Analysis and Applications, 9., Walter de Gruyter & Co., Berlin, 2003.

- [7] P. L. Butzer and R. J. Nessel, *Fourier Analysis and Approximation*, vol. I. Academic Press, New York, London, 1971.
- [8] A. D. Gadjiev, *On nearness to zero of a family of nonlinear integral operators of Hammerstein*, Izv. Akad. Nauk Azerbaïdžan. SSR Ser. Fiz.-Tehn. Mat. Nauk, (1966), no. 2, 32–34.
- [9] A. D. Gadjiev, *The order of convergence of singular integrals which depend on two parameters*, *Special Problems of Functional Analysis and their Appl. to the Theory of Diff. Eq. and the Theory of Func.*, Izdat. Akad. Nauk Azerbaïdžan. SSR., (1968), 40–44.
- [10] H. Karsli and E. Ibikli, *On convergence of convolution type singular integral operators depending on two parameters*, Fasc. Math. **38** (2007), 25–39.
- [11] R. G. Mamedov, *On the order of convergence of m -singular integrals at generalized Lebesgue points and in the space $L_p(-\infty, \infty)$* , Izv. Akad. Nauk SSSR Ser. Mat. **27** (2) (1963), 287–304.
- [12] J. Musielak, *On some approximation problems in modular spaces*, Constructive Function Theory 1981, (Proc. Int. Conf., Varna, June 1-5, 1981), Publ. House Bulgarian Acad. Sci., Sofia, (1983), 455–461.
- [13] J. Musielak, *Approximation by nonlinear singular integral operators in generalized Orlicz spaces*, Comment. Math. Prace Mat. **31** (1991), 79–88.
- [14] B. Rydzewska, *Approximation des fonctions par des intégrales singulières ordinaires*, Fasc. Math. **7** (1973), 71–81.
- [15] M. D. Spivak, *Calculus (3rd ed.)*, Publish or Perish, Inc., Houston, Texas, 1994.
- [16] T. Swiderski and E. Wachnicki, *Nonlinear singular integrals depending on two parameters*, Comment. Math. **40** (2000), 181–189.
- [17] R. Taberski, *Singular integrals depending on two parameters*, Prace Mat. **7** (1962), 173–179.
- [18] R. Taberski, *Exponential approximation on the real line*, Approximation and function spaces (Warsaw, 1986), Banach Center Publ., PWN, Warsaw, **22** (1989), 449–464.
- [19] G. Uysal, V. N. Mishra, O. O. Guller and E. Ibikli, *A generic research on nonlinear non-convolution type singular integral operators*, Korean J. Math. **24** (3) (2016), 545–565.

Sevgi Esen Almalı

Department of Mathematics

Kırıkkale University

Kırıkkale 71451, Turkey

E-mail: sevgi_esen@hotmail.com**Gümrah Uysal**

Department of Computer Technologies

Division of Technology of Information Security

Karabük University

Karabük 78050, Turkey

E-mail: fgumrahuysal@gmail.com**Vishnu Narayan Mishra**^{*1}Department of Mathematics

Indira Gandhi National Tribal University

Lalpur, Amarkantak 484 887, Madhya Pradesh, India

^{*2}L. 1627 Awadh Puri Colony Beniganj

Phase - III, Opposite - Industrial Training Institute (I.T.I.)

Ayodhya Main Road, Faizabad 224 001, U.P., India

E-mail: vishnunarayanmishra@gmail.com**Özge Özalp Güller**

Department of Mathematics

Ankara University

Ankara 06100, Turkey

E-mail: ozgeguller2604@gmail.com