

## SURFACES FOLIATED BY ELLIPSES WITH CONSTANT GAUSSIAN CURVATURE IN EUCLIDEAN 3-SPACE

AHMED T. ALI AND FATHI M. HAMDOON

ABSTRACT. In this paper, we study the surfaces foliated by ellipses in three dimensional Euclidean space  $\mathbf{E}^3$ . We prove the following results: **(1)** The surface foliated by an ellipse have constant Gaussian curvature  $K$  if and only if the surface is flat, i.e.  $K = 0$ . **(2)** The surface foliated by an ellipse is a flat if and only if it is a part of generalized cylinder or part of generalized cone.

### 1. Introduction

A cyclic surface in Euclidean space  $\mathbf{E}^3$  is a surface determined by a smooth uni-parametric family of pieces of circles [4, 5]. The first example of cyclic surface is a surface of revolution, that is, a surface which is stable under a group of rotations that leave a straight-line point wise fixed. The Riemann examples play a major role in the theory of minimal surfaces [10]. Enneper [4, 5] proved that for a cyclic minimal surface, the planes containing the circles must be parallel and it is one of the examples obtained by Riemann. Nitsche [9] Proved that the surface must be an open set of a sphere or, in non-spherical case, the circles must be lie in parallel planes. In the latter case, the only possibilities are the surfaces of revolution discovered by Delaunay [3].

---

Received September 21, 2017. Revised November 20, 2017. Accepted November 21, 2017.

2010 Mathematics Subject Classification: 53A05, 53A17.

Key words and phrases: Surfaces in Euclidean space, Gaussian curvature.

© The Kangwon-Kyungki Mathematical Society, 2017.

This is an Open Access article distributed under the terms of the Creative commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

Let  $\Gamma = \Gamma(u)$  be an orthogonal smooth curve to each  $u$ -plane of the foliation and denote by  $u$  its arc-length parameter. We assume that the planes of the foliation are not parallel and let  $\{\mathbf{t}, \mathbf{n}, \mathbf{b}\}$  be a moving frame of the curve  $\Gamma$ , where  $\mathbf{t}$ ,  $\mathbf{n}$  and  $\mathbf{b}$  denote the tangent, principal normal and binormal vectors respectively. Locally we parameterize the surface  $M(u, v)$  by

$$(1) \quad \mathbf{X}(u, v) = \mathbf{C}(u) + r(u) \left( a \cos[v] \mathbf{n} + b \sin[v] \mathbf{b} \right), \quad v \in [0, 2\pi], \quad a \neq b \in \mathbb{R}$$

where  $r = r(u) > 0$  and  $\mathbf{C} = \mathbf{C}(u)$  denote the center of each ellipse of the foliation, the Frenet equations of the curve  $\Gamma$  are

$$(2) \quad \begin{cases} \mathbf{t}' = \kappa \mathbf{n}. \\ \mathbf{n}' = -\kappa \mathbf{t} + \tau \mathbf{b}. \\ \mathbf{b}' = -\tau \mathbf{n}. \end{cases}$$

where the prime  $'$  denotes the derivative with respect to the  $u$ -parameter,  $\kappa$  and  $\tau$  are the curvature and torsion of  $\Gamma$  respectively. We assume that  $\kappa \neq 0$  because  $\Gamma$  is not straight line and let

$$(3) \quad \mathbf{C}'(u) = \alpha \mathbf{t} + \beta \mathbf{n} + \gamma \mathbf{b}$$

where  $\alpha$ ,  $\beta$  and  $\gamma$  are smooth functions on  $u$ .

Without loss of generality, we can assume that  $a = 1$ . It is worth noting that when  $b = 1$  the surface (1) is a cyclic surface foliated by a smooth one-parameter family of circles in three dimensional Euclidean space  $\mathbf{E}^3$  which studied in Lopez [6–8]. In [6] he studied a surface in Euclidean three space  $\mathbf{E}^3$  with constant Gauss curvature foliated by circles. In [7, 8] he studied surfaces in Euclidean 3-space that satisfy a special Linear Weingarten (LW) condition of linear type as  $\kappa_1 = c_1 \kappa_2 + c_2$  and  $c_3 H + c_4 K = c_5$ , where  $c_i$   $i = 1, 2, \dots, 5$  are real numbers and  $\kappa_1$  and  $\kappa_2$  denote the principal curvatures while  $H$  and  $K$  denotes the the mean and Gaussian curvatures at each point of the surface. Also, he proved that:

**(1):** All cyclic surface with non-zero constant Gauss curvature must be a surface of revolution [6].

**(2):** All cyclic LW-surface with  $\kappa_1 = c_1 \kappa_2$  must be of Riemann type [7].

**(3):** All cyclic LW-surface with  $c_3H + c_4K = c_5$  must be a surface of revolution, a Riemann minimal surface or a generalized cone [8].

In this paper, we will study the surface foliated by an ellipse. In this case, the surface takes the form (1) such that  $b \neq 1$ . We will prove the following main results for constant Gaussian curvature:

**THEOREM 1.1.** *The surface (1) foliated by an ellipse in  $\mathbf{E}^3$  is a flat if and only if it is a part of generalized cylinder or part of generalized cone.*

**THEOREM 1.2.** *The surface (1) foliated by an ellipse has constant Gaussian curvature  $K$  in  $\mathbf{E}^3$  if and only if  $K = 0$ .*

On the other hand, Let  $M$  be a surface in  $\mathbf{R}^3$  foliated by a pieces of ellipses in parallel planes. Without loss of generality, we assume that the planes of the foliation are parallel to  $x_1x_2$ -plane. Let

$$(4) \quad \mathbf{X}(u, v) = \left( f(u) + r(u) \cos[v], g(u) + br(u) \sin[v], u \right), \quad u \in I, v \in J,$$

be a local parametrization of  $M$ . Then, we have the following theorem:

**THEOREM 1.3.** *Let  $M$  be a surface (4) in  $\mathbf{E}^3$  with constant Gaussian curvature  $K = K_0$  and foliated by pieces of ellipses in parallel planes. Then*

- (1):**  $K_0 = 0$ .
- (2):**  $M$  must be parameterized, up a rigid motion of  $\mathbf{E}^3$ , as

$$(5) \quad \mathbf{X}(u, v) = \left( f_1 u + f_0, g_1 u + g_0, u \right) + \left( r_1 u + r_0 \right) \left( \cos[v], b \sin[v], 0 \right),$$

where  $f_0, f_1, g_0, g_1, r_0, r_1, b \in \mathbf{R}$ .

As a corollary of both theorem 1.2 and 1.3, we obtain:

**COROLLARY 1.4.** *All surfaces foliated by an ellipses with constant Gauss curvatures must be a surfaces of revolution.*

## 2. Gaussian curvatures

Consider  $M$  a surface in  $\mathbf{E}^3$  parameterized by  $\mathbf{X} = \mathbf{X}(u, v)$  and let  $\mathbf{U}$  denote the unit normal vector field on  $M$ . The tangent vectors to the

parametric curves of the surface  $\mathbf{X}(u, v)$  are

$$\mathbf{X}_u = \frac{\partial \mathbf{X}}{\partial u}, \quad \mathbf{X}_v = \frac{\partial \mathbf{X}}{\partial v},$$

and the unit normal on the surface is given by

$$\mathbf{U} = \frac{\mathbf{X}_u \times \mathbf{X}_v}{|\mathbf{X}_u \times \mathbf{X}_v|}$$

where  $\times$  denotes the cross product of  $\mathbf{E}^3$ . The first fundamental quadratic form on the surface is

$$I = \langle d\mathbf{X}, d\mathbf{X} \rangle = E du^2 + 2F du dv + G dv^2$$

with the first fundamental coefficients

$$E = \langle \mathbf{X}_u, \mathbf{X}_u \rangle, \quad F = \langle \mathbf{X}_u, \mathbf{X}_v \rangle, \quad G = \langle \mathbf{X}_v, \mathbf{X}_v \rangle.$$

The second fundamental quadratic form is given by

$$II = \langle \mathbf{U}, d^2\mathbf{X} \rangle = e du^2 + 2f du dv + g dv^2,$$

with second fundamental coefficients

$$e = \langle \mathbf{U}, \mathbf{X}_{uu} \rangle, \quad f = \langle \mathbf{U}, \mathbf{X}_{uv} \rangle, \quad g = \langle \mathbf{U}, \mathbf{X}_{vv} \rangle.$$

Under this parameterization of the surface  $\mathbf{X} = \mathbf{X}(u, v)$ , the Gaussian curvature  $K$  is

$$(6) \quad K = \frac{eg - f^2}{EG - F^2}.$$

The proof of our results depend on we can reduce the equation  $K = \text{constant}$  to an expression as a linear combination of the trigonometric function  $\{\cos[i v], \sin[i v]\}$ ,  $i \in \mathbb{N}$ , namely,

$$(7) \quad \sum_{i=0}^8 \left( E_i \cos[i v] + F_i \sin[i v] \right) = 0$$

and  $E_i$  and  $F_i$  are functions on the variable  $u$ . In particular, the coefficients must vanish. The work then is to compute explicitly these coefficients  $E_i$  and  $F_i$  by successive manipulations. The author were able to obtain the results using the symbolic program Mathematica to check their work. The computer was used in each calculation several times, giving understandable expressions of the coefficients  $E_i$  and  $F_i$ .

Although the explicit computation of the Gaussian curvature  $K$  can be obtained, for example, by using the Mathematica programme, its

expression is some cumbersome. However, the key in our proofs lies that one can write  $K$  as

$$(8) \quad K = \frac{\mathcal{P}(\cos[i v], \sin[i v])}{\mathcal{Q}(\cos[i v], \sin[i v])} = \frac{\sum_{i=0}^4 (A_i \cos[i v] + B_i \sin[i v])}{\sum_{i=0}^8 (C_i \cos[i v] + D_i \sin[i v])}.$$

The assumption of the constancy of the scalar curvature  $K$  implies that (8) converts into

$$(9) \quad K \mathcal{Q}(\cos[i v], \sin[i v]) - \mathcal{P}(\cos[i v], \sin[i v]) = 0.$$

Equation (9) means that if we write it as a linear combination of the functions  $\{\cos[i v], \sin[i v]\}$  namely, (7), the corresponding coefficients must vanish. From here, we will be able to describe all surfaces foliated by an ellipses with constant Gaussian curvature. As we will see, it is not necessary to give the (long) expression of  $K$  but only the coefficients of higher order for the trigonometric functions.

We distinguish the cases  $K = 0$  and  $K \neq 0$ . Without loss of generality, we can put  $\tau(u) = \lambda(u) \kappa(u)$  and  $\beta(u) = \mu(u) \kappa(u)$ , where  $\lambda$  and  $\mu$  are functions of  $u$ .

### 3. Proof Theorem 1.1

In this section we assume that  $K = 0$  on the surface  $\mathbf{X}(u, v)$ . From (9), we have

$$\mathcal{P}(\cos[i v], \sin[i v]) = \sum_{i=0}^4 (A_i \cos[i v] + B_i \sin[i v]) = 0.$$

Then the work consists in the explicit computations of the coefficients  $A_i$  and  $B_i$ . We distinguish different cases that fill all possible cases. Since

$$(10) \quad B_4 = \frac{b r^2 \kappa^3}{2} \left[ 2 \mu \gamma - (1 - b^2)(r^2 \lambda' - 2 \lambda \mu \alpha) \right]$$

must be vanished, then we have two possibilities for  $\mu$  as the following:

**3.1. When  $\mu \neq 0$  then**  $\gamma = \frac{(b^2 - 1) (2 \lambda \mu \alpha - r^2 \lambda')}{2 \mu}$ . A straightforward computation shows that:

$$(11) \quad A_3 = -b^2 \kappa^3 \left[ 3 f \alpha - (4 \mu r' - r \mu') r \right],$$

where  $f = [(b^2 - 1)\lambda^2 - 1]r^2$ . According to the condition  $A_3 = 0$ , let us distinguish two possibilities of  $f$  as as the following:

**3.1.1.** When  $f \neq 0$  then  $\alpha = \frac{(4\mu r' - r\mu')r}{3f}$ . From  $B_2 = 0$  and  $B_3 = 0$ , their imply

$$(12) \quad \left\{ \begin{array}{l} (b^2 - 1)\kappa f r^2 (2f r' - r f') \mu'' = 4f \kappa^3 \mu \left[ b^2 f (3r^2 r' - 4\mu^2 r' + r\mu\mu') \right. \\ \quad \left. + b^2 r^2 \mu (4\mu r' - r\mu') - 3f^2 r' \right] + (1 - b^2) \left[ 6f^3 \kappa^3 \mu r' \right. \\ \quad \left. + 4\kappa f' (r\mu' - 4\mu r') \left[ 8\mu^2 r' + r(f' - 2\mu\mu') \right] + f^2 \left( 40\kappa\mu r 6f^3 \right. \right. \\ \quad \left. \left. - 2r^2 r' \kappa' \mu' + r \left[ 3\mu\kappa^3 f' + 8\mu\kappa' r'^2 - 8\kappa r' (2r' \mu' + \mu r'') \right] \right) \right. \\ \quad \left. + f \left( 64\kappa\mu^3 r'^3 + r^3 \kappa' \mu' f' - 4\kappa\mu r r'^2 [3f' + 8\mu\mu'] \right. \right. \\ \quad \left. \left. + r^2 \left[ 6\kappa\mu' r' f' + 4\mu \left( \kappa r' \mu'^2 + f' [\kappa r' - r' \kappa'] \right) \right] \right) \right], \end{array} \right.$$

$$(13) \quad \left\{ \begin{array}{l} 2\kappa\mu f r^2 (f - r^2) f'' = 4\kappa\mu^2 r^3 f' (r\mu' - 4\mu r') \\ \quad + r f \left[ \mu \left( \kappa [r f'^2 + 32r\mu^2 r'^2 + 4f' r' (4\mu^2 - 3r^2)] - 2r^3 f' \kappa' \right) \right. \\ \quad \left. - 2\kappa r \mu' [f' (r^2 + 2\mu^2) + 4\mu^2 r r'] \right] \\ \quad + 4f^3 \left( r\kappa\mu r'' - r [r\mu\kappa' + \kappa (4\mu r' + r\mu')] \right) \\ \quad + 2f^2 \left[ 4\kappa\mu r r' (f' + \mu\mu') - 16\kappa\mu^3 r'^2 + r^2 (10\kappa\mu r'^2 + \mu\kappa f' \right. \\ \quad \left. + \kappa\mu' f') + 2r^3 [r' (\mu\kappa' + \kappa\mu') - \kappa\mu r''] \right]. \end{array} \right.$$

From the above equation, we will consider two cases:

**3.1.1.1**  $f \neq f_0 r^2$ , for all arbitrary constant  $f_0$ . If we substitute  $\mu''$  and  $f''$  from (12) and (13), in the coefficient  $B_1$  we obtain the following condition:

(14)

$$(2 f r' - r f') \left[ (r \mu' - 4 \mu r') \mu - 3 f r' \right] \left[ (r \mu' - 4 \mu r')^2 - 9 f^2 \kappa^2 \right] = 0.$$

The above condition reduces to two solutions:  $\mu' = \frac{4 \mu r' \pm 3 \kappa f}{r}$  and

$$\mu' = \frac{(3 f + 4 \mu^2) r'}{\mu r}.$$

**(I):**  $\mu' = \frac{4 \mu r' + 3 \kappa f}{r}$ . If we substitute  $\mu'$  from this equation and  $\mu''$  from (12), in the condition  $\mu'' - \frac{d\mu'}{du} = 0$ , then we can obtain the following condition:

(15)  $(\kappa \mu - r') \left[ 5 (b^2 - 1) (r f' - 2 f r') + 4 b^2 \kappa \mu (f - r^2) \right] = 0.$

The above condition yields two solutions:

**(I-A):**  $f' = 2 \left[ \frac{f r'}{r} - \frac{2 b^2 \kappa \mu (f - r^2)}{5 (b^2 - 1) r} \right]$ . Again the condition  $f'' - \frac{df'}{du} = 0$  reduced that:

(16)  $(f - r^2) \left[ 2 (14 b^2 - 15) \kappa \mu - 15 (b^2 - 1) r' \right] = 0.$

Because  $f \neq r^2$ , the above condition leads to  $r' = \frac{2 (14 b^2 - 15) \kappa \mu}{15 (b^2 - 1)}$ . Now, a straightforward computation shows that, the remain coefficients are:

$$(17) \quad \begin{cases} A_0 = \frac{b^2 r \kappa^4}{50(b^2 - 1)} \left[ 57 b^2 r^2 - 9(218 b^2 - 225) f - \frac{(51 b^2 - 55)(71 b^2 - 75) \mu^2}{b^2 - 1} \right], \\ A_1 = \frac{4 b^2 r \kappa^4}{45(b^2 - 1)} \left[ 18(b^2 r^2 - (34 b^2 - 35) f) - \frac{(25 b^2 - 27)(43 b^2 - 45) \mu^2}{b^2 - 1} \right], \\ A_2 = \frac{2 b^2 r \kappa^4}{225(b^2 - 1)} \left[ 18(3 b^2 r^2 - 2(49 b^2 - 50) f) - \frac{(2925 - 5580 b^2 + 2659 b^4) \mu^2}{b^2 - 1} \right], \\ A_4 = b^2 r^2 \kappa^4 \left[ \mu^2 - f - \frac{b^2(r^2 - f)}{25(b^2 - 1)} \right]. \end{cases}$$

The system  $\{A_0 = 0, A_1 = 0, A_2 = 0, A_4 = 0\}$  is an algebraic system of four unknown  $r, \mu, f$  and  $b$ . Solving this system by Mathematica Program, we have  $r = \pm\sqrt{6} f$ ,  $\mu = 0$  and  $b = \pm\frac{\sqrt{5}}{2}$  which is contradiction with  $\mu \neq 0$ .

**(I-B):**  $\mu = \frac{r'}{\kappa}$ . From this equation and equation of  $\mu'$  above, the condition  $\mu' - \frac{d\mu}{du} = 0$ , gives

$$(18) \quad r'' = \frac{3 f \kappa^3 + 4 \kappa r'^2 + r r' \kappa'}{r \kappa}.$$

From  $A_1 = 0$ , it implies

$$(19) \quad f' = \frac{18 f \kappa^2 r' + 16 r'^3}{r \kappa^2}.$$

The condition  $f'' - \frac{df'}{du} = 0$  gives

$$(20) \quad r'^2 (f \kappa^2 - r'^2) \left[ (17 f - 25 r^2) \kappa^2 - 8 r'^2 \right] = 0.$$

If  $r' = 0$ , then  $\mu = 0$  contradiction. Therefore the above condition reduce two solutions:

**(I-B1):**  $f = \frac{1}{17} \left[ 25 r^2 + \frac{8 r'^2}{\kappa^2} \right]$ . Again the condition  $f' - \frac{df}{du} = 0$  leads  $r'^2 (\kappa^2 r^2 + r'^2) = 0$  which is contradiction.

**(I-B2):**  $f = -\frac{r'^2}{\kappa^2}$ . Again the condition  $f' - \frac{df}{du} = 0$  gives

$$(21) \quad r r' \kappa' + \kappa (r'^2 - r r'') = 0.$$

The general solution of this equation is

$$(22) \quad r(s) = r_0 \exp \left[ r_1 \int \kappa(s) ds \right], \quad r_0, r_1 \in R.$$

Now, all coefficients  $A_i$  and  $A_j$  are equal zero. In this case we have:

$$(23) \quad \begin{cases} \lambda(u) = m, & m = \sqrt{\frac{1+r_1^2}{b^2-1}}, & \tau(u) = m \kappa(u), \\ \alpha(u) = -\kappa(u) r(s), & \beta(u) = r_1 \kappa(u) r(u), & \gamma(u) = -\left(\frac{1+r_1^2}{m}\right) \kappa(u) r(u). \end{cases}$$

Therefore, the parametrization of the tangent of the curve  $\mathbf{C}(u)$  is given by:

$$\mathbf{C}'(u) = -r'(u) \left[ \frac{(1+r_1^2) \mathbf{b} + m \mathbf{t}}{m r_1} \right] + r_1 \kappa(u) r(u) \mathbf{n}.$$

Since  $\lambda(u) = m$ , then  $\Gamma(u)$  is a general helix, we can prove that:

$$-\frac{d}{du} \left[ \frac{(1+r_1^2) \mathbf{b} + m \mathbf{t}}{m r_1} \right] = r_1 \kappa(u) \mathbf{n}.$$

Hence, there exists  $\mathbf{C}_0 \in \mathbf{R}^3$  such that

$$\mathbf{C}(u) = \mathbf{C}_0 - r(u) \left[ \frac{(1+r_1^2) \mathbf{b} + m \mathbf{t}}{m r_1} \right].$$

The parametrization of this surface is given by

$$(24) \quad X(u, v) = \mathbf{C}_0 - r(u) \left[ \frac{(1+r_1^2) \mathbf{b} + m \mathbf{t}}{m r_1} - \cos[v] \mathbf{n} - b \sin[v] \mathbf{b} \right],$$

where  $b = \frac{\sqrt{1+m^2+r_1^2}}{m}$ ,  $r(u) = r_0 \exp \left[ r_1 \int \kappa(u) du \right]$  such that the curvature of the base curve  $\kappa(u)$  is arbitrary function of  $u$  and  $r_0, r_1$  and

the ratio  $\frac{\tau(u)}{\kappa(u)} = m$  are arbitrary constants.

**(II):**  $\mu' = \frac{(3f + 4\mu^2)r'}{\mu r}$ . If, we compute the  $\mu''$  and compare it with (12) we can obtain the following condition:

$$(25) \quad r'' = \frac{\kappa^2 \mu^2}{r} + \frac{3r'^2}{r} + \frac{3f^2 r'^2}{r \mu^2} + \frac{r' \kappa'}{\kappa}.$$

If we substitute  $r''$  from the equation  $A_1 = 0$  and solve it we get:

$$(26) \quad f' = \frac{2f \kappa^2 \mu^2 r' + 2(17f + 16\mu^2)r'^3}{2r(\kappa^2 \mu^2 + r'^2)}.$$

Again, by computing  $f''$  from (26) and equating with  $f''$  from (13), yields the following condition:

$$r'^2 (f + \mu^2) \left[ \kappa^2 \mu^2 (f - r^2) (3\kappa^2 \mu^2 + 7r'^2) + 8(3f - 5r^2 - 2\mu^2)r'^4 \right] = 0.$$

The above condition implies three cases:

**(II-A):**  $f = \frac{8(2\mu^2 + 5r^2)r'^4 + \kappa^2 \mu^2 r^2 (3\kappa^2 \mu^2 + 7r'^2)}{3\kappa^4 \mu^4 + 7\kappa^2 \mu^2 r'^2 + 24r'^4}$ . In this case we have

$$(27) \quad A_1 = -\frac{32b^2 r \kappa r'^3 (r^2 + \mu^2)}{\mu^3 (3\kappa^4 \mu^4 + 7\kappa^2 \mu^2 r'^2 + 24r'^4)^2} \left[ 3\kappa^8 \mu^8 + 26\kappa^6 \mu^6 r'^2 + 217\kappa^4 \mu^4 r'^4 + 314\kappa^2 \mu^2 r'^6 + 840r'^8 \right].$$

It is easy to see that  $A_1 = 0$  if  $r = r_0$ , where  $r_0$  is an arbitrary constant.

The condition  $r'' - \frac{d^2 r}{du^2}$  gives  $\kappa^2 \mu^2 = 0$  contradiction.

**(II-B):** When  $f = -\mu^2$ , then  $\mu' = \frac{\mu r'}{r}$  which means  $\mu(u) = \mu_1 r(u)$ , where  $\mu_1$  is an arbitrary constant. From (25) we get  $r'' = \mu_1^2 r \kappa^2 + \frac{\kappa' r'}{r}$ . By solving this equation, one can obtain  $r(s) = R_0 \cosh \left[ R_1 + \mu_0 \int \kappa(s) ds \right]$ . Now, all coefficients  $A_i$  and  $A_j$  are zero. Therefore, we

have:

$$(28) \quad \begin{cases} \lambda(s) = m, & m = \sqrt{\frac{1 + \mu_1^2}{b^2 - 1}}, & \tau(s) = m \kappa(u), \\ \alpha(u) = -\frac{r'(u)}{\mu_1}, & \beta(u) = \mu_1 \kappa(u) r(u), & \gamma(u) = -\frac{(1 + \mu_1^2) r'(u)}{m \mu_1}. \end{cases}$$

The parametrization of the tangent of the curve  $\mathbf{C}(u)$  can be written in the following form:

$$\mathbf{C}'(u) = -r'(u) \left[ \frac{(1 + \mu_1^2) \mathbf{b} + m \mathbf{t}}{m r_1} \right] + \mu_1 \kappa(u) r(u) \mathbf{n}.$$

Because  $\lambda(u) = m$ , then  $\Gamma(u)$  is a general helix and then:

$$-\frac{d}{du} \left[ \frac{(1 + \mu_1^2) \mathbf{b} + m \mathbf{t}}{m \mu_1} \right] = \mu_1 \kappa(u) \mathbf{n}.$$

So that, there exists  $\mathbf{C}_0 \in \mathbf{R}^3$  such that

$$\mathbf{C}(u) = \mathbf{C}_0 - r(u) \left[ \frac{(1 + \mu_1^2) \mathbf{b} + m \mathbf{t}}{m \mu_1} \right].$$

Now, the parametrization of this surface is given by

$$(29) \quad X(u, v) = \mathbf{C}_0 - r(u) \left[ \frac{(1 + \mu_1^2) \mathbf{b} + m \mathbf{t}}{m \mu_1} - \cos[v] \mathbf{n} - b \sin[v] \mathbf{b} \right],$$

where  $b = \frac{\sqrt{1 + m^2 + \mu_1^2}}{m}$ ,  $r(u) = R_0 \cosh \left[ R_1 + \mu_0 \int \kappa(u) du \right]$  such that the curvature of the base curve  $\kappa(s)$  is arbitrary function of  $u$  and  $R_0$ ,  $R_1$ ,  $\mu_1$  and the ratio  $\frac{\tau(u)}{\kappa(u)} = m$  are arbitrary constants.

**3.1.1.2**  $f = f_0 r^2$ , where  $f_0$  is an arbitrary constant. In this case, the solution of the equation  $A_4 = 0$  is  $\mu(u) = \mu_2 r(u)$ , where  $\mu_2 = \sqrt{-f_0}$  is an arbitrary constant. Here, all coefficients  $A_i$  and  $A_j$  are zero.

Therefore, we can obtain the following:

$$(30) \quad \begin{cases} \lambda(s) = m, & m = \sqrt{\frac{1 + \mu_2^2}{b^2 - 1}}, & \tau(s) = m \kappa(u), \\ \alpha(u) = -\frac{r'(u)}{\mu_2}, & \beta(u) = \mu_2 \kappa(u) r(u), & \gamma(u) = -\frac{(1 + \mu_2^2) r'(u)}{m \mu_2}. \end{cases}$$

Hence, the parametrization of the tangent of the curve  $\mathbf{C}(u)$  is given by:

$$\mathbf{C}'(u) = -r'(u) \left[ \frac{(1 + \mu_2^2) \mathbf{b} + m \mathbf{t}}{m \mu_2} \right] + \mu_2 \kappa(u) r(u) \mathbf{n}.$$

Since  $\lambda(u) = m$ , then  $\Gamma(u)$  is a general helix, we can prove that:

$$-\frac{d}{du} \left[ \frac{(1 + \mu_2^2) \mathbf{b} + m \mathbf{t}}{m \mu_2} \right] = \mu_2 \kappa(u) \mathbf{n}.$$

Therefore, there exists  $\mathbf{C}_0 \in \mathbf{R}^3$  such that

$$\mathbf{C}(u) = \mathbf{C}_0 - r(u) \left[ \frac{(1 + \mu_2^2) \mathbf{b} + m \mathbf{t}}{m \mu_2} \right].$$

Now, the parametrization of this surface is given by

$$(31) \quad X(u, v) = \mathbf{C}_0 - r(u) \left[ \frac{(1 + \mu_2^2) \mathbf{b} + m \mathbf{t}}{m \mu_2} - \cos[v] \mathbf{n} - b \sin[v] \mathbf{b} \right],$$

where  $b = \frac{\sqrt{1 + m^2 + \mu_2^2}}{m}$ , while  $r(u)$  is an arbitrary function of  $u$ . Also,  $\mu_1$  and the ratio  $\frac{\tau(u)}{\kappa(u)} = m$  are an arbitrary constants while the curvature and  $\kappa(s)$  of the base curve is arbitrary function of  $u$ .

**3.1.2.** When  $f = 0$ , then  $\lambda = \pm \frac{1}{\sqrt{b^2 - 1}}$ . For computation,  $A_4 = \frac{1}{2} b^2 r \kappa^4 \mu$  leads contradiction with  $\mu \neq 0$ .

**3.2.**  $\mu = 0$ . Then  $B_4 = \frac{1}{2} b (b^2 - 1) \kappa^3 r^3 \lambda'$  is vanished when  $\lambda(u) = \lambda_0$ , where  $\lambda_0$  is arbitrary constant. Then  $A_4 = 3b^2 [\lambda_0^4 (b^2 - 1) - 1] r^2 \kappa^3 \alpha$ , that is  $\alpha = 0$  or  $\lambda = \frac{1}{\sqrt{b^2 - 1}}$ . If  $\alpha = 0$ , then  $(\alpha, \beta, \gamma) = (0, 0, 0)$  contradiction. Now,  $\lambda = \frac{1}{\sqrt{b^2 - 1}}$  and the equation  $A_4 = 0$ , lead  $\gamma(u) = \sqrt{b^2 - 1} \alpha(u)$ . Therefore  $A_2 = \frac{2b^3 r^2 \kappa^3 r'}{\sqrt{b^2 - 1}} = 0$ , gives  $r(u) = r_0$ , where  $r_0$  is arbitrary constant. Now all coefficients  $A_i, B_i$  are trivially zero and the parametrization of the tangent of the curve  $\mathbf{C}(u)$  is given by

$$\mathbf{C}'(u) = \frac{\alpha(s)}{m} (m \mathbf{t} + \mathbf{b}),$$

where  $m = \frac{1}{\sqrt{b^2 - 1}}$ . Since  $\Gamma(u)$  is a general helix, then  $\frac{d(m \mathbf{t} + \mathbf{b})}{du} = 0$  and hence,  $m \mathbf{t} + \mathbf{b}$  is a fixed vector. Therefore, there exists  $\mathbf{C}_0 \in \mathbf{R}^3$  such that

$$\mathbf{C}(u) = \mathbf{C}_0 + (m \mathbf{t} + \mathbf{b}) \Omega(u),$$

where  $\Omega(u) = \frac{1}{m} \int \alpha(u) du$  is arbitrary function of  $u$ . The parameterization of this surface is given by

$$(32) \quad X(u, v) = \mathbf{C}_0 + (m \mathbf{t} + \mathbf{b}) \Omega(u) + c \left( m \cos[v] \mathbf{n} + \sqrt{1 + m^2} \sin[v] \mathbf{b} \right),$$

where  $c = \frac{r_0}{m}$  and  $m$  are arbitrary constants.

**REMARK 3.1.** The developable surface (24) foliated by an ellipse is a special surface of (31) when  $r(u) = r_0 \exp \left[ r_1 \int \kappa(u) du \right]$ .

**REMARK 3.2.** The developable surface (29) foliated by an ellipse is a special surface of (31) when  $r(u) = R_0 \cosh \left[ R_1 + \mu_0 \int \kappa(u) du \right]$ .

**LEMMA 3.3.** *The surface (1) foliated by an ellipse is developable if and only if it takes the general forms (31) or (32).*

**LEMMA 3.4.** *If the base curve  $\Gamma(u)$  is not a general helix, then the surface (1) foliated by an ellipse is non-developable.*

Now, we will introduce the position vector of two obtained developable surfaces foliated by an ellipse in Euclidean 3-space. Firstly, we write the following Theorem

**THEOREM 3.5.** *The position vector  $\mathbf{C}$  of a general helix whose tangent vector makes a constant angle with a fixed straight line in the space, is expressed in the natural representation form as follows:*

(33)

$$\mathbf{C}(u) = \sqrt{1-n^2} \int \left( \cos \left[ \sqrt{1+m^2} \int \kappa(u) du \right], \sin \left[ \sqrt{1+m^2} \int \kappa(u) du \right], m \right) du,$$

where  $m = \frac{n}{\sqrt{1-n^2}}$ ,  $n = \cos[\phi]$  and  $\phi$  is the angle between the fixed straight line  $\mathbf{e}_3$  (axis of a general helix) and the tangent vector of the curve  $\mathbf{C}$ .

**(1):** The position vector  $\mathbf{X}(u, v) = (x_1, x_2, x_3)$  of the surface (31) takes the following form:

$$(34) \quad \begin{cases} x_1 = r(u) \left[ \left( Q - \sqrt{1+Q^2} \sin[v] \right) \cos[\Phi] - \cos[v] \sin[\Phi] \right], \\ x_2 = r(u) \left[ \left( Q - \sqrt{1+Q^2} \sin[v] \right) \sin[\Phi] + \cos[v] \cos[\Phi] \right], \\ x_3 = \frac{\sqrt{1+Q^2} r(u)}{mQ} \left[ Q \sin[v] - \sqrt{1+Q^2} \right], \end{cases}$$

where  $Q = \frac{\mu_2}{\sqrt{1+m^2}}$ , and  $\Phi = \sqrt{1+m^2} \int \kappa(u) du$ . Note that in this surface,  $r(s)$  and  $\kappa(s)$  are arbitrary function of the variable  $u$  while  $Q$  and  $m$  are arbitrary constants. It is worth noting that for the above surface we have

$$x_1^2 + x_2^2 = \left( \frac{m^2 Q^2}{1+Q^2} \right) x_3^2.$$

So that, we can write the following Lemma:

**LEMMA 3.6.** *The developable surface (31) foliated by an ellipse is represented a generalized cone in Euclidean 3-space.*

(2): The position vector of the surface (32) reduces to the following form:

$$(35) \quad \mathbf{X}(u, v) = (X_1, X_2, X_3) = \left( -M_0 \sin [\Theta], M_0 \cos [\Theta], \Omega \right)$$

where  $M_0 = m c$ ,  $\Theta = v - \sqrt{1 + m^2} \int \kappa(u) du$  and  $\Psi = \Omega(u) + c \sin \left[ \Theta + \sqrt{1 + m^2} \int \kappa(u) du \right]$ . Note that,  $\Omega(u)$  and  $\kappa(u)$  are arbitrary functions of the variable  $u$  while  $m$  and  $c$  are an arbitrary constant. It is easy to see that, for the above surface the following condition is satisfied:

$$X_1^2 + X_2^2 = M^2.$$

The above condition gives the following Lemma:

LEMMA 3.7. *The developable surface (32) foliated by an ellipse is represents a generalized cylinder in Euclidean 3-space.*

From the above, the main Theorem (1.1) is proved.

#### 4. Proof Theorem 1.2

In this section, we assume that the the surface (1) has a non-zero constant Gaussian curvature  $K$ . In this case equation (9) can be written in the form

$$\sum_{i=0}^8 E_i(u) \cos[i v] + F_i(u) \sin[i v] = 0$$

One begin to compute the coefficients  $E_i$  and  $F_i$ . The first coefficient

$$E_8 = \frac{1}{16} (b^2 - 1) K \kappa^4 r^6 [1 - (b^2 - 1) \lambda^2] = 0,$$

leads to

$$\lambda = \pm \frac{1}{\sqrt{b^2 - 1}}.$$

The coefficient  $F_6$  becomes

$$F_6 = 2 b (b^2 - 1) K \kappa^3 r^4 \mu [\gamma - \alpha \sqrt{b^2 - 1}].$$

For vanishes the above coefficient, we have two possibilities:

**4.1. When  $\mu(u) = 0$ , the coefficient  $E_6$  becomes:**

$$E_6 = (1 - b^2) K \kappa^2 r^4 \left( \alpha \sqrt{b^2 - 1} - \gamma \right)^2.$$

The vanishing of  $E_6$  gives  $\gamma = \alpha \sqrt{b^2 - 1}$ . Now,  $F_4 = 4 b^3 \sqrt{b^2 - 1} K \kappa^3 r^5 r' = 0$ , which means  $r = r_0$  where  $r_0$  is arbitrary constant. Finally, the coefficient  $E_4 = -b^4 K \kappa^4 r^6$  leads to contradiction.

**4.2.**  $\gamma = \alpha \sqrt{b^2 - 1}$ . Now,  $F_6 = b^2 (b^2 - 1) K \kappa^4 r^4 \mu^2 = 0$  gives  $\mu = 0$  and this case has been discussed previously. Therefore the proof of the Theorem 1.2 is completed.

### 5. Proof Theorem 1.3

Let  $M$  be a surface (4) in  $\mathbf{E}^3$  with constant Gauss curvature  $K = K_0$  and foliated by a pieces of ellipses in parallel planes. If we put  $K_0 = \frac{P}{W}$  in the computations of the Gauss curvature  $K$ , yields

(36)

$$W = r \left[ (1 + g'^2) (1 - \cos[2v]) + b^2 (1 + f'^2) (1 + \cos[2v]) + 2b \left( b r'^2 + 2 r' (g' \sin[v] + b f' \cos[v]) + 2 f' g' \sin[2v] \right) \right]^2,$$

$$(37) \quad P = -4b \left( b r'' + g'' \sin[v] + b f'' \cos[v] \right).$$

**5.1.**  $K_0 \neq 0$ . The identity (9) writes as:

$$\sum_{i=0}^4 (A_i \cos[i v] + B_i \sin[i v]) = 0.$$

A computations yields

(38)

$$\begin{cases} A_4 = -\frac{K_0 r}{2} \left( b^4 f'^4 - 6 b^2 f'^2 g'^2 + g'^4 + (b^2 - 1) \left[ b^2 (1 + 2 f'^2) - (1 + 2 g'^2) \right] \right), \\ B_4 = 2 b K_0 r f' g' \left[ 1 + g'^2 - b^2 (1 + f'^2) \right]. \end{cases}$$

According to the condition  $B_4 = 0$ , we have distinguish three possibilities as the following:

**5.1.1.**  $f' = 0$  and  $f(u) = f_0$ , where  $f_0$  is arbitrary constant. In this case  $A_4 = \frac{K_0 r}{2} (b^2 - 1 - g'^2)^2 = 0$  which implies  $g(u) = \pm \sqrt{b^2 - 1} u + g_0$ , where  $g_0$  is an arbitrary constant and  $|b| > 1$ . Therefore the coefficient  $A_2$  implies  $8(b^2 - 1) K_0 r r'^2 = 0$  and thus  $r(u) = r_0$ , where  $r_0$  is arbitrary constant. In this case,  $A_0 = 0$ , yields  $4b^4 K_0 r_0 = 0$  contradiction.

**5.1.2.**  $g' = 0$  and  $g(u) = g_0$ , where  $g_0$  is arbitrary constant. In this case  $A_4 = \frac{K_0 r}{2} [1 - b^2(1 + f'^2)]^2 = 0$  which implies  $f(u) = \pm \frac{\sqrt{1 - b^2} u}{b} + f_0$ , where  $f_0$  is an arbitrary constant and  $|b| < 1$ . Therefore the coefficient  $A_2$  implies  $8(b^2 - 1) K_0 r r'^2 = 0$  and thus  $r(u) = r_0$ , where  $r_0$  is arbitrary constant. With these conditions  $A_0 = 4 K_0 r_0 = 0$  contradiction.

**5.1.3.**  $f' \neq 0$  and  $g' \neq 0$  at some  $u$ -interval. Since  $B_4 = 0$ , then  $g' = \pm \sqrt{b^2(1 + f'^2) - 1}$ . Taking in account  $A_4 = 0$ , it follows

$$2b^2 K_0 r f'^2 [b^2(1 + f'^2) - 1] = 0,$$

which leads to  $f(u) = \left(\sqrt{\frac{1}{b^2} - 1}\right) u + f_0$ . Then, the coefficient  $A_2 = 8b^2 K_0 (b^2 - 1) r r'$  must be vanishes. Therefore  $r(u) = r_0$  and  $A_0 = -4 K_0 r_0 = 0$  contradiction.

**5.2.**  $K_0 = 0$ . Identity (9) leads to  $P = 0$ . In view of the above expression of  $P$ , it follows that  $r'' = f'' = g'' = 0$ . As consequence, there are constants  $r_0, r_1, f_0, f_1, g_0$  and  $g_1$  such that

$$(39) \quad \begin{cases} r(u) &= r_1 u + r_0, \\ f(u) &= f_1 u + f_0, \\ g(u) &= g_1 u + g_0, \end{cases}$$

that is, the functions  $f, g$  and  $r$  are linear on  $u$  and so, the surface is a generalized cone. Therefore the proof of Theorem 1.3 is completed.

### References

- [1] Ali A.T., *Position vectors of general helices in Euclidean 3-space*, Bull. Math. Anal. Appl. **3** (2) (2010), 198–205.
- [2] Ali A.T., *Position vectors of slant helices in Euclidean 3-space*, J. Egyptian Math. Soc. **20** (1) (2012), 1–6.
- [3] Delaunay C., *Sur la surface de révolution dont la courbure moyenne est constante*, J. Math. Pure Appl. **6** (1841), 309–320.
- [4] Enneper A., *Ueber die cyclischen Flächen*, Nach. K. önlgl. Ges. d. Wissensch. Göttingen, Math. Phys. **K1** (1866), 243–249.
- [5] Enneper A., *Die cyclischen Flächen*, Z. Math. Phys. **14** (1869), 393–421.
- [6] López R. *Cyclic surfaces of constant Gauss curvature*, Houston J. Math. **27** (4) (2001), 799–805.
- [7] López R. *On linear Weingarten surfaces*, Int. J. Math. **19** (2008), 439–448.
- [8] López R. *Special Weingarten surfaces foliated by circles*, Monatsh. Math. **154** (2008), 289–302.
- [9] Nitsche J. C. C., *Cyclic surfaces of constant mean curvature*, Nachr. Akad. Wiss. Göttingen Math. Phys. **II 1** (1989), 1–5.
- [10] Riemann, B. *Über die Flächen vom kleinsten Inhalt bei gegebener Begrenzung*, Abh. Königl. Ges. d. Wissensch. Göttingen, Mathema. C1, **13** (1868), 329–333.

#### Ahmed T. Ali

Department of Mathematics, Faculty of Science  
 King Abdul Aziz University  
 Jeddah 21589, Saudi Arabia.  
*E-mail*: atali71@yahoo.com

#### Fathi M. Hamdoon

\*<sup>1</sup>Mathematics Department, Faculty of Science  
 Al-Azhar University  
 Assiut branch 71524 Assiut, Egypt.  
 \*<sup>2</sup>Mathematics Department, Faculty of Science  
 Al-Azhar University  
 Nasr City 11884 Cairo, Egypt.  
*E-mail*: f\_hamdoon@yahoo.com