

## ***L*-FUZZY BI-CLOSURE SYSTEMS AND *L*-FUZZY BI-CLOSURE OPERATORS**

JUNG MI KO AND YONG CHAN KIM\*

ABSTRACT. In this paper, we introduced the notions of right and left closure systems on generalized residuated lattices. In particular, we study the relations between right (left) closure (interior) operators and right (left) closure (interior) systems. We give their examples.

### **1. Introduction**

The notion of closure systems and closure operators facilitated to study topological structures, logic and lattices. Gerla [5-7] introduced closure systems and closure operators in the unit interval  $[0,1]$ .

Ward et al.[16] introduced a complete residuated lattice which is an algebraic structure for many valued logic. It is an important mathematical tool for algebraic structure. By using the concepts of lower and upper approximation operators, information systems and decision rules are investigated in complete residuated lattices [1-4, 11-14]. Recently, Bělohlávek [1-4] investigate the properties of fuzzy relations and fuzzy closure systems on a residuated lattice which supports part of foundation of theoretic computer science. As an Bělohlávek's extension, Fang and Yue [8] introduced strong fuzzy closure systems and strong fuzzy

---

Received October 31, 2018. Revised May 20, 2019. Accepted June 12, 2019.

2010 Mathematics Subject Classification: 03E72, 03G10, 06A15, 54F05.

Key words and phrases: Right (resp. left, bi-) closure systems, generalized residuated lattices, right (resp. left, bi-) closure (interior) operators.

\* Corresponding author.

This work was supported by the Research Institute of Natural Science of Gangneung-Wonju National University.

© The Kangwon-Kyungki Mathematical Society, 2019.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

closure operators. On the other hand, Georgescu and Popescue [9,10] introduced non-commutative fuzzy Galois connection in a generalized residuated lattice which is induced by two implications.

In this paper, we introduced the notions of right and left closure (resp. interior) systems in a sense as the right and left least upper (resp. greatest lower) bound on a generalized residuated lattice. In particular, we investigated the relations between right (left) closure (interior) operators and and right (left) closure (interior) systems. We give their examples.

## 2. Preliminaries

**DEFINITION 2.1.** [9,10,17] A structure  $(L, \vee, \wedge, \odot, \rightarrow, \Rightarrow, \perp, \top)$  is called a *generalized residuated lattice* if it satisfies the following conditions:

(GR1)  $(L, \vee, \wedge, \top, \perp)$  is a bounded lattice where  $\top$  is the upper bound and  $\perp$  denotes the universal lower bound;

(GR2)  $(L, \odot, \top)$  is a monoid;

(GR3) it satisfies a residuation, i.e.

$$(a \odot b) \leq c \text{ iff } a \leq (b \rightarrow c) \text{ iff } b \leq (a \Rightarrow c).$$

**REMARK 2.2.** [9,10,15,17]

(1) A generalized residuated lattice is a residuated lattice  $(\rightarrow \Rightarrow)$  iff  $\odot$  is commutative.

(2) A left-continuous t-norm  $([0, 1], \leq, \odot)$  defined by  $a \rightarrow b = \bigvee \{c \mid a \odot c \leq b\}$  is a residuated lattice

(3) Let  $(L, \leq, \odot)$  be a quantale. For each  $x, y \in L$ , we define

$$x \rightarrow y = \bigvee \{z \in L \mid z \odot x \leq y\},$$

$$x \Rightarrow y = \bigvee \{z \in L \mid x \odot z \leq y\}.$$

Then it satisfies Galois correspondence, that is,

$(x \odot y) \leq z$  iff  $x \leq (y \rightarrow z)$  iff  $y \leq (x \Rightarrow z)$ . Hence  $(L, \vee, \wedge, \odot, \rightarrow, \Rightarrow, \perp, \top)$  is a generalized residuated lattice.

(4) A pseudo MV-algebra is a generalized residuated lattice with the law of double negation.

In this paper, we assume  $(L, \wedge, \vee, \odot, \rightarrow, \Rightarrow, \top, \perp)$  is a complete generalized residuated lattice with the law of double negation defined as  $a = (a^*)^0 = (a^0)^*$  where  $a^0 = a \rightarrow \perp$  and  $a^* = a \Rightarrow \perp$ .

LEMMA 2.3 [9,10,17] *For each  $x, y, z, x_i, y_i \in L$ , we have the following properties.*

- (1) *If  $y \leq z$ ,  $(x \odot y) \leq (x \odot z)$ ,  $(x \rightarrow y) \leq (x \rightarrow z)$  and  $(z \rightarrow x) \leq (y \rightarrow x)$  for  $\rightarrow \in \{\rightarrow, \Rightarrow\}$ .*
- (2)  *$x \rightarrow (\bigwedge_{i \in \Gamma} y_i) = \bigwedge_{i \in \Gamma} (x \rightarrow y_i)$  and  $(\bigvee_{i \in \Gamma} x_i) \rightarrow y = \bigwedge_{i \in \Gamma} (x_i \rightarrow y)$  for  $\rightarrow \in \{\rightarrow, \Rightarrow\}$ .*
- (3)  *$(x \odot y) \rightarrow z = x \rightarrow (y \rightarrow z)$  and  $(x \odot y) \Rightarrow z = y \Rightarrow (x \Rightarrow z)$ .*
- (4)  *$x \rightarrow (y \Rightarrow z) = y \Rightarrow (x \rightarrow z)$  and  $x \Rightarrow (y \rightarrow z) = y \rightarrow (x \Rightarrow z)$ .*
- (5)  *$x \odot (x \Rightarrow y) \leq y$  and  $(x \rightarrow y) \odot x \leq y$ .*
- (6)  *$(x \Rightarrow y) \odot (y \Rightarrow z) \leq x \Rightarrow z$  and  $(y \rightarrow z) \odot (x \rightarrow y) \leq x \rightarrow z$ .*
- (7)  *$(x \Rightarrow z) \leq (y \odot x) \Rightarrow (y \odot z)$  and  $(x \rightarrow z) \leq (x \odot y) \rightarrow (z \odot y)$ .*
- (8)  *$x \rightarrow y \leq (y \rightarrow z) \Rightarrow (x \rightarrow z)$  and  $(x \Rightarrow y) \leq (y \Rightarrow z) \rightarrow (x \Rightarrow z)$ .*
- (9)  *$y \rightarrow z \leq (x \rightarrow y) \rightarrow (x \rightarrow z)$  and  $(y \Rightarrow z) \leq (x \Rightarrow y) \Rightarrow (x \Rightarrow z)$ .*
- (10)  *$x \rightarrow y = \top$  iff  $x \leq y$ .*
- (11)  *$x \rightarrow y = y^0 \Rightarrow x^0$  and  $x \Rightarrow y = y^* \rightarrow x^*$ .*
- (12)  *$(x \rightarrow y)^* = x \odot y^*$  and  $(x \Rightarrow y)^0 = y^0 \odot x$ .*
- (13)  *$\bigwedge_{i \in \Gamma} x_i^* = (\bigvee_{i \in \Gamma} x_i)^*$  and  $\bigvee_{i \in \Gamma} x_i^* = (\bigwedge_{i \in \Gamma} x_i)^*$ .*
- (14)  *$\bigwedge_{i \in \Gamma} x_i^0 = (\bigvee_{i \in \Gamma} x_i)^0$  and  $\bigvee_{i \in \Gamma} x_i^0 = (\bigwedge_{i \in \Gamma} x_i)^0$ .*
- (15)  *$\bigwedge_{i \in \Gamma} x_i \rightarrow (\bigwedge_{i \in \Gamma} y_i) \geq \bigwedge_{i \in \Gamma} (x_i \rightarrow y_i)$  and  $\bigvee_{i \in \Gamma} x_i \rightarrow (\bigvee_{i \in \Gamma} y_i) \geq \bigwedge_{i \in \Gamma} (x_i \rightarrow y_i)$  for  $\rightarrow \in \{\rightarrow, \Rightarrow\}$ .*

DEFINITION 2.4. Let  $X$  be a set. A function  $e_X^r : X \times X \rightarrow L$  is called a *right partial order* if it satisfies the following conditions :

- (O1)  $e_X^r(x, x) = \top$  for all  $x \in X$ ,
- (O2) If  $e_X^r(x, y) = e_X^r(y, x) = \top$ , then  $x = y$ ,
- (R)  $e_X^r(x, y) \odot e_X^r(y, z) \leq e_X^r(x, z)$ , for all  $x, y, z \in X$ .

A function  $e_X^l : X \times X \rightarrow L$  is called a *left partial order* if it satisfies (O1), (O2) and

- (L)  $e_X^l(y, z) \odot e_X^l(x, y) \leq e_X^l(x, z)$ , for all  $x, y, z \in X$ .

The triple  $(X, e_X^r, e_X^l)$  is a bi-partial ordered set.

EXAMPLE 2.5.

- (1) We define a function  $e_L^r, e_L^l : L \times L \rightarrow L$  as

$$e_L^r(x, y) = (x \Rightarrow y), \quad e_L^l(x, y) = (x \rightarrow y).$$

By Lemma 2.3 (6),  $(L, e_L^r, e_L^l)$  is a bi-partial ordered set.

(2) We define a function  $e_{L^X}^r, e_{L^X}^l : L^X \times L^X \rightarrow L$  as

$$e_{L^X}^r(A, B) = \bigwedge_{x \in X} (A(x) \Rightarrow B(x)),$$

$$e_{L^X}^l(A, B) = \bigwedge_{x \in X} (A(x) \rightarrow B(x)).$$

By Lemma 2.3 (6),  $(L^X, e_{L^X}^r, e_{L^X}^l)$  is a bi-partial ordered set.

### 3. $L$ -fuzzy bi-closure systems and $L$ -fuzzy bi-closure operators

DEFINITION 3.1. A map  $S^r : L^X \rightarrow L$  is called an  $L$ -fuzzy right closure system if

(S1)  $S^r(\top_X) = \top$ ,

(S2)  $S^r(\bigwedge_{i \in \Gamma} A_i) \geq \bigwedge_{i \in \Gamma} S_X^r(A_i)$ , for all  $A_i \in L^X$ ,

(RS)  $S^r(\alpha \Rightarrow A) \geq S_X^r(A)$ , for all  $A \in L^X$  and  $\alpha \in L$ .

A map  $S^l : L^X \rightarrow L$  is called an  $L$ -fuzzy left closure system if it satisfies (S1), (S2) and

(LS)  $S^l(\alpha \rightarrow A) \geq S_X^l(A)$ , for all  $A \in L^X$  and  $\alpha \in L$ .

The triple  $(X, S^r, S^l)$  is called an  $L$ -fuzzy bi-closure system. A map  $f : (X, S_X^r, S_X^l) \rightarrow (Y, S_Y^r, S_Y^l)$  is called bi-continuous if  $S_X^r(f^{\leftarrow}(B)) \geq S_Y^r(B)$  and  $S_X^l(f^{\leftarrow}(B)) \geq S_Y^l(B)$  for each  $B \in L^Y$ .

DEFINITION 3.2. Let  $(X, e_{L^X}^r, e_{L^X}^l)$  be a bi-partial ordered set. An operator  $C^r : L^X \rightarrow L^X$  is called an  $L$ -fuzzy right closure operator on  $X$  if it satisfies the following conditions:

(CR1)  $e_{L^X}^r(A, C^r(A)) = \top$ , for all  $A \in L^X$ .

(CR2)  $e_{L^X}^r(A, B) \leq e_{L^X}^r(C^r(A), C^r(B))$  for all  $A, B \in L^X$ .

(CR3)  $\alpha \odot e_{L^X}^r(C^r(A)) \leq C^r(\alpha \odot A)$  for all  $A \in L^X$  and  $\alpha \in L$ .

An operator  $C^l : L^X \rightarrow L^X$  is called an  $L$ -fuzzy left closure operator on  $X$  if it satisfies the conditions

(CL1)  $e_{L^X}^l(A, C^l(A)) = \top$  for all  $A \in L^X$ ,

(CL2)  $e_{L^X}^l(A, B) \leq e_{L^X}^l(C^l(A), C^l(B))$  for all  $A, B \in L^X$ .

(CR3)  $e_{L^X}^l(C^l(A)) \odot \alpha \leq C^l(A \odot \alpha)$  for all  $A \in L^X$  and  $\alpha \in L$ .

The triple  $(X, C^r, C^l)$  is called an  $L$ -fuzzy bi-closure space. A map  $f : (X, C_X^r, C_X^l) \rightarrow (Y, C_Y^r, C_Y^l)$  is called an  $L$ -fuzzy bi-closed map if

$C_{S^r}^r(f^{\leftarrow}(B)) \leq f^{\leftarrow}(C_Y^r(B))$  and  $C_X^l(f^{\leftarrow}(B)) \leq f^{\leftarrow}(C_Y^l(B))$  for each  $B \in L^Y$ .

**THEOREM 3.3.** *Let  $(L^X, e_{L^X}^r, e_{L^X}^l)$  be a bi-partial ordered set and  $(X, S^r, S^l)$  be an L-fuzzy bi-closure system. Define two maps  $C_{S^r}^r, C_{S^l}^l : L^X \rightarrow L^X$  as follows:*

$$C_{S^r}^r(A) = \bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot S^r(B) \rightarrow B),$$

$$C_{S^l}^l(A) = \bigwedge_{B \in L^X} (S^l(B) \odot e_{L^X}^l(A, B) \Rightarrow B).$$

Then  $(X, C_{S^r}^r, C_{S^l}^l)$  is an L-fuzzy bi-closure space.

*Proof.* (CR1) For each  $A \in L^X$ , by Lemma 2.3(2,4),

$$\begin{aligned} & e_{L^X}^r(A, C_{S^r}^r(A)) \\ &= \bigwedge_{x \in X} (A(x) \Rightarrow (\bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot S^r(B) \rightarrow B))) \\ &= \bigwedge_{x \in X} \bigwedge_{B \in L^X} (A(x) \Rightarrow ((e_{L^X}^r(A, B) \odot S^r(B) \rightarrow B))) \\ &= \bigwedge_{x \in X} \bigwedge_{B \in L^X} ((e_{L^X}^r(A, B) \odot S^r(B) \rightarrow (A(x) \Rightarrow B))) \\ &= \bigwedge_{B \in L^X} ((e_{L^X}^r(A, B) \odot S^r(B) \rightarrow \bigwedge_{x \in X} (A(x) \Rightarrow B))) = \top. \end{aligned}$$

(CR2) We will show that  $e_{L^X}^r(A, B) \leq e_{L^X}^r(C_{S^r}^r(A), C_{S^r}^r(B))$ .

$$\begin{aligned} & C_{S^r}^r(A) \odot e_{L^X}^r(A, B) \odot e_{L^X}^r(B, D) \odot S^r(D) \\ & \leq \bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot S^r(B) \rightarrow B) \odot e_{L^X}^r(A, D) \odot S^r(D) \\ & \leq D \text{ (by Lemma 2.3(5)).} \end{aligned}$$

Then  $C_{S^r}^r(A) \odot e_{L^X}^r(A, B) \leq C_{S^r}^r(B)$ . Hence  $e_{L^X}^r(A, B) \leq e_{L^X}^r(C_{S^r}^r(A), C_{S^r}^r(B))$ .

(CR3) For each  $A, B \in L^X$ , since  $S^r(\alpha \Rightarrow B) \geq S^r(B)$ ,

$$\begin{aligned} C_{S^r}^r(A) &= \bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot S^r(B) \rightarrow B) \\ &= \bigwedge_{B \in L^X} (e_{L^X}^r(A, \alpha \Rightarrow B) \odot S^r(\alpha \Rightarrow B) \rightarrow (\alpha \Rightarrow B)) \\ &= \bigwedge_{B \in L^X} (e_{L^X}^r(\alpha \odot A, B) \odot S^r(\alpha \Rightarrow B) \rightarrow (\alpha \Rightarrow B)) \\ &= \alpha \Rightarrow \bigwedge_{B \in L^X} (e_{L^X}^r(\alpha \odot A, B) \odot S^r(\alpha \Rightarrow B) \rightarrow B) \\ &\leq \alpha \Rightarrow \bigwedge_{B \in L^X} (e_{L^X}^r(\alpha \odot A, B) \odot S^r(B) \rightarrow B) \\ &= \alpha \Rightarrow C_{S^r}^r(\alpha \odot A) \end{aligned}$$

Hence  $\alpha \odot C_{S^r}^r(A) \leq C_{S^r}^r(\alpha \odot A)$ . Thus  $C_{S^r}^r$  is an L-fuzzy right closure operator. Similarly,  $C_{S^l}^l$  is an L-fuzzy left closure operator. □

**THEOREM 3.4.** *Let  $(L^X, e_{L^X}^r, e_{L^X}^l)$  be a bi-partial ordered set and  $(X, C^r, C^l)$  be an  $L$ -fuzzy bi-closure operator. Define two maps  $S_{C^r}^r, S_{C^l}^l : L^X \rightarrow L$  as follows:*

$$S_{C^r}^r(A) = e_{L^X}^r(C^r(A), A),$$

$$S_{C^l}^l(A) = e_{L^X}^l(C^l(A), A).$$

(1)  $(X, S_{C^r}^r, S_{C^l}^l)$  is an  $L$ -fuzzy bi-closure system such that  $C^r \leq C_{S_{C^r}^r}^r$  and  $C^l \leq C_{S_{C^l}^l}^l$ .

(2) If  $(X, S^r, S^l)$  is an  $L$ -fuzzy bi-closure system, then  $S^r \leq S_{C^r}^r$  and  $S^l \leq S_{C^l}^l$ .

*Proof.* (1) (S1)  $S_{C^r}^r(\top_X) = e_{L^X}^r(C^r(\top_X), \top_X) = \top$ .

(S2) For all  $A_i \in L^X$ , by Lemma 2.3(15),

$$\begin{aligned} S_{C^r}^r(\bigwedge_{i \in \Gamma} A_i) &= e_{L^X}^r(C^r(\bigwedge_{i \in \Gamma} A_i), \bigwedge_{i \in \Gamma} A_i) \\ &\geq e_{L^X}^r(\bigwedge_{i \in \Gamma} C^r(A_i), \bigwedge_{i \in \Gamma} A_i) \\ &\geq \bigwedge_{i \in \Gamma} e_{L^X}^r(C^r(A_i), A_i) \\ &= \bigwedge_{i \in \Gamma} S_{C^r}^r(A_i) \end{aligned}$$

(RS) For all  $A \in L^X$  and  $\alpha \in L$ , by Lemma 2.3(3),

$$\begin{aligned} S_{C^r}^r(\alpha \Rightarrow A) &= e_{L^X}^r(C^r(\alpha \Rightarrow A), \alpha \Rightarrow A) \\ &= e_{L^X}^r(\alpha \odot C^r(\alpha \Rightarrow A), A) \\ &\geq e_{L^X}^r(C^r(\alpha \odot (\alpha \Rightarrow A)), A) \\ &\geq e_{L^X}^r(C^r(A), A) = S_{C^r}^r(A). \end{aligned}$$

Hence  $S_{C^r}^r$  is an  $L$ -fuzzy right closure system. Moreover, for all  $A \in L^X$ ,

$$\begin{aligned} C_{S_{C^r}^r}^r(A) &= \bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot S_{C^r}^r(B) \rightarrow B) \\ &= \bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot e_{L^X}^r(C^r(B), B) \rightarrow B) \\ &\geq \bigwedge_{B \in L^X} (e_{L^X}^r(C^r(A), C^r(B)) \odot e_{L^X}^r(C^r(B), B) \rightarrow B) \\ &\geq \bigwedge_{B \in L^X} (e_{L^X}^r(C^r(A), B) \rightarrow B) \\ &\geq C^r(A). \end{aligned}$$

Similarly,  $S_{C^l}^l$  is an  $L$ -fuzzy left closure system.

(2) Since  $(a \rightarrow b) \odot a \leq b$  iff  $a \leq (a \rightarrow b) \Rightarrow$ ,

$$\begin{aligned} S_{C_{S^r}^r}^r(A) &= e_{L^X}^r(C_{S^r}^r(A), A) \\ &= e_{L^X}^r(\bigwedge_{B \in L^X} (e_{L^X}^r(A, B) \odot S^r(B) \rightarrow B), A) \\ &\geq e_{L^X}^r((e_{L^X}^r(A, A) \odot S^r(A) \rightarrow A), A) \\ &\geq S^r(A). \end{aligned}$$

Similarly,  $S^l \leq S_{C^l_{S^l}}^l$ . □

**THEOREM 3.5.** (1) *If  $f : (X, C_X^r, C_X^l) \rightarrow (Y, C_Y^r, C_Y^l)$  is an  $L$ -fuzzy bi-closure map, then  $f : (X, S_{C_X^r}^r, S_{C_X^l}^l) \rightarrow (Y, S_{C_Y^r}^r, S_{C_Y^l}^l)$  is a bi-continuous map.*

(2) *If  $f : (X, S_X^r, S_X^l) \rightarrow (Y, S_Y^r, S_Y^l)$  is a bi-continuous map, then  $f : (X, C_{S_X^r}^r, C_{S_X^l}^l) \rightarrow (Y, C_{S_Y^r}^r, C_{S_Y^l}^l)$  is an  $L$ -fuzzy bi-closure map.*

*Proof.* (1) Since  $C_X^r(f^{\leftarrow}(B)) \leq f^{\leftarrow}(C_Y^r(B))$ ,

$$\begin{aligned} S_{C_X^r}^r(f^{\leftarrow}(B)) &= e_{L^X}^r(C^r(f^{\leftarrow}(B)), f^{\leftarrow}(B)) \\ &\geq e_{L^X}^r(f^{\leftarrow}(C_Y^r(B)), f^{\leftarrow}(B)) \\ &\geq \bigwedge_{x \in X} (C_Y^r(B)(f(x)) \Rightarrow B(f(x))) \\ &\geq \bigwedge_{y \in Y} (C_Y^r(B)(y) \Rightarrow B(y)) \\ &= S_{C_Y^r}^r(B). \end{aligned}$$

Similarly,  $S_{C_X^l}^l(f^{\leftarrow}(B)) \geq S_{C_Y^l}^l(B)$  for all  $B \in L^Y$ .

(2) Since  $S_X^r(f^{\leftarrow}(D)) \geq S_Y^r(D)$  for all  $D \in L^Y$ ,

$$\begin{aligned} f^{\leftarrow}(C_{S_Y^r}^r(B)) &= f^{\leftarrow}(\bigwedge_{D \in L^Y} (e_{L^Y}^r(B, D) \odot S_Y^r(D) \rightarrow D)) \\ &= \bigwedge_{D \in L^Y} (e_{L^Y}^r(B, D) \odot S_Y^r(D) \rightarrow f^{\leftarrow}(D)) \\ &\geq \bigwedge_{D \in L^Y} (e_{L^X}^r(f^{\leftarrow}(B), f^{\leftarrow}(D)) \odot S_X^r(f^{\leftarrow}(D)) \rightarrow f^{\leftarrow}(D)) \\ &\geq \bigwedge_{E \in L^X} (e_{L^X}^r(f^{\leftarrow}(B), E) \odot S_X^r(E) \rightarrow E) \\ &= C_{S_X^r}^r(f^{\leftarrow}(B)). \end{aligned}$$

Similarly,  $f^{\leftarrow}(C_{S_Y^l}^l(B)) \geq C_{S_X^l}^l(f^{\leftarrow}(B))$  for all  $B \in L^Y$ . □

**DEFINITION 3.6.** [1] Suppose that  $F : \mathcal{D} \rightarrow \mathcal{C}$ ,  $G : \mathcal{C} \rightarrow \mathcal{D}$  are concrete functors. The pair  $(F, G)$  is called a *Galois correspondence* between  $\mathcal{C}$  and  $\mathcal{D}$  if for each  $Y \in \mathcal{C}$ ,  $id_Y : F \circ G(Y) \rightarrow Y$  is a  $\mathcal{C}$ -morphism, and for each  $X \in \mathcal{D}$ ,  $id_X : X \rightarrow G \circ F(X)$  is a  $\mathcal{D}$ -morphism.

If  $(F, G)$  is a Galois correspondence, then it is easy to check that  $F$  is a left adjoint of  $G$ , or equivalently that  $G$  is a right adjoint of  $F$ .

Let **BFC** be denote the category of  $L$ -fuzzy bi-closure spaces and bi-closure mappings for morphisms.

Let **BCS** be denote the category of  $L$ -fuzzy bi-closure systems and continuous mappings for morphisms.

**THEOREM 3.7.** (1)  $\mathbf{F} : \mathbf{BFC} \rightarrow \mathbf{BCS}$  defined as  $\mathbf{F}(X, C_X^r, C_X^l) = (X, S_{C_X^r}^r, S_{C_X^l}^l)$  is a functor.

(2)  $\mathbf{G} : \mathbf{BCS} \rightarrow \mathbf{BFC}$  defined as  $\mathbf{G}(X, S_X^r, S_X^l) = (X, C_{S_X^r}^r, C_{S_X^l}^l)$  is a functor.

(3) The pair  $(\mathbf{F}, \mathbf{G})$  is a Galois correspondence between  $\mathbf{BFC}$  and  $\mathbf{BCS}$ .

*Proof.* (1) and (2) follows from Theorem 3.5.

(3) By Theorem 3.4(2), if  $(X, S_X^r, S_X^l)$  is an  $L$ -fuzzy bi-closure system, then  $\mathbf{F}(\mathbf{G}(X, S_X^r, S_X^l)) = (X, S_{C_{S_X^r}^r}, S_{C_{S_X^l}^l}^l) \geq (X, S_X^r, S_X^l)$ . Hence, the identity map  $id_X : (X, S_{C_{S_X^r}^r}, S_{C_{S_X^l}^l}^l) = \mathbf{F}(\mathbf{G}(X, S_X^r, S_X^l)) \rightarrow (X, S_X^r, S_X^l)$  is a bi-continuous map. Moreover, if  $(X, C_X^r, C_X^l)$  is an  $L$ -fuzzy bi-closure system, by Theorem 3.4(1),  $\mathbf{G}(\mathbf{F}(X, C_X^r, C_X^l)) = (X, C_{S_{C_X^r}^r}, C_{S_{C_X^l}^l}^l) \geq (X, C_X^r, C_X^l)$ . Hence the identity map  $id_X : (X, C_{S_{C_X^r}^r}, C_{S_{C_X^l}^l}^l) = \mathbf{G}(\mathbf{F}(X, C_X^r, C_X^l)) \rightarrow (X, C_X^r, C_X^l)$  is a continuous map. Therefore  $(\mathbf{F}, \mathbf{G})$  is a Galois correspondence. □

**EXAMPLE 3.8.** Let  $M = \{(x, y) \in R^2 \mid y > 0\}$  be a set and we define an operation  $\otimes : M \times M \rightarrow M$  as follows:

$$(x_1, y_1) \otimes (x_2, y_2) = (x_1 + y_1x_2, y_1y_2).$$

Then  $(M, \otimes)$  is a group with  $e = (0, 1)$ ,  $(x, y)^{-1} = (-\frac{x}{y}, \frac{1}{y})$ .

We have a positive cone  $P = \{(a, b) \in R^2 \mid b = 1, a \geq 0, \text{ or } y > 1\}$  because  $P \cap P^{-1} = \{(0, 1)\}$ ,  $P \otimes P \subset P$ ,  $(a, b)^{-1} \otimes P \times (a, b) = P$  and  $P \cup P^{-1} = L$ . For  $(x_1, y_1), (x_2, y_2) \in M$ , we define

$$\begin{aligned} (x_1, y_1) &\leq (x_2, y_2) \\ \Leftrightarrow (x_1, y_1)^{-1} \otimes (x_2, y_2) &\in P, (x_2, y_2) \otimes (x_1, y_1)^{-1} \in P \\ \Leftrightarrow y_1 < y_2 \text{ or } y_1 = y_2, x_1 &\leq x_2. \end{aligned}$$

Then  $(M, \leq, \otimes)$  is a lattice-group. Put  $L = \{(x, y) \in M \mid (1, \frac{1}{2}) \leq (x, y) \leq (0, 1)\}$ . Then  $(L, \odot, \Rightarrow, \rightarrow, (1, \frac{1}{2}), (0, 1))$  is a generalized residuated lattice where  $(1, \frac{1}{2})$  is the least element and  $(0, 1)$  is the greatest



element from the following statements:

$$\begin{aligned} (x_1, y_1) \odot (x_2, y_2) &= (x_1, y_1) \otimes (x_2, y_2) \vee (1, \frac{1}{2}) \\ &= (x_1 + y_1x_2, y_1y_2) \vee (1, \frac{1}{2}), \\ (x_1, y_1) \Rightarrow (x_2, y_2) &= ((x_1, y_1)^{-1} \otimes (x_2, y_2)) \wedge (0, 1) \\ &= (\frac{-x_1+x_2}{y_1}, \frac{y_2}{y_1}) \wedge (0, 1), \\ (x_1, y_1) \rightarrow (x_2, y_2) &= ((x_2, y_2) \otimes (x_1, y_1)^{-1}) \wedge (0, 1) \\ &= (x_2 - \frac{x_1y_2}{y_1}, \frac{y_2}{y_1}) \wedge (0, 1). \end{aligned}$$

It is not commutative because

$$(\frac{2}{3}, \frac{3}{4}) \odot (4, \frac{1}{2}) = (3 + \frac{2}{3}, \frac{3}{8}) \neq (4, \frac{1}{2}) \odot (\frac{2}{3}, \frac{3}{4}) = (4 + \frac{1}{3}, \frac{3}{8}).$$

Furthermore, we have  $(x, y) = (x, y)^{\circ} = (x, y)^{\circ*}$  from:

$$\begin{aligned} (x, y)^* &= (x, y) \Rightarrow (1, \frac{1}{2}) = (\frac{-x+1}{y}, \frac{1}{2y}), \\ (x, y)^{\circ} &= (x, y) \rightarrow (1, \frac{1}{2}) = (1 - \frac{x}{2y}, \frac{1}{2y}). \end{aligned}$$

Let  $X = \{a, b, c\}$  and  $A \in L^X$  as follows:

$$A(a) = (1, 0.6), A(b) = (2, 0.8), A(c) = (0, 0.6).$$

Define two maps  $S^r, S^l : L^X \rightarrow L$  as follows:

$$\begin{aligned} S^r(B) &= \begin{cases} (0, 1), & \text{if } B = \alpha \Rightarrow A, \\ (1, \frac{1}{2}), & \text{otherwise,} \end{cases} \\ S^l(B) &= \begin{cases} (0, 1), & \text{if } B = \alpha \rightarrow A, \\ (1, \frac{1}{2}), & \text{otherwise.} \end{cases} \end{aligned}$$

Then  $(X, S^r, S^r)$  is an *L*-fuzzy bi-closure system. For each  $D \in L^X$ , by Lemma 2.3(9) and Theorem 3.3,

$$\begin{aligned} C_{S^r}(D) &= \bigwedge_{B \in L^X} (e_{L^X}^r(D, B) \odot S^r(B) \rightarrow B) \\ &= \bigwedge_{\alpha \in L} (e_{L^X}^r(D, \alpha \Rightarrow A) \rightarrow (\alpha \Rightarrow A)) \\ C_{S^l}(D) &= \bigwedge_{B \in L^X} (S^l(B) \odot e_{L^X}^l(D, B) \Rightarrow B) \\ &= \bigwedge_{\alpha \in L} (e_{L^X}^l(D, \alpha \rightarrow A) \Rightarrow (\alpha \rightarrow A)) \end{aligned}$$

By Theorems 3.3 and 3.4,  $(X, C_{S^r}, C_{S^l})$  is an *L*-fuzzy bi-closure space. Moreover, by Theorem 3.4, we have

$$\begin{aligned} S_{C_{S^r}}^r(D) &= e_{L^X}^r(C_{S^r}(D), D) \\ &= e_{L^X}^r(\bigwedge_{\alpha \in L} (e_{L^X}^r(D, \alpha \Rightarrow A) \rightarrow (\alpha \Rightarrow A)), D), \\ S_{C_{S^l}}^l(D) &= e_{L^X}^l(C_{S^l}(D), D) \\ &= e_{L^X}^l(\bigwedge_{\alpha \in L} (e_{L^X}^l(D, \alpha \rightarrow A) \Rightarrow (\alpha \rightarrow A)), D). \end{aligned}$$

Since  $S_{C_{Sr}}^r(\alpha \Rightarrow A) = S_{C_{Sl}}^l(D)(\alpha \Rightarrow A) = (0, 1)$ , we have

$$S_{C_{Sr}}^r \geq S^r, \quad S_{C_{Sl}}^l \geq S^l.$$

### References

- [1] R. Bělohlávek, *Fuzzy Relational Systems*, Kluwer Academic Publishers, New York, 2002.
- [2] R. Bělohlávek, *Fuzzy Galois connection*, Math. Log. Quart., **45** (2000), 497–504.
- [3] R. Bělohlávek, *Fuzzy closure operator*, J. Math. Anal. Appl. **262** (2001), 473–486.
- [4] R. Bělohlávek, *Lattices of fixed points of Galois connections*, Math. Logic Quart. **47** (2001), 111–116.
- [5] L. Biacino and G. Gerla, *Closure systems and L-subalgebras*, Inf. Sci. **33** (1984), 181–195.
- [6] L. Biacino and G. Gerla, *An extension principle for closure operators*, J. Math. Anal. Appl. **198** (1996), 1–24.
- [7] G. Gerla, *Graded consequence relations and fuzzy closure operators*, J. Appl. Non-classical Logics **6** (1996), 369–379.
- [8] Jinming Fang and Yueli Yue, *L-fuzzy closure systems*, Fuzzy Sets and Systems **161** (2010), 1242–1252.
- [9] G. Georgescu and A. Popescue, *Non-commutative Galois connections*, Soft Computing **7** (2003), 458–467.
- [10] G. Georgescu and A. Popescue, *Non-dual fuzzy connections*, Arch. Math. Log. **43** (2004), 1009–1039.
- [11] P. Hájek, *Metamathematics of Fuzzy Logic*, Kluwer Academic Publishers, Dordrecht, 1998.
- [12] U. Höhle and E. P. Klement, *Non-classical logic and their applications to fuzzy subsets*, Kluwer Academic Publisher, Boston, 1995.
- [13] U. Höhle and S.E. Rodabaugh, *Mathematics of Fuzzy Sets: Logic, Topology, and Measure Theory, The Handbooks of Fuzzy Sets Series 3*, Kluwer Academic Publishers, Boston, 1999.
- [14] H. Lai and D. Zhang, *Concept lattices of fuzzy contexts: Formal concept analysis vs. rough set theory*, Int. J. Approx. Reasoning **50** (2009), 695–707.
- [15] C.J. Mulvey, *Quantales*, Suppl. Rend. Cric. Mat. Palermo Ser.II 12,1986,99-104.
- [16] M. Ward and R.P. Dilworth, *Residuated lattices*, Trans. Amer. Math. Soc. **45** (1939), 335–354.
- [17] E. Turunen, *Mathematics Behind Fuzzy Logic*, A Springer-Verlag Co., 1999.

**Jung Mi Ko**

Department of Mathematics  
Gangneung-Wonju National University  
Gangneung 25457, Korea  
*E-mail*: [jmko@gwnu.ac.kr](mailto:jmko@gwnu.ac.kr)

**Yong Chan Kim**

Department of Mathematics  
Gangneung-Wonju National University  
Gangneung 25457, Korea  
*E-mail*: [yck@gwnu.ac.kr](mailto:yck@gwnu.ac.kr)