

SOME METRIC ON EINSTEIN LORENTZIAN WARPED PRODUCT MANIFOLDS

SOO-YOUNG LEE

ABSTRACT. In this paper, let $M = B \times_{f^2} F$ be an Einstein Lorentzian warped product manifold with 2-dimensional base. We study the geodesic completeness of some metric with constant curvature. First of all, we discuss the existence of nonconstant warping functions on M . As the results, we have some metric g admits nonconstant warping functions f . Finally, we consider the geodesic completeness on M .

1. Introduction

R.L. Bishop and B. O’Neill introduced singly warped products or simply warped products to construct Riemannian manifolds with negative sectional curvature([5]). Later, we study the existence of some metric on Riemannian warped product manifolds([7], [12], [18]). And we consider the existence and the completeness of some metric on Lorentzian warped product manifolds([2], [3], [4], [8], [11], [14], [15], [16], [17], [19], [25], [26]).

In the present work, we study multiply warped products or multiply-warped products. One can also generalize singly warped products to

Received November 16, 2018. Revised December 20, 2019. Accepted December 24, 2019.

2010 Mathematics Subject Classification: 53C25, 53C50, 58E10, 53C21, 58D17.

Key words and phrases: Einstein manifold, Lorentzian warped product manifold, geodesics, warping function, multiply warped product.

© The Kangwon-Kyungki Mathematical Society, 2019.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

multiply warped products. A multiply warped product (M, g) is a product manifold of the form $M = B \times_{f_1} F_1 \times \cdots \times_{f_m} F_m$ with the metric $g = g_B \oplus f_1^2 g_{F_1} \oplus \cdots \oplus f_m^2 g_{F_m}$, where for each $i \in \{1, \dots, m\}$, $f_i : B \rightarrow (0, \infty)$ is smooth and (F_i, g_{F_i}) is a pseudo-Riemannian manifold. In particular, when $B = (a, b)$ with the negative definite metric $g_B = -dt^2$, the corresponding multiply warped product $M = (a, b) \times_{f_1} F_1 \times \cdots \times_{f_m} F_m$ with the metric $g = -dt^2 \oplus f_1^2 g_{F_1} \oplus \cdots \oplus f_m^2 g_{F_m}$ is called a Lorentzian multiply warped product, where for each $i \in \{1, \dots, m\}$, (F_i, g_{F_i}) is a Riemannian manifold and $-\infty \leq a < b \leq \infty$ ([27]).

In a recently, we study an Einstein manifold. We obtain some results an Einstein warped product manifold ([6], [9], [10], [13], [20], [21], [22], [23]). In [1], the author may also consider for that purpose special case of an Einstein warped product manifold $M = B \times_{f^2} F$ with 2-dimensional base, $B = (a, b) \times_{f^2} \mathbb{R}$, where $-\infty \leq a < b \leq \infty$. And we study the existence of nonconstant warping functions on M ([24]).

In this paper, we study an Einstein Lorentzian warped product manifold $M = B \times_{f^2} F$ with 2-dimensional base, $B = (a, b) \times_{f^2} \mathbb{R}$ when (a, b) with the negative definite metric $-dt^2$, where $-\infty \leq a < b \leq \infty$. First of all, we study the existence of nonconstant warping functions f depends on the signs of λ_0 . As a results, we have some metric g admits nonconstant warping functions f . Finally, we consider the geodesic completeness on M .

2. Preliminaries

We denote by Ric_F be the Ricci curvature of (F, g_F) and Ric_B be the Ricci curvature of (B, g_B) . We denote by Ric^B and Ric^F the lifts to M of Ricci curvatures of B and F , respectively.

PROPOSITION 2.1. *The Ricci curvature Ric of the warped product manifold $M = B \times_{f^2} F$ satisfies*

- (i) $Ric(V, W) = Ric^F(V, W) + g(V, W) \left[\left(\frac{\Delta f}{f} - (p-1) \frac{\|df\|^2}{f^2} \right) \pi \right]$,
- (ii) $Ric(X, V) = 0$,

$$(iii) Ric(X, Y) = Ric^B(X, Y) - \frac{p}{f}H^f(X, Y)$$

for any vertical vectors V, W and any horizontal vectors X, Y . We are defined by df is the gradient of f for g_B and H^f is the Hessian of f for g_B . We denote by Δf is the Laplacian of f for g_B and $p = \dim F$ ([1]).

COROLLARY 2.2. *The warped product $M = B \times_{f^2} F$ is Einstein manifold (with $Ric = \lambda g$) if and only if g_F, g_B and f satisfy*

$$(i) (F, g_F) \text{ is Einstein (with } Ric_F = \lambda_0 g_F),$$

$$(ii) \frac{\Delta f}{f} - (p - 1) \frac{\|df\|^2}{f^2} + \frac{\lambda_0}{f^2} = \lambda,$$

$$(iii) Ric_B - \frac{p}{f}H^f = \lambda g_B.$$

Obviously, (ii) and (iii) are two differential equations for f on (B, g_B) ([1]).

REMARK 2.3. Using Corollary 2.2 (ii) and (iii), we replace the unique equation

$$(2.1) Ric_B - \frac{p}{f}H^f = \frac{1}{2} [s_B + 2p \frac{\Delta f}{f} - p(p - 1) \frac{\|df\|^2}{f^2} + p \frac{\lambda_0}{f^2} - (p + q - 2)\lambda] g_B,$$

where $q = \dim B$.

PROPOSITION 2.4. *In the special case of a warped product $B \times_{f^2} F$ over 2-dimensional base, we have $Ric_B = \frac{1}{2}s_B g_B$ and $q = 2$. Hence equation (2.1) implies that*

$$(2.2) H^f = -\frac{1}{2} [2\Delta f - (p - 1) \frac{\|df\|^2}{f} + \frac{\lambda_0}{f} - \lambda f] g_B.$$

LEMMA 2.5. *Let $B = (a, b) \times_{f'(t)^2} \mathbb{R}$ be 2-dimensional manifold for $t \in (a, b)$ and $u \in \mathbb{R}$, where $-\infty \leq a < b \leq \infty$. On (B, g_B) the equation $H^f = -f'' g_B$ admits a nonconstant solution f if and only if, locally at*

points where $df \neq 0$, there exists local coordinates (t, u) such that f is a function of t alone.

Proof. By a proof similar Lemma 9.117 in [1], then $H^f = -f''g_B$. \square

With the notations of the Lemma 2.5, we have an ordinary differential equation for in the variable t

$$(2.3) \quad 2f''(t) + (p - 1)\frac{f'(t)^2}{f(t)} + \frac{\lambda_0}{f(t)} - \lambda f(t) = 0,$$

where $\|df\|^2 = -[f'(t)]^2$ and $\Delta f = 2f''(t)$.

The following notation and Remark 2.6 are needed to show the geodesic completeness.

NOTATION. Let $M = (a, b) \times_{f_1} F_1 \times \cdots \times_{f_m} F_m$ be a Lorentzian multiply warped product with metric $g = -dt^2 \oplus f_1^2 g_{F_1} \oplus \cdots \oplus f_m^2 g_{F_m}$, where $-\infty \leq a < b \leq \infty$. If $\mathcal{B} = \{f_1, \dots, f_m\}$ and for some $k \in \{1, \dots, m\}$ and for some subset $\{\bar{f}_1, \dots, \bar{f}_k\}$ of \mathcal{B} , then

$$r[\bar{f}_1, \dots, \bar{f}_k] = \prod_{i=1}^k \bar{f}_i \quad \text{and} \quad h[\bar{f}_1, \dots, \bar{f}_k] = \sum_{i=1}^k \bar{f}_i^2.$$

Also, it is assumed that $h[\bar{f}_1] = 1$ for any \bar{f}_1 ([27]).

REMARK 2.6. Let $M = (a, b) \times_{f_1} F_1 \times \cdots \times_{f_m} F_m$ be a Lorentzian multiply warped product with metric $g = -dt^2 \oplus f_1^2 g_{F_1} \oplus \cdots \oplus f_m^2 g_{F_m}$, where $-\infty \leq a < b \leq \infty$. Suppose that (F_i, g_{F_i}) is a complete Riemannian manifold for any $i \in \{1, \dots, m\}$ and $\mathcal{B} = \{f_1, \dots, f_m\}$. Then

every future directed time-like geodesic is future(respectively past) complete if and only if $\lim_{t \rightarrow b^-} \int_{t_1}^t \frac{r[\bar{f}_1, \dots, \bar{f}_k](s)}{\sqrt{r[\bar{f}_1, \dots, \bar{f}_k]^2(s) + h[\bar{f}_1, \dots, \bar{f}_k]}} ds = \infty$

(respectively $\lim_{t \rightarrow a^+} \int_t^{t_1} \frac{r[\bar{f}_1, \dots, \bar{f}_k](s)}{\sqrt{r[\bar{f}_1, \dots, \bar{f}_k]^2(s) + h[\bar{f}_1, \dots, \bar{f}_k]}} ds = \infty$) for some $t_1 \in (a, b)$ and for any $k \in \{1, \dots, m\}$ and for any subset $\{\bar{f}_1, \dots, \bar{f}_k\}$ of \mathcal{B} (cf. Theorem 4.8 in [27]).

3. The existence of nonconstant warping functions

Let $M = (a, b) \times_{f'(t)^2} \mathbb{R} \times_{f(t)^2} F$ be an Einstein Lorentzian warped product manifold, where $f(t)$ and $f'(t)$ are smooth functions and $-\infty \leq a < b \leq \infty$. Let $\dim F = p > 1$.

First of all, if we denote $f(t) = z(t)^{\frac{2}{p+1}}$, then equation (2.3) can be changed into

$$(3.1) \quad [z'(t)]^2 = -\frac{(p+1)^2 \lambda_0}{4(p-1)} z(t)^{2-\frac{4}{p+1}} + \frac{(p+1)\lambda}{4} [z(t)]^2,$$

where $z(t)$ is a positive function. Thus we study positive solution $z(t)$ of equation (3.1).

THEOREM 3.1. *Suppose that $\lambda_0 = 0$. If λ is a constant, then there exists a nonconstant solution $z(t)$ of equation (3.1).*

(i) *For $\lambda = 0$, there does not exist a nonconstant solution of equation (3.1).*

(ii) *For $\lambda > 0$, we have a solution $z(t) = e^{\pm\sqrt{\frac{(p+1)\lambda}{4}} t+c}$, where c is a constant.*

(iii) *For $\lambda < 0$, there does not exist a solution of equation (3.1).*

Proof. For $\lambda_0 = 0$, equation (3.1) implies that

$$(3.2) \quad [z'(t)]^2 = \frac{(p+1)\lambda}{4} [z(t)]^2.$$

(i) For $\lambda = 0$, equation (3.2) implies that $[z'(t)]^2 = 0$ and $z'(t) = 0$. An integration gives $z(t) = c$, where c is a positive constant. Because $z(t) = c$ is not a nonconstant, thus $z(t) = c$ is not our solution.

(ii) For $\lambda > 0$, equation (3.2) implies that we get $z'(t) = \pm\sqrt{\frac{(p+1)\lambda}{4}} u(t)$.

An integration gives $\ln |z(t)| = \pm\sqrt{\frac{(p+1)\lambda}{4}} t + c$, where c is a constant.

Therefore we have $z(t) = e^{\pm\sqrt{\frac{(p+1)\lambda}{4}} t+c}$, where c is a constant.

(iii) For $\lambda < 0$, equation (3.2) implies that $[z'(t)]^2 < 0$. Which is a contradiction. Hence there does not exist a solution of equation (3.1). \square

REMARK 3.2. Let M be an Einstein Lorentzian warped product manifold. From above Theorem 3.1 (ii), for $\lambda_0 = 0$ and $\lambda > 0$, we have that equation (2.3) satisfies a nonconstant warping function $f(t) = e^{\pm\sqrt{\frac{\lambda}{p+1}} t + \frac{2c}{p+1}}$ on $(-\infty, \infty)$, where c is a constant.

THEOREM 3.3. Suppose that $\lambda_0 > 0$. If λ is a constant, then there exists a nonconstant solution $z(t)$ of equation (3.1).

(i) For $\lambda \leq 0$, there does not exist a solution of equation (3.1).

(ii) For $\lambda > 0$, we have $z(t) = \left(\sqrt{\frac{(p+1)\lambda_0}{(p-1)\lambda}} \cosh\left(\sqrt{\frac{\lambda}{p+1}} t + c\right) \right)^{\frac{p+1}{2}}$, where c is a constant.

Proof. (i) For $\lambda \leq 0$, equation (3.1) implies that $[z'(t)]^2 < 0$. Which is a contradiction. Therefore there does not exist a solution of equation (3.1).

(ii) For $\lambda > 0$, first of all, equation (3.1) implies that we rewritten as

$$\int \frac{1}{z(t) \sqrt{-\frac{(p+1)^2 \lambda_0}{4(p-1)} z(t)^{-\frac{4}{p+1}} + \frac{(p+1)\lambda}{4}}} du = \pm \int dt.$$

Putting $\frac{(p+1)^2 \lambda_0}{4(p-1)} = I > 0$ and $\frac{(p+1)\lambda}{4} = J > 0$, then we get the equation

$$\int \frac{1}{z(t) \sqrt{J - I z(t)^{-\frac{4}{p+1}}}} du = \pm \int dt.$$

By using trigonometric substitution, $z(t)^{-\frac{2}{p+1}} = \frac{\sqrt{J}}{\sqrt{I}} \sin \theta$, then we obtain

$$- \int \csc \theta d\theta = \pm \int \frac{2\sqrt{J}}{p+1} dt.$$

Upon integration, we become $\ln | \csc \theta + \cot \theta | = \pm \frac{2\sqrt{J}}{p+1} t + c$, where c is a constant. Here we have $\ln |\sqrt{J}z(t)^{\frac{2}{p+1}} + \sqrt{Jz(t)^{\frac{4}{p+1}} - I}| = \pm \frac{2\sqrt{J}}{p+1} t + c + \ln \sqrt{I}$, where c is a constant.

Therefore we have $z(t) = \left(\sqrt{\frac{(p+1)\lambda_0}{(p-1)\lambda}} \cosh\left(\sqrt{\frac{\lambda}{p+1}}t + c\right) \right)^{\frac{p+1}{2}}$, where c is a constant. □

REMARK 3.4. Let M be an Einstein Lorentzian warped product manifold. From above Theorem 3.3 (ii), for $\lambda_0 > 0$ and $\lambda > 0$, we have that equation (2.3) satisfies a nonconstant warping function $f(t) = \sqrt{\frac{(p+1)\lambda_0}{(p-1)\lambda}} \cosh\left(\sqrt{\frac{\lambda}{p+1}}t + c\right)$ on $\left(-\sqrt{\frac{p+1}{\lambda}}c, \infty\right)$, where c is a constant.

THEOREM 3.5. Suppose that $\lambda_0 < 0$. If λ is a constant, then there exist nonconstant solutions $z(t)$ of equation (3.1).

(i) For $\lambda = 0$, we have $z(t) = \left(\pm \sqrt{\frac{-\lambda_0}{p-1}}t + \frac{2c}{p+1}\right)^{\frac{p+1}{2}}$, where c is a constant.

(ii) For $\lambda > 0$, we get $z(t) = \left(\sqrt{\frac{-(p+1)\lambda_0}{(p-1)\lambda}} \sinh\left(\pm \sqrt{\frac{\lambda}{p+1}}t + c\right)\right)^{\frac{p+1}{2}}$, where c is a constant.

(iii) For $\lambda < 0$, we become $z(t) = \left(\sqrt{\frac{(p+1)\lambda_0}{(p-1)\lambda}} \cos\left(\pm \sqrt{\frac{-\lambda}{p+1}}t + c\right)\right)^{\frac{p+1}{2}}$, where c is a constant.

Proof. (i) For $\lambda = 0$, equation (3.1) implies that we have equation

$$z'(t) = \pm \sqrt{\frac{-(p+1)^2\lambda_0}{4(p-1)}} z(t)^{1-\frac{2}{p+1}}.$$

Therefore we have $z(t) = \left(\pm \sqrt{\frac{-\lambda_0}{p-1}}t + \frac{2c}{p+1}\right)^{\frac{p+1}{2}}$, where c is a constant.

(ii) For $\lambda > 0$. By a proof similar to Theorem 3.3 (ii), equation (3.1) implies that we rewritten as

$$\int \frac{1}{z(t) \sqrt{-\frac{(p+1)^2 \lambda_0}{4(p-1)} z(t)^{-\frac{4}{p+1}} + \frac{(p+1)\lambda}{4}}} du = \pm \int dt.$$

Putting $-\frac{(p+1)^2 \lambda_0}{4(p-1)} = I > 0$ and $\frac{(p+1)\lambda}{4} = J > 0$, then we have the equation

$$\int \frac{1}{z(t) \sqrt{Iz(t)^{-\frac{4}{p+1}} + J}} du = \pm \int dt.$$

By using trigonometric substitution, $z(t)^{-\frac{2}{p+1}} = \frac{\sqrt{J}}{\sqrt{I}} \tan \theta$, then we obtain

$$-\int \csc \theta d\theta = \pm \int \frac{2\sqrt{J}}{p+1} dt.$$

Therefore we have $z(t) = \left(\sqrt{\frac{-(p+1)\lambda_0}{(p-1)\lambda}} \sinh(\pm \sqrt{\frac{\lambda}{p+1}} t + c) \right)^{\frac{p+1}{2}}$, where c is a constant.

(iii) For $\lambda < 0$. By a proof similar to Theorem 3.3 (ii) and Theorem 3.5 (ii), putting $-\frac{(p+1)^2 \lambda_0}{4(p-1)} = I > 0$ and $-\frac{(p+1)\lambda}{4} = J > 0$, then we have

$$\int \frac{1}{z(t) \sqrt{Iz(t)^{-\frac{4}{p+1}} - J}} du = \pm \int dt.$$

By using trigonometric substitution, $z(t)^{-\frac{2}{p+1}} = \frac{\sqrt{J}}{\sqrt{I}} \sec \theta$, then we get

$$\int d\theta = \pm \int \frac{2\sqrt{J}}{p+1} dt.$$

Therefore we have $z(t) = \left(\sqrt{\frac{(p+1)\lambda_0}{(p-1)\lambda}} \cos(\pm \sqrt{\frac{-\lambda}{p+1}} t + c) \right)^{\frac{p+1}{2}}$, where c is a constant. \square

REMARK 3.6. Let M be an Einstein Lorentzian warped product manifold. From above Theorem 3.5, we consider that equation (2.3) satisfies nonconstant warping functions $f(t)$.

- (i) For $\lambda_0 < 0$ and $\lambda = 0$, we become $f(t) = \sqrt{\frac{-\lambda_0}{p-1}} t + \frac{2c}{p+1}$ on $\left(-\frac{2c}{p+1} \sqrt{\frac{p-1}{-\lambda_0}}, \infty\right)$, where c is a constant.
- (ii) For $\lambda_0 < 0$ and $\lambda > 0$, we get $f(t) = \sqrt{\frac{-(p+1)\lambda_0}{(p-1)\lambda}} \sinh\left(\sqrt{\frac{\lambda}{p+1}} t + c\right)$ on $\left(-\sqrt{\frac{p+1}{\lambda}} c, \infty\right)$, where c is a constant.
- (iii) For $\lambda_0 < 0$ and $\lambda < 0$, we have $f(t) = \sqrt{\frac{(p+1)\lambda_0}{(p-1)\lambda}} \cos\left(\sqrt{\frac{-\lambda}{p+1}} t + c\right)$ on $\left((2n\pi - \frac{\pi}{4} - c)\sqrt{\frac{p+1}{-\lambda}}, (2n\pi + \frac{3\pi}{4} - c)\sqrt{\frac{p+1}{-\lambda}}\right)$, where c is a constant and n is an integer.

REMARK 3.7. The behaviour of the nonconstant warping functions depends on the signs of λ_0 and λ . Then we reduced to the following sets of solutions besides the constant case when $c = 0$ and $p > 1$ is an integer.

λ_0	0	$p - 1$	$-(p - 1)$	$-(p - 1)$	$-(p - 1)$
λ	$p + 1$	$p + 1$	0	$p + 1$	$-(p + 1)$
$f(t)$	$e^{\pm t}$	$\cosh(t)$	t	$\sinh(t)$	$\cos(t)$

4. The existence and the completeness of some metric

Let $M = (a, b) \times_{f'(t)^2} \mathbb{R} \times_{f(t)^2} F$ be an Einstein Lorentzian warped product manifold, where $f(t)$ and $f'(t)$ are smooth functions and $-\infty \leq a < b \leq \infty$. Let $\dim F = p > 1$.

REMARK 4.1. From the Remark 3.2, we have positive smooth functions $f(t)$ and $f'(t)$. Then we have the metric

$$g = -dt^2 + \frac{\lambda}{p+1} e^{\sqrt{\frac{\lambda}{p+1}} 2t + \frac{4c}{p+1}} du^2 + e^{\sqrt{\frac{\lambda}{p+1}} 2t + \frac{4c}{p+1}} g_F,$$

where c is a constant.

THEOREM 4.2. *Let M be an Einstein Lorentzian warped product manifold. Suppose that (\mathbb{R}, du^2) and (F, g_F) are complete. If $\lambda_0 = 0$ and $\lambda > 0$, then on M the resulting metric is that every future directed time-like geodesics is future (or past) complete.*

Proof. For $\lambda_0 = 0$ and $\lambda > 0$, the metric of Remark 4.1 simplifies to

$$g = -dt^2 + \alpha^2 e^{2\alpha t} du^2 + e^{2\alpha t} g_F$$

on $(-\infty, \infty)$, where α is a positive constant.

Let $\mathcal{B} = \{\alpha e^{\alpha t}, e^{\alpha t}\}$, where α is a positive constant. For some $t_1 \in (-\infty, \infty)$, then

(i) $\int_{t_1}^{\infty} \frac{\alpha e^{\alpha t}}{\sqrt{\alpha^2 e^{2\alpha t} + 1}} dt \geq \int_{t_1}^{\infty} \frac{1}{\sqrt{2}} dt = +\infty,$

(ii) $\int_{t_1}^{\infty} \frac{e^{\alpha t}}{\sqrt{e^{2\alpha t} + 1}} dt \geq \int_{t_1}^{\infty} \frac{1}{\sqrt{2}} dt = +\infty,$

(iii) $\int_{t_1}^{\infty} \frac{\alpha e^{\alpha t} e^{\alpha t}}{\sqrt{\alpha^2 e^{2\alpha t} + e^{2\alpha t} + \alpha^2 e^{2\alpha t} e^{2\alpha t}}} dt \geq \int_{t_1}^{\infty} \frac{1}{\sqrt{3}} dt = +\infty,$ where α is a positive constant.

Therefore from the Remark 2.6, on M every future directed time-like geodesic is future complete. On the other hand, by similar methods, on M every future directed time-like geodesic is past incomplete. □

REMARK 4.3. From the Remark 3.4, we have positive smooth functions $f(t)$ and $f'(t)$. Then we have the metric

$$g = -dt^2 + \frac{\lambda_0}{p-1} \sinh^2\left(\sqrt{\frac{\lambda}{p+1}} t + c\right) du^2 + \frac{(p+1)\lambda_0}{(p-1)\lambda} \cosh^2\left(\sqrt{\frac{\lambda}{p+1}} t + c\right) g_F,$$

where c is a constant.

THEOREM 4.4. *Let M be an Einstein Lorentzian warped product manifold. Suppose that (\mathbb{R}, du^2) and (F, g_F) are complete. If $\lambda_0 > 0$ and $\lambda > 0$, then on M the resulting metric is that every future directed time-like geodesics is future (or past) complete.*

Proof. For $\lambda_0 > 0$ and $\lambda > 0$, the metric of Remark 4.3 simplifies to

$$g = -dt^2 + \alpha^2 \sinh^2(\alpha t) du^2 + \cosh^2(\alpha t) g_F$$

on $(0, \infty)$, where α is a positive constant.

Let $\mathcal{B} = \{\alpha \sinh(\alpha t), \cosh(\alpha t)\}$, where α is a positive constant. For some $t_1 \in (0, \infty)$, then

(i)
$$\int_{t_1}^{\infty} \frac{\alpha \sinh(\alpha t)}{\sqrt{\alpha^2 \sinh^2(\alpha t) + 1}} dt \geq \int_{t_1}^{\infty} \frac{1}{\sqrt{2}} dt = +\infty,$$

(ii)
$$\int_{t_1}^{\infty} \frac{\cosh(\alpha t)}{\sqrt{\cosh^2(\alpha t) + 1}} dt \geq \int_{t_1}^{\infty} \frac{1}{\sqrt{2}} dt = +\infty,$$

(iii)
$$\int_{t_1}^{\infty} \frac{\alpha \sinh(\alpha t) \cosh(\alpha t)}{\sqrt{\alpha^2 \sinh^2(\alpha t) + \cosh^2(\alpha t) + \alpha^2 \sinh^2(\alpha t) \cosh^2(\alpha t)}} dt \geq \int_{t_1}^{\infty} \frac{1}{\sqrt{3}} dt$$

$= +\infty$, where α is a positive constant.

Therefore from the Remark 2.6, on M every future directed time-like geodesic is future complete but past incomplete. □

REMARK 4.5. From the Remark 3.6, we have positive smooth functions $f(t)$ and $f'(t)$. Then we have the metrics.

(i) For $\lambda_0 < 0$ and $\lambda = 0$, we have

$$g = -dt^2 + \frac{-\lambda_0}{p-1} du^2 + \left(\sqrt{\frac{-\lambda_0}{p-1}} t + \frac{2c}{p+1} \right)^2 g_F,$$

where c is a constant.

(ii) For $\lambda_0 < 0$ and $\lambda > 0$, we become

$$g = -dt^2 + \frac{-\lambda_0}{p-1} \cosh^2\left(\sqrt{\frac{\lambda}{p+1}} t+c\right) du^2 + \frac{-(p+1)\lambda_0}{(p-1)\lambda} \sinh^2\left(\sqrt{\frac{\lambda}{p+1}} t+c\right) g_F,$$

where c is a constant.

(iii) For $\lambda_0 < 0$ and $\lambda < 0$, we get

$$g = -dt^2 + \frac{-\lambda_0}{p-1} \sin^2\left(\sqrt{\frac{-\lambda}{p+1}} t+c\right) du^2 + \frac{(p+1)\lambda_0}{(p-1)\lambda} \cos^2\left(\sqrt{\frac{-\lambda}{p+1}} t+c\right) g_F,$$

where c is a constant and n is an integer.

THEOREM 4.6. *Let M be an Einstein Lorentzian warped product manifold. Suppose that (\mathbb{R}, du^2) and (F, g_F) are complete. If $\lambda_0 < 0$ and $\lambda = 0$, then on M the resulting metric is that every future directed time-like geodesic is future (or past) complete.*

Proof. For $\lambda_0 < 0$ and $\lambda = 0$, from the Remark 4.5 (i), the warping function $f(t)$ is a linear function and $f'(t)$ is a constant function. Because $f'(t)$ is not a nonconstant, thus we can not discuss geodesic complete. \square

THEOREM 4.7. *Let M be an Einstein Lorentzian warped product manifold. Suppose that (\mathbb{R}, du^2) and (F, g_F) are complete. If $\lambda_0 < 0$ and $\lambda > 0$, then on M the resulting metric is that every future directed time-like geodesic is future (or past) complete.*

Proof. For $\lambda_0 < 0$ and $\lambda > 0$, the metric of Remark 4.5 (ii) simplifies to

$$g = -dt^2 + \alpha^2 \cosh^2(\alpha t) du^2 + \sinh^2(\alpha t) g_F,$$

on $(0, \infty)$, where α is a positive constant.

Let $\mathcal{B} = \{\alpha \cosh(\alpha t), \sinh(\alpha t)\}$, where α is a positive constant. By a proof similar to Theorem 4.4, for some $t_1 \in (0, \infty)$, from the Remark 2.6 implies that on M every future directed time-like geodesic is future complete but past incomplete. \square

THEOREM 4.8. *Let M be an Einstein Lorentzian warped product manifold. Suppose that (\mathbb{R}, du^2) and (F, g_F) are complete. If $\lambda_0 < 0$ and $\lambda < 0$, then on M the resulting metric is that every future directed time-like geodesic is not future (or past) complete.*

Proof. For $\lambda_0 < 0$ and $\lambda < 0$, from the Remark 4.5 (iii), we have $f(t) = \cos\left(\sqrt{\frac{-\lambda}{p+1}} t + c\right)$ and $f'(t) = \sqrt{\frac{-\lambda}{p+1}} \sin\left(\sqrt{\frac{-\lambda}{p+1}} t + c\right)$, where c is a constant. Because we can consider the existence of a nonconstant warping function on only a finite interval, thus we can not discuss the completeness. □

REMARK 4.9. Let $M = (a, b) \times_{f'(t)^2} \mathbb{R} \times_{f(t)^2} F$ be an Einstein Lorentzian warped product manifold. The behaviour of the metrics depends on the signs of λ_0 and λ . Then we reduced to the following sets of metrics besides the constant case when $c = 0$ and $p > 1$ is an integer.

	λ_0	λ	metric
(i)	0	$p + 1$	$g = -dt^2 + e^{2t} du^2 + e^{2t} g_F$
(ii)	$p - 1$	$p + 1$	$g = -dt^2 + \sinh^2 t du^2 + \cosh^2 t g_F$
(iii)	$-(p - 1)$	0	$g = -dt^2 + du^2 + t^2 g_F$
(iv)	$-(p - 1)$	$p + 1$	$g = -dt^2 + \cosh^2 t du^2 + \sinh^2 t g_F$
(v)	$-(p - 1)$	$-(p + 1)$	$g = -dt^2 + \sin^2 t du^2 + \cos^2 t g_F$

References

- [1] A.L. Besse, *Einstein manifold*, Springer-Verlag, New York, 1987.
- [2] J.K. Beem and P.E. Ehrlich, *Global Lorentzian geometry*, Pure and Applied Mathematics, Vol.67, Dekker, New York, 1981.
- [3] J.K. Beem, P.E. Ehrlich, and K.L. Easley, *Global Lorentzian Geometry* (2nd ed)., Marcel Dekker, Inc., New York (1996).
- [4] J.K. Beem, P.E. Ehrlich, and Th.G. Powell, *Warped product manifolds in relativity*, Selected Studies (Th.M.Rassias, eds.), North-Holland, 1982, 41–56.
- [5] R.L. Bishop and O’Neill, *Manifolds if negative curvature*, Trans., Am. Math. Soc. **145** (1969), 1–49.
- [6] J. Case, Y.J. Shu, and G. Wei, *Rigidity of quasi-Einstein metricc*, Diff. Geo. and its applications **29** (2011), 93–100.

- [7] E.H. Choi, Y.H. Yang, and S.Y. Lee, *The nonexistence of warping functions on Riemannian warped product manifolds*, J. Chungcheong Math. Soc. **24** (2) (2011), 171–185.
- [8] P.E. Ehrlich, Y.T. Jung, and S.B. Kim, *Constant scalar curvatures on warped product manifolds*, Tsukuba J. Math. **20** (1) (1996), 239–256.
- [9] C. He, P.Petersen, and W. Wylie, *On the classification of warped product Einstein metrics*, math.DG. **24**, Jan.(2011).
- [10] C. He, P.Petersen, and W. Wylie, *Uniqueness of warped product Einstein metrics and applications*, math.DG. **4**, Feb.(2013).
- [11] Y.T. Jung, *Partial differential equations on semi-Riemannian manifolds*, J. Math. Anal. Appl. **241** (2000), 238–253.
- [12] Y.T. Jung, S.H. Chae, and S.Y. Lee, *The existence of some metrics on Riemannian warped product manifolds with fiber manifold of class (B)*, Korean J. Math, **23** (4) (2015), 733–740.
- [13] Y.T. Jung, E.H. Choi, and S.Y. Lee, *Nonconstant warping functions on Einstein Lorentzian warped product manifold*, Honam Math. J. **39** (3) (2018), 447–456.
- [14] Y.T. Jung, A.R. Kim, and S.Y. Lee, *The existence and the completeness of some metrics on Lorentzian warped product manifolds with fiber manifold of class (B)*, Honam Math. J. **39** (2) (2017), 187–197.
- [15] Y.T. Jung, Y.J. Kim, S.Y. Lee, and C.G. Shin, *Scalar curvature on a warped product manifold*, Korean Annals of Math. **15** (1998), 167–176.
- [16] Y.T. Jung, Y.J. Kim, S.Y. Lee, and C.G. Shin, *Partial differential equations and scalar curvature on semi-Riemannian manifolds(I)*, J. Korea Soc. Math. Educ. Ser. B: Pre Appl. Math. **5** (2) (1998), 115–122.
- [17] Y.T. Jung, Y.J. Kim, S.Y. Lee, and C.G. Shin, *Partial differential equations and scalar curvature on semi-Riemannian manifold(II)*, J. Korea Soc. Math. Educ. Ser. B: Pre Appl. Math. **6** (2) (1999), 95–101.
- [18] Y.T. Jung, G.Y. Lee, A.R. Kim, and S.Y. Lee, *The existence of warping functions on Riemannian warped product manifolds with fiber manifold of class (B)*, Honam Mathematical J. **36** (3) (2014), 597–603.
- [19] Y.T. Jung, J.M. Lee, and G.Y. Lee, *The completeness of some metrics on Lorentzian warped product manifolds with fiber manifold of class (B)*, Honam Math. J. **37** (1) (2015), 127–134.
- [20] D.S. Kim, *Einstein warped product spaces*, Honam Mathematical J. **22** (1) (2000), 107–111.
- [21] D.S. Kim, *Compact Einstein warped product spaces*, Trends in Mathematics, Information center for Mathematical Sciences, **5** (2) December (2002), 1–5.
- [22] D.S. Kim and Y.H. Kim, *Compact Einstein warped product spaces with nonpositive scalar curvature*, proceeding of A.M.S. **131** (8) (2003), 2573–2576.
- [23] W.Kühnel and H.B. Rademacher, *Conformally Einstein product spaces*, math.DG. **12**, Jul.(2016).
- [24] S.Y. Lee, *Nonconstant warping functions on Einstein warped product manifolds with 2–dimensional base*, Korean J. Math. **26** (1) (2018), 75–85.
- [25] B. O’Neill, *Semi-Riemannian Geometry*, Academic, New York, 1983.

- [26] T.G. Powell, *Lorentzian manifolds with non-smooth metrics and warped products*, ph. D. thesis, Univ. of Missouri-Columbia (1982).
- [27] B. Ünal, *Multiply warped products*, J. Geom. Phys. **34** (2000), 287–301.

Soo-Young Lee

Department of Mathematics

Chosun University,

Kwangju, 61452, Republic of Korea.

E-mail: `skdmlskan@hanmail.net`