

CROSSED SEMIMODULES AND CAT^1 -MONOIDS

SEDAT TEMEL

ABSTRACT. The main idea of this paper is to introduce the notion of cat^1 -monoids and to prove that the category of crossed semimodules $\mathcal{C} = (A, B, \partial)$ where A is a group is equivalent to the category of cat^1 -monoids. This is a generalization of the well known equivalence between category of cat^1 -groups and that of crossed modules over groups.

1. Introduction

Crossed modules as defined by Whitehead [30, 31] have been widely used in homotopy theory [5], the theory of group representation (see [7] for a survey), in algebraic K-theory [13] and homological algebra [12, 15]. Crossed modules can be viewed as 2-dimensional groups [6]. In [9] Brown and Spencer proved that the category of internal groupoids within the groups, which are also called in [9] under the name of \mathcal{G} -groupoids and alternative names, quite generally used are group-groupoids [8] or 2-groups (see for example [2]) is equivalent to the category of crossed modules of groups. In [14] Loday defined an algebraic object called cat^1 -group as a group G with two endomorphisms s, t of G such that $st = t, ts = s$ and $[\text{Ker } s, \text{Ker } t] = 0$, where $[\text{Ker } s, \text{Ker } t]$ represents the commutator subgroup of G ; and proved that the categories of cat^1 -groups and crossed modules are equivalent.

Received February 7, 2019. Revised June 14, 2019. Accepted June 15, 2019.

2010 Mathematics Subject Classification: 20L05, 18D05, 18D35, 20J15.

Key words and phrases: Crossed module, crossed semimodule, cat^1 -group, cat^1 -monoid.

© The Kangwon-Kyungki Mathematical Society, 2019.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

In [25] Porter proved a similar result to one in [9] holds for certain algebraic categories, introduced by Orzech [23], which definition was adapted by him and called category of groups with operations. The equivalences of these categories given in [9] and [25] are very important and useful which enable to generalize some results on group-groupoids, equivalently cat^1 -groups, to the more general internal groupoids for a certain algebraic category (see for example [1, 18, 19] and [21]). There are also other generalizations of the Brown-Spencer Theorem [9]. First one is the generalization to the category of monoids, that is the natural equivalence of the category of crossed semimodules satisfying certain conditions and that of Schreier internal categories within the category of monoids which is given by Patchkoria [24]. The other one is the generalization to the category of monoid with operations given in [22] by Martins-Ferreira et. al. In [24], Patchkoria also defined Schreier internal groupoids in the category of monoids which are naturally equivalent to the category of crossed semimodules $\mathcal{C} = (A, B, \partial)$ where A is a group. This is the special case of the equivalence given in [26] by Porter. Topological aspect of the results of Patchkoria is given in [28]. See [29] for 2-categorical approach to Schreier internal categories within monoids using Schreier 2-categories with one object. Equivalences given in [3, 9, 24, 26, 28] are very important and enable one to interpret special objects such as normal subobject, quotient object, etc. and special morphisms such as covering, lifting, etc. which are known in one of the categories in equivalences. See for example [1, 17, 19, 20, 27] for such interpretations.

The natural equivalence between crossed modules and cat^1 -groups is useful for generalisation of crossed modules to higher dimensions (see [14]). In this note we introduce the notion of cat^1 -monoids and prove the natural equivalence between the category of cat^1 -monoids and the category of crossed semimodules $\mathcal{C} = (A, B, \partial)$ where A is a group. This is a new way of thinking crossed semimodules as simple algebraic structures.

2. Preliminaries

Let C be a finitely complete category. An *internal category* $D = (D_0, D_1, d_0, d_1, \varepsilon, \mu)$ in C consists of a set of objects D_0 and a set of morphisms D_1 together with morphisms $d_0, d_1: D_1 \rightarrow D_0$, $\varepsilon: D_0 \rightarrow D_1$ in C called the source, the target and the identity object maps,

respectively,

$$D_1 \begin{array}{c} \xleftarrow{\varepsilon} \\ \xrightarrow{s} \\ \xrightarrow{t} \end{array} D_0$$

such that $d_0\varepsilon = d_1\varepsilon = 1_{D_0}$ and a morphism $\mu: D_1 \times_{D_0} D_1 \rightarrow D_1$ of \mathcal{C} called the composition map (usually expressed as $\mu(f, g) = g \circ f$) where $D_1 \times_{D_0} D_1$ is the pullback of d_0, d_1 such that $\varepsilon d_0(f) \circ f = f = f \circ \varepsilon d_0(f)$ [16]. An *internal groupoid* in \mathcal{C} is an internal category in which all morphisms are invertible up to groupoid composition.

An internal category (hence a groupoid) in the category GP of groups is called group-groupoid. A group-groupoid can also be obtained as a group object in CAT .

A crossed module over groups is a pair of groups A, B with an action $\bullet: B \times A \rightarrow A$ of B on A denoted by $b \bullet a$ for $a \in A$ and $b \in B$ and a morphism $\partial: A \rightarrow B$ of groups satisfies the following conditions

- [CM 1] $\partial(b \bullet a) = b\partial(a)b^{-1}$
- [CM 2] $\partial(a) \bullet a_1 = aa_1a^{-1}$,

for $a, a_1 \in A$ and $b \in B$ [30, 31]. For the basic examples of crossed modules see [10].

The following theorem is proved in [9] by Brown and Spencer.

THEOREM 2.1. *The category GPGD of group-groupoids is equivalent to the category CM of crossed modules over groups.*

Let MON denote the category of monoids. Recall from [24] that a Schreier internal category $M = (M_0, M_1, d_0, d_1, \varepsilon, \mu)$ in MON is an internal category which satisfies the Schreier condition: for any $f \in M_1$ there exists a unique $g \in \text{Ker } d_0$ such that

$$f = g\varepsilon d_0(f).$$

A Schreier internal groupoid in MON is a Schreier internal category in which all morphisms are invertible up to composition.

We recall the definition of crossed semimodules from [24]. A crossed semimodule $\mathcal{C} = (A, B, \partial)$ consists of a pair of monoids A, B and a homomorphism $\partial: A \rightarrow B$ of monoids with an action $\bullet: B \times A \rightarrow A$ of B on A satisfying

$$[\text{CSM 1}] \quad \partial(b \bullet a)b = b\partial(a)$$

$$[\text{CSM } 2] \quad (\partial(a) \bullet a_1)a = aa_1,$$

for $a, a_1 \in A$ and $b \in B$.

Let $\mathcal{C} = (A, B, \partial)$ and $\mathcal{C}' = (A', B', \partial')$ be crossed semimodules. A crossed semimodule morphism is a mapping $\lambda = (\lambda_1, \lambda_2): \mathcal{C} \rightarrow \mathcal{C}'$ where $\lambda_1: A \rightarrow A'$ and $\lambda_2: B \rightarrow B'$ are monoid homomorphisms such that $\lambda_1(b \bullet a) = \lambda_2(b) \bullet' \lambda_1(a)$ and $\lambda_2\partial = \partial'\lambda_1$.

$$\begin{array}{ccccc} B \times A & \xrightarrow{\bullet} & A & \xrightarrow{\partial} & B \\ \lambda_2 \times \lambda_1 \downarrow & & \lambda_1 \downarrow & & \downarrow \lambda_2 \\ B' \times A' & \xrightarrow{\bullet'} & A' & \xrightarrow{\partial'} & B' \end{array}$$

Hence crossed semimodules and their morphisms form a category which we denoted by CSM.

The following theorem and corollary are proved in [24]:

THEOREM 2.2. *The category SIC of Schreier internal categories in MON is equivalent to the category CSM of crossed semimodules.*

Note that this theorem is obtained as a special case of the equivalence given in [29].

Let CSM* denote the category of crossed semimodules $\mathcal{C} = (A, B, \partial)$ such that A is a group. Then we obtain the following corollary as a restriction of above equivalence.

COROLLARY 2.3. *The category SIG of Schreier internal groupoids in MON is equivalent to the category CSM*.*

Note that this corollary is obtained as a special case of the equivalence given in [26]. Restricting this corollary to the case of groups, Theorem 2.1 is obtained as given in [9] and [24].

Let G be a group. We recall that from [10] a cat^1 -group (or 1-cat-group [14]) is a triple $\mathcal{G} = (G, s, t)$ with two group homomorphisms $s, t: G \rightarrow G$ called structural maps satisfying following conditions

$$[\text{C1G } 1] \quad st = t \text{ and } ts = s$$

$$[\text{C1G } 2] \quad [\text{Ker } s, \text{Ker } t] = 1 \text{ where } [\text{Ker } s, \text{Ker } t] \text{ is the commutator subgroup of Ker } s \text{ and Ker } t.$$

A cat^1 -group morphism is a group homomorphism $f: (G, s, t) \rightarrow (G', s', t')$ where $\mathcal{G} = (G, s, t)$ and $\mathcal{G}' = (G', s', t')$ are cat^1 -groups such that $s'f = fs$ and $t'f = ft$.

$$\begin{array}{ccc} G & \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} & G \\ f \downarrow & & \downarrow f \\ G' & \begin{array}{c} \xrightarrow{s'} \\ \xrightarrow{t'} \end{array} & G' \end{array}$$

Hence cat^1 -groups form a category which we denoted by $\text{CAT}^1\text{-GP}$. For the categorical equivalence between crossed modules over groups and cat^1 -groups, see [10].

3. Cat^1 -monoids

DEFINITION 3.1. A cat^1 -monoid is a triple $\mathcal{M} = (M, s, t)$ where M is a monoid and $s, t: M \rightarrow M$ are monoid homomorphisms of M satisfying

- [C1M 1] $st = t$ and $ts = s$,
- [C1M 2] $xy = yx$ for $x \in \text{Ker } s, y \in \text{Ker } t$,
- [C1M 3] $\text{Ker } s$ is a group and
- [C1M 4] there exists a unique $\tilde{m} \in \text{Ker } s$ for all $m \in M$ such that $m = \tilde{m}s(m)$.

EXAMPLE 3.2. Consider the monoid (\mathbb{Z}_4, \cdot) and the group (\mathbb{Z}_5, \cdot) . Then we construct a cat^1 -monoid $(\mathbb{Z}_4 \times \mathbb{Z}_5, s, t)$ with direct product where $s(\bar{x}^4, \bar{y}^5) = (\bar{x}^4, \bar{1}^5)$ and $t(\bar{x}^4, \bar{y}^5) = (\bar{y} \cdot \bar{x}^4, \bar{1}^5)$. Clearly $\text{Ker } s$ is a group. Since $(\bar{x}^4, \bar{y}^5) = (\bar{1}^4, \bar{y}^5) \cdot (\bar{x}^4, \bar{1}^5)$, all pairs satisfy the condition [C1M4].

EXAMPLE 3.3. Every cat^1 -group is a cat^1 -monoid. Note that $\tilde{m} = ms(m^{-1})$.

PROPOSITION 3.4. Given any cat^1 -monoid (M, s, t) , we have

1. $\text{Im } s = \text{Im } t$,
2. s and t are identities on $\text{Im } s$,
3. $s^2 = s$ and $t^2 = t$.

Proof. The proof is straightforward. So it is omitted. □

DEFINITION 3.5. A morphism $f: (M, s, t) \rightarrow (M', s', t')$ of cat^1 -monoids is a morphism of monoids such that $s'f = fs$ and $t'f = ft$.

$$\begin{array}{ccc} M & \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} & M \\ f \downarrow & & \downarrow f \\ M' & \begin{array}{c} \xrightarrow{s'} \\ \xrightarrow{t'} \end{array} & M' \end{array}$$

Hence we can construct the category of cat^1 -monoids which we denoted by $\text{CAT}^1\text{-MON}$.

PROPOSITION 3.6. Let $\mathcal{C} = (A, B, \partial)$ be an object of CSM^* . Then $\mathcal{M} = (M, s, t)$ is a cat^1 -monoid where $M = B \times A$ is the semi-direct product of monoids, $s(b, a) = (b, 1)$ and $t(b, a) = (\partial(a)b, 1)$.

Proof. We define a functor $\psi: \text{CSM}^* \rightarrow \text{CAT}^1\text{-MON}$ such that $\psi(A, B, \partial) = (M, s, t)$. The product of M is defined by

$$(b, a)(b', a') = (bb', a(b \bullet a'))$$

for $a, a' \in A$, $b, b' \in B$. Clearly s is a monoid homomorphism. We prove that t is a homomorphism of monoids.

$$\begin{aligned} t\left((b, a)(b', a')\right) &= t\left(bb', a(b \bullet a')\right) \\ &= \left(\partial(a)\partial(b \bullet a')bb', 1\right) \\ &= (\partial(a)b\partial(a')b', 1) \\ &= (\partial(a)b, 1)(\partial(a')b', 1) \\ &= t(b, a)t(b', a') \end{aligned}$$

It is easy to show that s and t satisfy [C1M1]. With respect to [C1M2], for elements $(1, a) \in \text{Ker } s$ and $(b', a') \in \text{Ker } t$, we have

$$\begin{aligned} (b', a')(1, a) &= (b', a'(b' \bullet a)) \\ &= (b', (\partial(a') \bullet (b' \bullet a))a') \\ &= (b', (\partial(a')b' \bullet a)a') \\ &= (b', (1 \bullet a)a') \\ &= (b', aa') \\ &= (b', a(1 \bullet a')) \\ &= (1, a)(b', a'). \end{aligned}$$

Clearly $\text{Ker } s$ is a group. Since $(b, a) = (1, a)(b, 1) = (1, a)s(b, a)$, that is $\widetilde{(b, a)} = (1, a)$ and so each element of M satisfies [C1M4]. \square

PROPOSITION 3.7. *Let $\mathcal{M} = (M, s, t)$ be a cat^1 -monoid. Then $\gamma(M, s, t) = (A, B, \partial)$ is an object of CSM^* where $A = \text{Ker } s$, $B = \text{Im } s$, $\partial = t|_{\text{Ker } s}$ and an action of B on A is defined by $(n \bullet x)n = nx$, for $x \in \text{Ker } s$ and $n \in \text{Im } s$.*

Proof. We define a functor

$$\gamma: \text{CAT}^1\text{-MON} \rightarrow \text{CSM}^*$$

as a weak inverse of ψ such that $\gamma(M, s, t) = (A, B, \partial)$. First we show that \bullet is an action of $\text{Im } s$ on $\text{Ker } s$. Let $n, n' \in \text{Im } s$ and $x, x' \in \text{Ker } s$. Since

$$(nn' \bullet x)nn' = nn'x = n(n' \bullet x)n' = (n \bullet (n' \bullet x))nn',$$

under the condition [C1M4] we write

$$(nn') \bullet x = n \bullet (n' \bullet x).$$

Since

$$(n \bullet (xx'))n = nxx' = (n \bullet x)nx' = (n \bullet x)(n \bullet x')n,$$

under the condition [C1M4]

$$n \bullet (xx') = (n \bullet x)(n \bullet x').$$

Clearly

$$1 \bullet x = (1 \bullet x)1 = 1x = x.$$

Now we will verify [CSM1] and [CSM2]. Since $n = t(n) = s(n)$, by Proposition 3.4. we get

$$t(n \bullet x)n = t(n \bullet x)t(n) = t((n \bullet x)n) = t(nx) = t(n)t(x) = nt(x).$$

On the other hand

$$(t(x) \bullet x')x = (t(x) \bullet x')t(x)t(x^{-1})x = t(x)x't(x^{-1})x.$$

Since $t(x^{-1})x \in \text{Ker } t$ and $x' \in \text{Ker } s$, by [C1M2] we have

$$(t(x) \bullet x')x = t(x)x't(x^{-1})x = t(x)t(x^{-1})xx' = xx'.$$

□

As a corollary of Propositions 3.6 and 3.7 we can give the following theorem.

THEOREM 3.8. *The categories $\text{CAT}^1\text{-MON}$ and CSM^* are equivalent.*

Proof. Let $f: (M, s, t) \rightarrow (M', s', t')$ be a morphism of cat^1 -monoids, $n \in \text{Im } s, x \in \text{Ker } s$. Since $f(n) \in N'$, the condition [C1M4] allows us to write

$$f(n \bullet x)f(n) = f((n \bullet x)n) = f(nx) = f(n)f(x) = \left(f(n) \bullet f(x) \right) f(n)$$

and so under the same condition we have

$$f(n \bullet x) = f(n) \bullet f(x).$$

This means that $\gamma(f) = (f, f)$ is a morphism of crossed semimodules.

On the other hand, given a morphism $\lambda = (\lambda_1, \lambda_2): \mathcal{C} \rightarrow \mathcal{C}'$ where $\mathcal{C} = (A, B, \partial)$ and $\mathcal{C}' = (A', B', \partial')$ are crossed semimodules such that A, A' are groups, $\psi(\lambda_1, \lambda_2) = (\lambda_2, \lambda_1)$ is a morphism of cat^1 -monoids as shown in the following diagram.

$$\begin{array}{ccc} B \times A & \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} & B \times A \\ (\lambda_2, \lambda_1) \downarrow & & \downarrow (\lambda_2, \lambda_1) \\ B' \times A' & \begin{array}{c} \xrightarrow{s'} \\ \xrightarrow{t'} \end{array} & B' \times A' \end{array}$$

We will verify that this diagram is commutative.

$$\begin{aligned}
(\lambda_2, \lambda_1)s(b, a) &= (\lambda_2, \lambda_1)(b, 1) \\
&= (\lambda_2(b), 1) \\
&= s'(\lambda_2(b), \lambda_1(a)) \\
&= s'(\lambda_2, \lambda_1)(b, a)
\end{aligned}$$

$$\begin{aligned}
(\lambda_2, \lambda_1)t(b, a) &= (\lambda_2, \lambda_1)(\partial(a)b, 1) \\
&= (\lambda_2\partial(a)\lambda_2(b), 1) \\
&= (\partial\lambda_1(a)\lambda_2(b), 1) \\
&= t'(\lambda_2(b), \lambda_1(a)) \\
&= t'(\lambda_2, \lambda_1)(b, a)
\end{aligned}$$

A natural equivalence $S: 1_{\text{CAT}^1\text{-MON}} \rightarrow \psi\gamma$ is given via a mapping

$$S(M, s, t) = (\text{Im } s \times \text{Ker } s, s', t')$$

such that $S_M(m) = (s(m), \tilde{m})$ and $s'(n, x) = (n, 1)$, $t'(n, x) = (t(x)n, 1)$ is an isomorphism. We will verify that S_M is a homomorphism.

$$\begin{aligned}
S_M(mm_1) &= S_M\left(\tilde{m}s(m)\tilde{m}_1s(m_1)\right) \\
&= S_M\left(\tilde{m}(s(m) \bullet \tilde{m}_1)s(m)s(m_1)\right) \\
&= S_M\left(\tilde{m}(s(m) \bullet \tilde{m}_1)s(mm_1)\right) \\
&= \left(s(mm_1), \tilde{m}(s(m) \bullet \tilde{m}_1)\right) \\
&= \left(s(m), \tilde{m}\right)\left(s(m_1), \tilde{m}_1\right) \\
&= S_M(m)S_M(m_1)
\end{aligned}$$

On the other hand, a natural equivalence $T: 1_{\text{CSM}^*} \rightarrow \gamma\psi$ is defined by

$$T(\mathcal{C}) = (\text{Ker } s, s(B \times A), t)$$

for $\mathcal{C} = (A, B, \partial)$ such that $T_{\mathcal{C}}(b) = (b, 1)$, $T_{\mathcal{C}}(a) = (1, a)$.

Other details are straightforward and so is omitted. \square

Since every group is a monoid then the following equivalence given in [10, 14] is a consequence of Theorem 3.8.

THEOREM 3.9. [10] *The category of crossed modules over groups is equivalent to the category of cat^1 -groups.*

References

- [1] Akız, H.F., Alemdar, N., Mucuk, O. and Şahan, T., *Coverings of internal groupoids and crossed modules in the category of groups with operations*, Georgian Mathematical Journal **20** (2) (2013), 223–238.
- [2] Baez, J.C. and Lauda, A.D., *Higher Dimensional Algebra V: 2-Groups*, Theory and Applications of Categories **12** (14) (2004), 423–491.
- [3] Baez J.C. and Stevenson D. *The classifying space of a topological 2-group*, Algebraic Topology Abel Symposia; 4: pp 1-31 (2009).
- [4] Baez, J.C., Baratin, A., Freidel, L. and Wise, D.K., *Infinite-Dimensional Representations of 2-Groups*, Memoirs of the American Mathematical Society, Volume 219, Number 1032, (2012).
- [5] Brown, R., *Groupoids and crossed objects in algebraic topology*, Homology Homotopy Appl. **1** (1999), 1–78.
- [6] Brown, R., *Higher dimensional group theory*. In: *Low Dimensional Topology*, London Math. Soc. Lect. Notes, 48, pp. 215–238. Cambridge Univ. Press, (1982)
- [7] Brown, R. and Huebschmann, J., *Identities among relations*. In: *Low Dimensional Topology*, London Math. Soc. Lect. Notes, 48, pp. 153–202. Cambridge Univ. Press (1982).
- [8] Brown, R. and Mucuk, O., *Covering groups of non-connected topological groups revisited*, Math. Proc. Camb. Phil. Soc. **115** (1994), 97–110.
- [9] Brown, R. and Spencer, C.B., *\mathcal{G} -groupoids, crossed modules and the fundamental groupoid of a topological group*, Proc. Konn. Ned. Akad. **79** (1976), 296–302.
- [10] Brown, R., Higgins, P. J. and Sivera, R., *Nonabelian Algebraic Topology: Filtered spaces, crossed complexes, cubical homotopy groupoids*, European Mathematical Society Tracts in Mathematics **15** (2011).
- [11] Brown, R. and Huebschmann, J., *Identities among relations*, pages 153–202. London Mathematical Society Lecture Note Series. Cambridge University Press, (1982).
- [12] Huebschmann, J., *Crossed n -fold extensions of groups and cohomology*, Comment. Math. Helvetici **55** (1980), 302–314.
- [13] Loday, J.-L., *Cohomologie et groupes de Steinberg relatifs*, J. Algebra **54** (1978), 178–202.
- [14] Loday, J.-L., *Spaces with finitely many non-trivial homotopy groups*, J.Pure Appl. Algebra **24** (2) (1982), 179–202 .
- [15] Lue, A.S.T., *Cohomology of groups relative to a variety*, J. Algebra **69** (1981), 155–174.
- [16] MacLane, S., *Categories for the Working Mathematician, Graduate Text in Mathematics*, Volume 5. Springer-Verlag, New York (1971).

- [17] Mucuk, O., Şahan, T. and Alemdar, N., *Normality and quotients in crossed modules and group-groupoids*, Applied Categorical Structures **23** (3) (2015), 415–428.
- [18] Mucuk, O., Kılıçarslan, B., Şahan T. and Alemdar N., *Group-groupoids and monodromy groupoids*, Topology Appl. **158** (15) (2011), 2034–2042.
- [19] Mucuk, O. and Şahan, T., *Coverings and crossed modules of topological groups with operations*, Turkish Journal of Mathematics **38** (5), 833–845.
- [20] Mucuk, O. and Şahan, T., *Group-groupoid actions and liftings of crossed modules*, Georgian Mathematical Journal, Doi: 10.1515/gmj-2018-0001.
- [21] Mucuk, O. and Akız, H. F., *Monodromy groupoids of an internal groupoid in topological groups with operations*, Filomat **29**:10 (2015), 2355–2366.
- [22] N. Martins-Ferreira, A. Montoli and M. Sobral., *Semidirect products and crossed modules in monoids with operations*, J. Pure Appl. Algebra **217** (2013), 334–347.
- [23] Orzech, G., *Obstruction theory in algebraic categories I, II.*, J. Pure Appl. Algebra **2** (1972), 287–314, 315–340.
- [24] Patchkoria, A., *Crossed Semimodules and Schreier Internal Categories In The Category of Monoids*, Georgian Mathematical Journal **5** (6) (1998), 575–581.
- [25] Porter, T., *Extensions, crossed modules and internal categories in categories of groups with operations*, Proc. Edinb. Math. Soc. **30** (1987) , 373–381.
- [26] Porter, T., *Crossed Modules in Cat and a Brown-Spencer Theorem for 2-Categories*, Cahiers de Topologie et Geometrie Differentielle Categoriques, Vol. XXVI-4 (1985).
- [27] Şahan, T., *Further remarks on liftings of crossed modules*, Hacettepe J. Math. Stat. **48** (3) (2019), 743–752. Doi: 10.15672/HJMS.2018.554.
- [28] Temel, S., *Topological Crossed Semimodules and Schreier Internal Categories in the Category of Topological Monoids*, Gazi University Journal of Science **29** (4) (2016), 915–921 .
- [29] Temel, S., *Crossed semimodules of categories and Schreier 2-categories*, Tbilisi Math. J. **11** (2) (2018), 47–57.
- [30] Whitehead, J.H.C., *Combinatorial homotopy II*, Bull. Amer. Math. Soc. **55** (1949), 453–496.
- [31] Whitehead, J.H.C., *Note on a previous paper entitled "On adding relations to homotopy group"*, Ann. Math. **47** (1946), 806–810.

Sedat Temel

Department of Mathematics

Recep Tayyip Erdogan University, Rize, Turkey

E-mail: sedat.temel@erdogan.edu.tr