

FIXED POINT THEOREMS FOR ASYMPTOTICALLY REGULAR MAPPINGS IN FUZZY METRIC SPACES

NILAKSHI GOSWAMI AND BIJOY PATIR*

ABSTRACT. The aim of this paper is to extend some existing fixed point results for asymptotically regular mappings to fuzzy metric spaces. For this purpose some contractive type conditions with respect to an altering distance function are used. Some new common fixed point results have been derived for such mappings. We provide suitable examples to justify our study.

1. Introduction and preliminaries

In 1966, Browder and Petryshyn introduced the concept of asymptotic regularity of self mapping at a point in a metric space [1]. After that several researchers (refer to [6, 8, 12, 13, 15]) derived different results in fixed point theory for such type of mappings.

After the initiation of the study of fixed point theory in fuzzy metric spaces by Grabiec in 1988 [3], extensions of many existing fixed point results as well as new fixed point results have been established by several research workers in fuzzy metric spaces (refer to [8, 10, 11, 14–18]). The introduction of the concept of altering distance function by Khan et al. [5] enhanced the fixed point theory in different spaces.

Received April 5, 2019. Revised September 17, 2019. Accepted September 19, 2019.

2010 Mathematics Subject Classification: 47H10, 54H25.

Key words and phrases: fuzzy metric space, fixed point, asymptotically regular mappings.

* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2019.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

In this paper, we prove some fixed point theorems for asymptotically regular mappings in fuzzy metric spaces which extend some existing fixed point results. Moreover using the concept of compatibility, we derive some common fixed point results for asymptotically regular mappings.

First we provide some basic concepts.

DEFINITION 1.1. [4] A mapping $T : [0, 1] \times [0, 1] \longrightarrow [0, 1]$ is called a triangular norm or t-norm if

- i) $T(x_1, 1) = x_1 \forall x_1 \in [0, 1]$,
- ii) $T(x_1, x_2) = T(x_2, x_1) \forall x_1, x_2 \in [0, 1]$,
- iii) $x_1 \geq x_2, x_3 \geq x_4 \Rightarrow T(x_1, x_3) \geq T(x_2, x_4)$,
 $\forall x_1, x_2, x_3, x_4 \in [0, 1]$,
- iv) $T(x_1, T(x_2, x_3)) = T(T(x_1, x_2), x_3), \forall x_1, x_2, x_3 \in [0, 1]$.

Some basic examples of t-norms are $T_m(x_1, x_2) = \min(x_1, x_2)$, $T_p(x_1, x_2) = x_1 \cdot x_2$, $T_L(x_1, x_2) = \max(x_1 + x_2 - 1, 0)$.

DEFINITION 1.2. [2] For an arbitrary set X , let T be a continuous t-norm and M be a fuzzy set on $X \times X \times (0, \infty)$. The 3-tuple (X, M, T) is called a fuzzy metric space if the following conditions are satisfied :

- a) $M(x_1, x_2, t) > 0, \forall x_1, x_2 \in X, t > 0$,
- b) $M(x_1, x_2, t) = 1 \Leftrightarrow x_1 = x_2, \forall t > 0$,
- c) $M(x_1, x_2, t) = M(x_2, x_1, t), \forall x_1, x_2 \in X, t > 0$,
- d) $T(M(x_1, x_2, t), M(x_2, x_3, s)) \leq M(x_1, x_3, t + s)$,
 $\forall x_1, x_2, x_3 \in X, t, s > 0$,
- e) $M(x_1, x_2, \cdot) : (0, \infty) \longrightarrow [0, 1]$ is continuous $\forall x_1, x_2 \in X$.

For a fuzzy metric space (X, M, T) , the function M is a continuous function on $X \times X \times (0, \infty)$ (refer to [7]).

EXAMPLE 1.3. Let $X = [0, 1]$. For a continuous function $g : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ and $t \in (0, \infty)$, let $M(x, y, t) = e^{\left(\frac{-d(x,y)}{g(t)}\right)}$ where $d(x, y) = |x - y|$. Then (X, M, T) is a fuzzy metric space with respect to the t-norm $T_p(x, y) = x \cdot y, x, y \in X$ (refer to [4]).

DEFINITION 1.4. [3] Let (X, M, T) be a fuzzy metric space and $\{x_n\}$ be a sequence in X .

$\{x_n\}$ is a Cauchy sequence in X if $\forall \varepsilon \in (0, 1), \exists n_0 \in \mathbb{N}$ such that $M(x_n, x_m, t) > 1 - \varepsilon, \forall n, m \geq n_0$, or equivalently, if $\lim_{n, m \rightarrow \infty} M(x_n, x_m, t) = 1, \forall t > 0$.

$\{x_n\}$ converges to x if $\forall \varepsilon \in (0, 1), \exists n_0 \in \mathbb{N}$ such that $M(x_n, x, t) > 1 - \varepsilon, \forall n \geq n_0$.

X is said to be complete if and only if every Cauchy sequence converges in X .

DEFINITION 1.5. [14] A mapping $\phi : [0, 1] \rightarrow [0, 1]$ is called an altering distance function if

(i) ϕ is strictly decreasing and left continuous.

(ii) $\phi(\lambda) = 0$ if and only if $\lambda = 1$

i.e, $\lim_{\phi \rightarrow 1^-} \phi(1) = 0$.

DEFINITION 1.6. [9, 13] Let f and g be self mappings on a fuzzy metric space (X, M, T) and $\{x_n\}$ be a sequence in X .

f is said to be asymptotically regular at a point $x_0 \in X$ if $\lim_{n \rightarrow \infty} M(f^n(x_0), f^{n+1}(x_0), t) = 1, \forall t > 0$.

Also the sequence $\{x_n\}$ is said to be asymptotically regular with respect to the pair (f, g) if $\lim_{n \rightarrow \infty} M(f(x_n), g(x_n), t) = 1, \forall t > 0$.

DEFINITION 1.7. [9] Two self mappings f and g on a fuzzy metric space (X, M, T) are said to be compatible if $\lim_{n \rightarrow \infty} M(fg(x_n), gf(x_n), t) = 1, \forall t > 0$, where $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} g(x_n) = x$, for some $x \in X$.

2. Results and discussion

In 2013, Shukla et al. [15] proved a fixed point result in a complete partial metric space using asymptotic regularity. Following is an extension of this result in the setting of fuzzy metric space.

THEOREM 2.1. Let (X, M, T) be a complete fuzzy metric space, ϕ be the altering distance function and $f : X \rightarrow X$ be such that the following condition is satisfied:

$$\begin{aligned} \phi(M(f(x), f(y), t)) &\leq b_1(x, y)\theta[\min\{\phi(M(x, f(x), t)), \phi(M(y, f(y), t))\}] \\ &\quad + b_2(x, y)\psi[\phi(M(x, f(x), t)) \cdot \phi(M(y, f(y), t))] + b_3(x, y)\phi(M(x, y, t)) \\ &\quad + b_4(x, y)(\phi(M(x, f(x), t)) + \phi(M(y, f(y), t))) \\ (1) \quad &\quad + b_5(x, y)[\phi(M(x, f(y), t)) + \phi(M(f(x), y, t))] \end{aligned}$$

$\forall x, y \in X, t > 0$ where $b_i : X \times X \rightarrow [0, \infty), i = 1, 2, 3, 4, 5$ are such that for some arbitrarily fixed $\lambda_1 > 0, 0 < \lambda_2 < 1$,

$$(2) \quad b_1(x, y) + b_2(x, y) \leq \lambda_1,$$

$$(3) \quad b_3(x, y) + b_4(x, y) + 2b_5(x, y) \leq \lambda_2$$

and $\theta, \psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are continuous functions at 0 and $\theta(0) = \psi(0) = 0$. If f is asymptotically regular at some point $x_0 \in X$, then f has a unique fixed point in X .

Proof. Suppose that $\{x_n\}$ is a sequence in X where $x_0 \in X$ and $x_{n+1} = f(x_n) \forall n \geq 0$. Now if for some $n \geq 0$, $x_n = x_{n+1}$, then x_n is a fixed point of f . Suppose that $x_n \neq x_{n+1} \forall n$. We show that the sequence $\{x_n\}$ is Cauchy.

Suppose to the contrary $\exists 0 < \varepsilon < 1, t > 0$ and two sequences of integers $\{r_n\}$ and $\{s_n\}$ such that $r_n > s_n > n$,

$$(4) \quad \begin{aligned} M(x_{r_n}, x_{s_n}, t) &\leq 1 - \varepsilon, \\ M(x_{r_{n-1}}, x_{s_{n-1}}, t) &> 1 - \varepsilon, \\ M(x_{r_{n-1}}, x_{s_n}, t) &> 1 - \varepsilon, \quad \forall n \in \mathbb{N} \cup \{0\}. \end{aligned}$$

Now we have

$$\begin{aligned} 1 - \varepsilon &\geq M(x_{r_n}, x_{s_n}, t) \geq T\left(M(x_{r_n}, x_{r_{n-1}}, \frac{t}{2}), M(x_{r_{n-1}}, x_{s_n}, \frac{t}{2})\right) \\ &\geq T\left(M(x_{r_n}, x_{r_{n-1}}, t), 1 - \varepsilon\right) \\ \Rightarrow 1 - \varepsilon &\geq \lim_{n \rightarrow \infty} M(x_{r_n}, x_{s_n}, t) \geq T(1, 1 - \varepsilon) \\ &\quad \text{(since } f \text{ asymptotically regular at } x_0) \\ (5) \Rightarrow \lim_{n \rightarrow \infty} M(x_{r_n}, x_{s_n}, t) &= 1 - \varepsilon \end{aligned}$$

Again,

$$(6) \quad \begin{aligned} M(x_{r_n}, x_{s_{n-1}}, t) &\geq T\left(M(x_{r_n}, x_{s_n}, \frac{t}{2}), M(x_{s_n}, x_{s_{n-1}}, \frac{t}{2})\right) \\ \Rightarrow \lim_{n \rightarrow \infty} M(x_{r_n}, x_{s_{n-1}}, t) &> T(1 - \varepsilon, 1) \\ \Rightarrow \lim_{n \rightarrow \infty} M(x_{r_n}, x_{s_{n-1}}, t) &> 1 - \varepsilon \end{aligned}$$

Taking $x = x_{r_{n-1}}$ and $y = x_{s_{n-1}}$ in (1), we have

$$\begin{aligned} \phi(M(x_{r_n}, x_{s_n}, t)) &\leq b_1(x, y)\theta(\min\{\phi(M(x_{r_{n-1}}, x_{r_n}, t)), \phi(M(x_{s_{n-1}}, x_{s_n}, t))\}) \\ &\quad + b_2(x, y)\psi(\phi(M(x_{r_{n-1}}, x_{r_n}, t)) \cdot \phi(M(x_{s_{n-1}}, x_{s_n}, t))) \\ &\quad + b_3(x, y)\phi(M(x_{r_{n-1}}, x_{s_{n-1}}, t)) + b_4(x, y)[\phi(M(x_{r_{n-1}}, x_{r_n}, t)) \\ &\quad + \phi(M(x_{s_{n-1}}, x_{s_n}, t))] + b_5(x, y)[\phi(M(x_{r_{n-1}}, x_{s_n}, t)) \\ &\quad + \phi(M(x_{r_n}, x_{s_{n-1}}, t))] \end{aligned}$$

Taking $n \rightarrow \infty$ and by (4), (5), (6) and using the fact that f is asymptotically regular at x_0 we have,

$$\phi(1 - \varepsilon) \leq b_3(x, y)\phi(1 - \varepsilon) + 2b_5(x, y)\phi(1 - \varepsilon) < \phi(1 - \varepsilon)$$

which is a contradiction.

Thus $\{x_n\}$ is a Cauchy sequence. Since (X, M, T) is a complete fuzzy metric space, $\exists z \in X$ such that $x_n \rightarrow z$.

$$\begin{aligned} \text{Now, } & \phi(M(f(x_n), f(z), t)) \\ & \leq b_1(x, y)\theta(\min\{\phi(M(x_n, x_{n+1}, t)), \phi(M(z, f(z), t))\}) \\ & + b_2(x, y)\psi(\phi(M(x_n, x_{n+1}, t)) \cdot \phi(M(z, f(z), t))) \\ & + b_3(x, y)\phi(M(x_n, z, t)) + b_4(x, y)[\phi(M(x_n, x_{n+1}, t)) \\ & + \phi(M(z, f(z), t))] + b_5(x, y)[\phi(M(x_n, f(z), t)) \\ & + \phi(M(z, x_{n+1}, t))] \end{aligned}$$

For $n \rightarrow \infty$,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \phi(M(z, f(z), t)) \leq (b_4(x, y) + b_5(x, y)) \lim_{n \rightarrow \infty} \phi(M(z, f(z), t)) \\ \Rightarrow & (1 - b_4(x, y) - b_5(x, y)) \lim_{n \rightarrow \infty} \phi(M(z, f(z), t)) \leq 0 \\ \Rightarrow & \lim_{n \rightarrow \infty} \phi(M(z, f(z), t)) = 0 \quad (\text{since } 0 < b_3(x, y) + b_4(x, y) + 2b_5(x, y) < 1) \\ \Rightarrow & f(z) = z. \end{aligned}$$

If u is another fixed point of f in X , then

$$\begin{aligned} & \phi(M(f(u), f(z), t)) \\ & \leq b_1(x, y)\theta(\min\{\phi(M(u, f(u), t)), \phi(M(z, f(z), t))\}) \\ & + b_2(x, y)\psi(\phi(M(u, f(u), t)) \cdot \phi(M(z, f(z), t))) \\ & + b_3(x, y)\phi(M(u, z, t)) + b_4(x, y)[\phi(M(u, f(u), t)) \\ & + \phi(M(z, f(z), t))] + b_5(x, y)[\phi(M(u, f(z), t)) \\ & + \phi(M(z, f(u), t))] \\ \Rightarrow & \phi(M(u, z, t)) \leq b_3(x, y)\phi(M(u, z, t)) + 2b_5(x, y)\phi(M(u, z, t)) \\ \Rightarrow & (1 - b_3(x, y) - 2b_5(x, y))\phi(M(u, z, t)) \leq 0 \\ \Rightarrow & \phi(M(u, z, t)) = 0 \quad (\text{since } 0 < b_3(x, y) + b_4(x, y) + 2b_5(x, y) < 1) \\ \Rightarrow & u = z, \end{aligned}$$

establishes that the fixed point of f in X is unique. \square

EXAMPLE 2.2. Consider the complete metric space (X, d) where $X = [0, 1]$ and $d(x, y) = |x - y| \forall x, y \in X$. Let M be a fuzzy set on $X \times X \times (0, \infty)$ given by $M(x, y, t) = \frac{t}{t + d(x, y)}$, if $t > 0$.

Then (X, M, T) is a complete fuzzy metric space with respect to the t-norm $T(a, b) = \min\{a, b\}$, $a, b \in [0, 1]$. Define $f : X \rightarrow X$ by

$$f(x) = \begin{cases} \frac{x}{3}, & x \in [0, \frac{1}{2}] \\ \frac{1}{6}, & x \in [\frac{1}{2}, 1] \end{cases} \quad \text{and } \phi(\lambda) = 1 - \lambda, \lambda \in [0, 1].$$

Let $\theta(p) = \sqrt{p}$ and $\psi(q) = q^2$, $\forall p, q \in \mathbb{R}^+$.

Also let $b_1(x, y) = |x - y|$, $b_2(x, y) = |x^2 - y^2|$, $b_3(x, y) = \begin{cases} \frac{1}{|x - y|}, & x \neq y \\ 0, & x = y \end{cases}$,

$$b_4(x, y) = b_5(x, y) = \frac{1 - b_3(x, y)}{3}.$$

Then f satisfies (1).

Hence from Theorem 2.1 we can say that f has a unique fixed point.

COROLLARY 2.3. Let $f, g : X \rightarrow X$ be mappings on a complete fuzzy metric space (X, M, T) and ϕ be the altering distance function. Let f and g be asymptotically regular at a point $x_0 \in X$ and both satisfy the inequality (1). Moreover, if

$$(7) \quad \phi(M(f(x), g(y), t)) \leq k(\phi(M(x, y, t)) + \phi(M(x, f(x), t)) + \phi(M(y, g(y), t))),$$

where $0 < k < 1$ and $x, y \in X$,

then f and g have a unique common fixed point in X .

Proof. From Theorem 2.1, both f and g have unique fixed points say, z and v respectively.

Since f and g satisfy (7),

$$\begin{aligned} & \phi(M(f(z), g(v), t)) \leq k(\phi(M(z, v, t)) + \phi(M(z, f(z), t)) + \phi(M(v, g(v), t))) \\ \Rightarrow & \phi(M(z, v, t)) \leq k(\phi(M(z, v, t)) + \phi(M(z, z, t)) + \phi(M(v, v, t))) \\ \Rightarrow & (1 - k)\phi(M(z, v, t)) \leq 0 \\ \Rightarrow & \phi(M(z, v, t)) \leq 0 \quad (\text{since } k < 1) \\ \Rightarrow & z = v \end{aligned}$$

i.e., f and g have a unique common fixed point. \square

EXAMPLE 2.4. Let $X = \{P, Q, R, S\} \subseteq \mathbb{R}^2$, where $P = (0, 0)$, $Q = (0, 1)$, $R = (1, 0)$ and $S = (1, 1)$. Let $M(x, y, t) = \frac{t}{t + d(x, y)}$, $t > 0$, where

$d(x, y)$ is the Euclidean distance in \mathbb{R}^2 . Let $\phi(\lambda) = 1 - \lambda, \lambda \in [0, 1]$. Then (X, M, T) is a complete fuzzy metric space with respect to the t-norm $T(a, b) = \min\{a, b\}$.

Let $f, g : X \rightarrow X$ be defined by

$$f(P) = f(Q) = P, f(R) = Q, f(S) = R, \text{ and}$$

$$g(P) = g(R) = P, g(Q) = R, g(S) = Q.$$

we can be shown easily that, for $k = \frac{4}{5}$ every pair of points in X satisfies the condition (7).

Hence by Theorem 2.3, f and g have a unique common fixed point which is P in this case.

In the following result, we use a minimum condition.

THEOREM 2.5. *Let $f : X \rightarrow X$ be a mapping on a complete fuzzy metric space (X, M, T) and ϕ is the altering distance function. If f is asymptotically regular at a point $x_0 \in X$ and f satisfies,*

$$(8) \quad \phi(M(f(x), f(y), t)) \leq h_1 \min\{\phi(M(x, y, t)), \phi(M(f(x), x, t)), \phi(M(f(x), y, t))\} \\ + h_2 \min\{\phi(M(x, y, t)), \phi(M(f(y), y, t)), \phi(M(x, f(y), t))\}$$

for all $x, y \in X, t > 0$, where $h_1, h_2 > 0$ are constants such that $h_1 + h_2 < 1$,

then f has a unique fixed point in X .

Proof. As in Theorem 2.1, we construct a sequence $\{x_n\}$ in X by $x_{n+1} = f(x_n) \forall n \in \mathbb{N} \cup \{0\}$ where $x_0 \in X$. If there exists n with $x_n = x_{n+1}$, then x_n is a fixed point of f . Suppose that $x_n \neq x_{n+1}$ for all n .

To show that $\{x_n\}$ is a Cauchy sequence.

Let $m, n \in \mathbb{N} \cup \{0\}$. From (8),

$$\phi(M(f(x_n), f(x_m), t)) \leq h_1 \min\{\phi(M(x_n, x_m, t)), \phi(M(f(x_n), x_n, t)), \\ \phi(M(f(x_n), x_m, t))\} + h_2 \min\{\phi(M(x_n, x_m, t)), \phi(M(f(x_m), x_m, t)), \\ \phi(M(x_n, f(x_m), t))\}.$$

Since, f is asymptotically regular at $x_0 \in X$, taking $n, m \rightarrow \infty$,

$$\lim_{n, m \rightarrow \infty} \phi(M(f(x_n), f(x_m), t)) = 0 \\ \Rightarrow \lim_{n, m \rightarrow \infty} M(f(x_n), f(x_m), t) = 1,$$

i.e. $\{x_n\}$ is a Cauchy sequence in (X, M, T) . Since (X, M, T) is complete, therefore $x_n \rightarrow z$ (say) in X .

Using (8),

$$\begin{aligned}
 \phi(M(x_{n+1}, f(z), t)) &= \phi(M(f(x_n), f(z), t)) \\
 &\leq h_1 \min\{\phi(M(x_n, z, t)), \phi(M(f(x_n), x_n, t)), \phi(M(f(x_n), z, t))\} \\
 &\quad + h_2 \min\{\phi(M(x_n, z, t)), \phi(M(f(z), z, t)), \phi(M(x_n, f(z), t))\} \\
 &\Rightarrow \lim_{n \rightarrow \infty} \phi(M(x_{n+1}, f(z), t)) = 0 \\
 &\Rightarrow \phi(M(z, f(z), t)) = 0 \\
 &\Rightarrow f(z) = z, \text{ establishes that } z \text{ is a fixed point for } f.
 \end{aligned}$$

Uniqueness can be shown easily.

Hence, z is the unique fixed point of f . \square

Prudhvi [12] proved a common fixed point theorem for asymptotically regular self mappings in cone metric spaces. We obtain an extension of this result for fuzzy metric spaces.

THEOREM 2.6. *Let (X, M, T) be a fuzzy metric space, ϕ be the altering distance function and f and g be two commutative self mappings on X such that*

$$\begin{aligned}
 \phi(M(f(x), f(y), t)) &\leq k_1[\phi(M(g(x), g(y), t)) + k_2(\phi(M(g(x), f(x), t)) \\
 (9) \quad &\quad + \phi(M(g(y), f(y), t)))]
 \end{aligned}$$

where $x, y \in X$, $t > 0$ and $k_1 : \mathbb{R} \rightarrow [0, 1]$, $0 < k_1, k_2 < 1$.

Moreover if

- (i) f and g are asymptotically regular at x_0 ,
- (ii) $f(X) \subseteq g(X)$,
- (iii) $f(X)$ or $g(X)$ is a complete subspace of X ,

then f and g have a unique common fixed point.

Proof. Let $x_0 \in X$. Since $f(X) \subseteq g(X)$, define a sequence $\{u_n\}$ by $u_{n+1} = f(x_n) = g(x_{n+1})$, $n \in \mathbb{N} \cup \{0\}$. Again since f and g are asymptotically regular at x_0 ,

$$(10) \quad \lim_{n \rightarrow \infty} \phi(M(u_n, u_{n+1}, t)) = 0$$

To show that the sequence $\{u_n\}$ is Cauchy.

Suppose, there exists $0 < \varepsilon < 1$ and two sequences of integers $\{r_n\}$ and

$\{s_n\}$ such that $r_n > s_n > n$,

$$(11) \quad \begin{aligned} M(u_{r_n}, u_{s_n}, t) &\leq 1 - \varepsilon, \\ M(u_{r_{n-1}}, u_{s_{n-1}}, t) &> 1 - \varepsilon, \\ M(u_{r_{n-1}}, u_{s_n}, t) &> 1 - \varepsilon, \quad \forall n \in \mathbb{N} \cup \{0\}. \end{aligned}$$

Following the technique applied in Theorem 2.2 we can show that

$$(12) \quad \lim_{n \rightarrow \infty} M(u_{r_n}, u_{s_n}, t) = 1 - \varepsilon, \quad t > 0$$

Now from (9)

$$\begin{aligned} \phi(M(u_{r_{n+1}}, u_{s_{n+1}}, t)) &= \phi(M(f(x_{r_n}), f(x_{s_n}), t)) \\ &\leq k_1[\phi(M(g(x_{r_n}), g(x_{s_n}), t)) + k_2(\phi(M(g(x_{r_n}), f(x_{r_n}), t)) \\ &\quad + \phi(M(g(x_{s_n}), f(x_{s_n}), t)))] \end{aligned}$$

Taking $n \rightarrow \infty$ and using (10) and (12) we have

$$\phi(1 - \varepsilon) \leq k_1 \phi(1 - \varepsilon) < \phi(1 - \varepsilon)$$

a contradiction. Hence $\{u_n\}$ is Cauchy sequence.

Suppose that $g(X)$ is complete, then there exists $v \in g(X)$ such that $\lim_{n \rightarrow \infty} u_n = v$. Also, for some $z \in X$ we have $g(z) = v$.

Now,

$$\begin{aligned} \phi(M(f(z), u_{n+1}, t)) &= \phi(M(f(z), f(x_n), t)) \\ &\leq k_1[\phi(M(g(z), g(x_n), t)) + k_2(\phi(M(f(z), g(z), t)) \\ &\quad + \phi(M(f(x_n), g(x_n), t)))] \end{aligned}$$

For $n \rightarrow \infty$,

$$\begin{aligned} \phi(M(f(z), v, t)) &\leq k_1[k_2 \phi(M(f(z), v, t))] \\ \Rightarrow (1 - k_1 \cdot k_2) \phi(M(f(z), v, t)) &= 0 \\ \Rightarrow \phi(M(f(z), v, t)) &= 0 \\ \Rightarrow f(z) &= v \end{aligned}$$

Therefore $f(z) = v = g(z)$ i.e., z is the coincident point of f and g .
Next, from (9),

$$\begin{aligned} \phi(M(f(f(z)), f(z), t)) &\leq k_1[\phi(M(g(f(z)), g(z), t)) + k_2(\phi(M(f(f(z)), \\ &\quad g(f(z)), t)) + \phi(M(f(z), g(z), t)))] \\ &= k_1[\phi(M(f(g(z)), g(z), t)) + k_2(\phi(M(f(f(z)), f(g(z)), t)) \\ &\quad + \phi(M(f(z), g(z), t)))] \quad (\text{Since } fg = gf) \\ &= k_1[\phi(M(f(f(z)), f(z), t)) + k_2(\phi(M(f(f(z)), f(f(z)), t)) \\ &\quad + \phi(M(f(z), f(z), t)))] \\ &= k_1\phi(M(f(f(z)), f(z), t)) \\ \Rightarrow (1 - k_1)\phi(M(f(f(z)), f(z), t)) &= 0 \\ \Rightarrow \phi(M(f(f(z)), f(z), t)) &= 0 \\ \Rightarrow f(f(z)) = f(z) = v \end{aligned}$$

Similarly, $g(g(z)) = g(z) = v$.

Hence v is a common fixed point of f and g . If v_1 is another common fixed point of f and g , then

$$\begin{aligned} \phi(M(f(v), f(v_1), t)) &\leq k_1[\phi(M(g(v), g(v_1), t)) + k_2(\phi(M(f(v), g(v), t)) \\ &\quad + \phi(M(f(v_1), g(v_1), t)))] \\ \Rightarrow \phi(M(v, v_1, t)) &\leq k_1[\phi(M(v, v_1, t)) + k_2(\phi(M(v, v, t)) + \phi(M(v_1, v_1, t)))] \\ \Rightarrow (1 - k_1)\phi(M(v, v_1, t)) &= 0 \\ \Rightarrow v = v_1 \end{aligned}$$

Hence f and g have a unique common fixed point in X . \square

EXAMPLE 2.7. For $X = \{1, 2, 3, 4, 5\}$, let M be a fuzzy set on $X \times X \times [0, \infty[$ given by $M(x, y, t) = \frac{t}{t + d(x, y)}$, $t > 0$, where $d(x, y) = |x - y| \forall x, y \in X$.

Then (X, M, T) is a complete fuzzy metric space with respect to the t -norm $T(a, b) = \min\{a, b\}$, $a, b \in [0, 1]$. Define $f, g : X \rightarrow X$ by $f(1) = f(2) = 2, f(3) = f(4) = f(5) = 1$ and $g(1) = g(2) = 2, g(3) = g(4) = 3, g(5) = 1$.

Also let $\phi(\lambda) = 1 - \lambda$. Then f and g satisfy (9), and hence from Theorem 2.6, we have f and g have a unique common fixed point.

In view of Theorem 2.6, we formulate the following result without proof.

THEOREM 2.8. *Let (X, M, T) be a fuzzy metric space, ϕ be the altering distance function and f and g be two self mappings on X such that*

$$(13) \quad \begin{aligned} \phi(M(f(x), g(y), t)) \leq & k_1[\phi(M(f(x), g(x), t)) + \phi(M(f(y), g(y), t))] + \\ & k_2[\phi(M(f(x), f(y), t)) + \phi(M(g(x), g(y), t))] \\ & + k_3\phi(M(f(y), g(x), t)) \end{aligned}$$

where $x, y \in X$, $t > 0$ and $k_1 + 2k_2 + k_3 < 1$, $0 < k_1, k_2, k_3 < 1$. Moreover (f, g) is asymptotically regular at x_0 and commutes with each other. If the range of f is a subset of range of g and if $f(X)$ or $g(X)$ is a complete subspace of X then f and g have a unique common fixed point.

Now we derive a common fixed point theorem for three compatible self mappings.

THEOREM 2.9. *Let f, g and h be self mappings on the complete fuzzy metric space (X, M, T) and ϕ be altering distance function such that the following conditions are satisfied:*

- (I) $\phi(M(f(x), f(y), t))$
 $\leq k_1\phi(M(f(x), g(y), t))[\phi(M(g(x), h(x), t)) + \phi(M(g(y), h(y), t))]$
 $+ k_2\phi(M(g(x), h(y), t))[\phi(M(h(x), f(x), t)) + \phi(M(h(y), f(y), t))]$
 $+ k_3\phi(M(h(x), f(y), t))[\phi(M(f(x), g(x), t)) + \phi(M(f(y), g(y), t))]$
 for all $x, y \in X$ and $0 < k_1, k_2, k_3 < 1$,
- (II) g and h are continuous mappings, and each pair (f, g) , (f, h) and (g, h) are compatible,
- (III) there exists a sequence $\{x_n\}$ in X which is asymptotically regular with respect to each pair (f, g) , (f, h) and (g, h) .

Then f, g and h have a common unique fixed point in X .

Proof. From condition (I), for $n, m \in \mathbb{N}, t > 0$

$$\begin{aligned} \phi(M(f(x_n), f(x_m), t)) &\leq k_1 \phi(M(f(x_n), g(x_m), t)) [\phi(M(g(x_n), h(x_n), t)) \\ &\quad + \phi(M(g(x_m), h(x_m), t))] + k_2 \phi(M(g(x_n), h(x_m), t)) \\ &\quad [\phi(M(h(x_n), f(x_n), t)) + \phi(M(h(x_m), f(x_m), t))] \\ &\quad + k_3 \phi(M(h(x_n), f(x_m), t)) [\phi(M(f(x_n), g(x_n), t)) \\ &\quad + \phi(M(f(x_m), g(x_m), t))], \end{aligned}$$

which shows that

$$\begin{aligned} \lim_{n, m \rightarrow \infty} \phi(M(f(x_n), f(x_m), t)) &= 0, \\ \text{i.e., } \lim_{n, m \rightarrow \infty} M(f(x_n), f(x_m), t) &= 1 \end{aligned}$$

So, the sequence $\{f(x_n)\}$ is a Cauchy sequence.

Since X is complete, $\{f(x_n)\}$ converges to some $z \in X$ (say).

Now,

$$M(g(x_n), z, t) \leq T(M(g(x_n), f(x_n), t), M(f(x_n), z, t))$$

Since $\{x_n\}$ is asymptotically regular with respect to the pair (f, g)

and $f(x_n)$ converges to z ,

$$\lim_{n \rightarrow \infty} M(g(x_n), z, t) \leq T(1, 1) = 1,$$

$$\text{i.e., } \lim_{n \rightarrow \infty} M(g(x_n), z, t) = 1,$$

shows that $\{g(x_n)\}$ converges to z .

Similarly, we can show that $\{h(x_n)\}$ converges to z .

Again,

$$M(f(g(x_n)), g(z), t) \geq T(M(f(g(x_n)), g(f(x_n)), t), M(g(f(x_n)), g(z), t))$$

Since f and g are compatible and g is continuous,

$$\begin{aligned} \lim_{n \rightarrow \infty} M(f(g(x_n)), g(z), t) &\geq T(1, 1) \\ \Rightarrow \lim_{n \rightarrow \infty} M(f(g(x_n)), g(z), t) &= 1 \\ \text{i.e., } f(g(x_n)) &\longrightarrow g(z). \end{aligned}$$

Similarly, $f(h(x_n)) \rightarrow h(z)$.

Now from condition (I), using the continuity of g and h ,

$$\begin{aligned} \phi(M(f(g(x_n)), f(h(x_n)), t)) &\leq k_1\phi(M(f(g(x_n)), g(h(x_n)), t)) \\ &\quad [\phi(M(g(g(x_n)), h(g(x_n)), t)) + \phi(M(g(h(x_n)), h(h(x_n)), t))] \\ &\quad + k_2\phi(M(g(g(x_n)), h(h(x_n)), t))[\phi(M(h(g(x_n)), f(g(x_n)), t)) \\ &\quad + \phi(M(h(h(x_n)), f(h(x_n)), t))] + k_3\phi(M(h(g(x_n)), f(h(x_n)), t)) \\ &\quad [\phi(M(f(g(x_n)), g(g(x_n)), t)) + \phi(M(f(h(x_n)), g(h(x_n)), t))] \end{aligned}$$

Taking limit as $n \rightarrow \infty$,

$$\phi(M(g(z), h(z), t)) \leq k_2\phi(M(g(z), h(z), t)) \cdot \phi(M(g(z), h(z), t))$$

If $\phi(M(g(z), h(z), t)) \neq 0$, then

$$1 \leq k_2\phi(M(g(z), h(z), t))$$

which is a contradiction as $k_2 < 1$. Therefore

$$\begin{aligned} \phi(M(g(z), h(z), t)) &= 0 \\ \Rightarrow M(g(z), h(z), t) &= 1 \\ \Rightarrow g(z) &= h(z) \end{aligned}$$

Again,

$$\begin{aligned} (14) \quad \phi(M(f(g(x_n)), f(z), t)) &\leq k_1\phi(M(f(g(x_n)), g(z), t))[\phi(M(g(g(x_n)), h(g(x_n)), t)) \\ &\quad + \phi(M(g(z), h(z), t))] + k_2\phi(M(g(g(x_n)), h(z), t)) \\ &\quad [\phi(M(h(g(x_n)), f(g(x_n)), t)) + \phi(M(h(z), f(z), t))] + \\ &\quad k_3\phi(M(h(g(x_n)), f(z), t))[\phi(M(f(g(x_n)), g(g(x_n)), t)) \\ &\quad + \phi(M(f(z), g(z), t))] \end{aligned}$$

Taking limit as $n \rightarrow \infty$

$$\begin{aligned} &\Rightarrow \phi(M(g(z), f(z), t)) \\ (15) \quad &\leq k_3\phi(M(h(z), f(z), t)) \cdot \phi(M(f(z), g(z), t)) \quad (\text{since } g(z) = h(z)) \end{aligned}$$

Now, $M(h(z), f(z), t) \leq T(M(h(z), g(z), t), M(g(z), f(z), t))$

$$(16) \quad \Rightarrow M(h(z), f(z), t) \leq T(1, M(g(z), f(z), t)) = M(g(z), f(z), t)$$

Hence from (14) and (16),

$$\begin{aligned}\phi(M(g(z), f(z), t)) &\leq k_3 \phi(M(g(z), f(z), t)) \cdot \phi(M(g(z), f(z), t)) \\ \Rightarrow \phi(M(g(z), f(z), t)) &= 0 \\ \Rightarrow g(z) &= f(z)\end{aligned}$$

Similarly,

$$\begin{aligned}\phi(M(f(z), f(x_n), t)) &\leq k_1 \phi(M(f(z), g(x_n), t)) [\phi(M(g(z), h(z), t)) \\ &\quad + \phi(M(g(x_n), h(x_n), t))] + k_2 \phi(M(g(z), h(x_n), t)) \\ &\quad [\phi(M(h(z), f(z), t)) + \phi(M(h(x_n), f(x_n), t))] + \\ &\quad k_3 \phi(M(h(z), f(x_n), t)) [\phi(M(f(z), g(z), t)) + \\ &\quad \phi(M(f(x_n), g(x_n), t))]\end{aligned}$$

Taking limit as $n \rightarrow \infty$ and using asymptotic regularity of x_n with respect to each pair of mappings (f, g) , (f, h) and (g, h) ,

$$\begin{aligned}\phi(M(f(z), z, t)) &= 0 \\ \Rightarrow f(z) &= z\end{aligned}$$

Hence $f(z) = g(z) = h(z) = z$ i.e., z is a common fixed point of f , g and h .

For the uniqueness, suppose there is another common fixed point of f , g and h , say v . Then from condition (I)

$$\begin{aligned}\phi(M(f(z), f(v), t)) &\leq k_1 \phi(M(f(z), g(v), t)) [\phi(M(g(z), h(z), t)) \\ &\quad + \phi(M(g(v), h(v), t))] + k_2 \phi(M(g(z), h(v), t)) \\ &\quad [\phi(M(h(z), f(z), t)) + \phi(M(h(v), f(v), t))] + \\ &\quad k_3 \phi(M(h(z), f(x_n), t)) [\phi(M(f(z), g(z), t)) + \\ &\quad \phi(M(f(v), g(v), t))]\end{aligned}$$

$$\begin{aligned}\Rightarrow \phi(M(z, v, t)) &= 0 \\ \Rightarrow M(z, v, t) &= 1 \\ \Rightarrow z &= v\end{aligned}$$

Therefore z is a unique common fixed point of f , g and h . \square

EXAMPLE 2.10. Let $X = [0, 2]$, $d(x, y) = |x - y|$ and $M(x, y, t) = 1 - \frac{d(x, y)}{g(t)}$ where $g : (0, \infty) \rightarrow]2, \infty[$. Then (X, M, T) is a complete fuzzy metric space with respect to the t-norm $T(a, b) = \min\{a, b\}$, $a, b \in [0, 1]$.

Consider the self mappings on f, g and h on X defined by $f(x) = \begin{cases} 2, & \text{if } x = 0, \\ 1, & \text{if } x \neq 0 \end{cases}$, $g(x) = x$, $\forall x \in X$ and $h(x) = \begin{cases} 2 - x, & \text{if } x \in [0, 1], \\ 1, & \text{if } x \in (1, 2] \end{cases}$.

Then each pair (f, g) , (f, h) and (g, h) are compatible. Let the sequence $\{x_n\} \in X$ defined by $x_n = 1 - \frac{1}{3^n}$. Then $\{x_n\}$ is asymptotically regular with respect to (f, g) , (f, h) and (g, h) .

Moreover for $k_1 = k_2 = k_3 = \frac{3}{4}$, the condition (I) of Theorem 2.9 is satisfied.

Hence from Theorem 2.9, f, g and h have a common unique fixed point in X which is 1 in this case.

3. Conclusion

Throughout this paper, we have discussed some fixed point results for the class of asymptotically regular mappings in fuzzy metric spaces. The practical applications of these results for solving systems of fuzzy differential as well as integral equations is a scope for future study.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

References

- [1] F. E Browder and W. V. Petryshyn, *The solution by iteration of nonlinear functional equations in Banach Spaces*, Bull. Amer. Math. Soc. **72** (1966), 571–575.
- [2] A. George and P. Veeramani, *On some results in fuzzy metric spaces*, Fuzzy Set. Syst. **64** (1994), 395–399.
- [3] M. Grabiec, *Fixed points in fuzzy metric spaces*, Fuzzy Set. Syst. **27** (1988), 385–389.
- [4] A. A. Harandi and H. Emami, *A fixed point theorem for contraction type maps in partially ordered metric spaces and application to ordinary different equations*, Nonlinear Analysis **72** (2010), 2238–2242.
- [5] M. S. Khan, M. Swaleh and S. Sessa, *Fixed point theorems by altering distances between the points*, Bull. Aust. Math. Soc. **30** (1984), 1–9.

- [6] B. E. Rhoades, S. Seesa, M. S. Khan and M. D. Khan, *Some fixed point theorems for hardy-rogers type mappings*, International J. Math. and Math. Sci. **7** (1984), 75–87.
- [7] J. Rodriguez-Lopez and S. Romaguera, *The Hausdorff fuzzy metric on compact sets*, Fuzzy Sets Syst. **147** (2004), 273–283.
- [8] H. K. Nashine and Z. Kadelburg, *Common fixed point theorems for asymptotically regular mappings on ordered orbitally complete metric spaces with an application to systems of integral equations*, Filomat **30** (12) (2016), 3277–3289.
- [9] P. Nigam and S. S. Pagey, *Some fixed point theorems for a pair of asymptotically regular and compatible mappings in fuzzy 2-metric space*, Int. J. Open Problems Comput. Math. **5** (1) (2012), 71–84.
- [10] B. Patir, N. Goswami and L. N. Mishra, *Fixed point theorems in fuzzy metric spaces for mappings with some contractive type conditions*, Korean J. Math. **26** (2) (2018), 307–326.
- [11] B. Patir, N. Goswami and V. N. Mishra, *Some results on fixed point theory for a class of generalized nonexpansive mappings*, Fixed Point Theory and Applications **2018** (19) (2018).
- [12] K. Prudhvi, *A common fixed point theorem for asymptotically regular in cone metric spaces*, Asian Journal of Fuzzy and Applied Mathematics **2**(1), 12-16 (2014).
- [13] K. P. R. Sastry, V. S. R. Naidu, I. H. N. Rao and K. P. R. Rao, *Common fixed points for asymptotically regular mappings*, Indian J. Pure Appl. Math. **15** (8) (1984), 849–854.
- [14] Y. Shen, D. Qiu and W. Chen, *Fixed point theorems in fuzzy metric spaces*, Appl. Math. Lett. **25** (2012), 138–141.
- [15] S. Shukla, I. Altun and R. Sen, *Fixed point theorems and asymptotically regular mappings in partial metric spaces*, ISRN Computational Mathematics **2013**, 6 pages (2013).
- [16] B. C. Tripathy, S. Paul and N. R. Das, *Banach's and Kannan's fixed point results in fuzzy 2-metric spaces*, Proyecciones J. Math. **32**(4) (2013), 359–375.
- [17] B. C. Tripathy, S. Paul and N. R. Das, *A fixed point theorem in a generalized fuzzy metric space*, Bol. Soc. Paran. Mat. **32**(2) (2014), 221–227.
- [18] B. C. Tripathy, S. Paul and N. R. Das, *Fixed point and periodic point theorems in fuzzy metric space*, Songklanakarin Journal of Science and Technology, **37** (1) (2015), 89–92.

Nilakshi Goswami

Department of Mathematics

Gauhati University

Guwahati-781014, Assam, India

E-mail: nila_g2003@yahoo.co.in

Bijoy Patir

Department of Mathematics

Gauhati University

Guwahati-781014, Assam, India

E-mail: bpatir07@gmail.com