Kangweon-Kyungki Math. Jour. **10** (2002), No. 1, pp. 19–23

# FIXED POINTS OF SUMS OF NONEXPANSIVE MAPS AND COMPACT MAPS

Jongsook Bae and Daejong An

Abstract. Let $X$ be a Banach space satisfying Opial's condition, $C$ a weakly compact convex subset of $X$, $F : C \to X$ a nonexpansive map, and let $G : C \to X$ be a compact and demiclosed map. We prove that $F + G$ has a fixed point in $C$ if $F + G : C \to X$ is a weakly inward map.

## 1. Introduction

Let $C$ be a nonempty subset of a Banach space $X$. A map $T : C \to X$ is called nonexpansive if $\|T(x) - T(y)\| \leq \|x - y\|$ for all $x, y \in C$. A map $T : C \to X$ is said to be a Lipschitzian if there exists $k \geq 0$ such that for all $x, y \in C$, $\|T(x) - T(y)\| \leq k\|x - y\|$; it is a contraction if $0 < k < 1$ and is a compact if it is continuous and maps bounded sets to relatively compact sets.

In 1955, Krasnoselskii proved the following theorem which can be found in [1].

Theorem 1. *Let $C$ be a nonempty bounded closed convex set in a Banach space $X$. Let $F : C \to X$ be a contraction map, and let $G : C \to X$ be a compact map. If $(F + G)(C) \subseteq C$, then $F + G$ has a fixed point in $C$.*

Let $X$ be a Banach space and $\mathcal{B}$ the family of its bounded subsets. Then $\alpha : \mathcal{B} \to [0, \infty]$, defined by

$$\alpha(B) = \inf\{d > 0 | B \text{ admits a finite cover by sets of diameter} \leq d \},$$

is called the Kuratowski measure of noncompactness ([2]). It is not only natural but also useful since $\alpha$ has interesting properties, some of which are listed in Proposition 2 ([4]).

Received December 22, 2001.

2000 Mathematics Subject Classification: 47H10.

Key words and phrases: nonexpansive map, compact map, Krasnoselskii fixed point theorem, Opial's condition, weakly inward map.

PROPOSITION 2. *Let $X$ be a Banach space, $\mathcal{B}$ the family of all bounded sets of $X$, and let $\alpha : \mathcal{B} \to [0, \infty)$ be the Kuratowski measure of noncompactness. Then*

*(a)* $\alpha(B) = 0$ *iff* $\bar{B}$ *is compact.*

*(b)* $\alpha$ *is a seminorm, i.e.,*

$$\alpha(\lambda B) = |\lambda|\alpha(B) \text{ and}$$
$$\alpha(B_1 + B_2) \leq \alpha(B_1) + \alpha(B_2).$$

*(c)* $B_1 \subseteq B_2$ *implies* $\alpha(B_1) \leq \alpha(B_2)$;
$\alpha(B_1 \cup B_2) = \max\{\alpha(B_1), \alpha(B_2)\}$.

*(d)* $\alpha(convB) = \alpha(B)$.

*(e)* $\alpha$ *is continuous with respect to the Hausdorff metric* $H$, *defined by* $H(B_1, B_2) = \max\{\sup_{B_1} d(x, B_2), \sup_{B_2} d(x, B_1)\}$; *in particular* $\alpha(\bar{B}) = \alpha(B)$.

For $D \subset X$, $T : D \to X$ is called condensing if $\alpha(T(B)) < \alpha(B)$ for any bounded subset $B$ of $D$ with $\alpha(B) > 0$. Let $K$ be a convex subset of a Banach space $X$ and $x \in K$. The inward set $I_K(x)$ of $K$ at $x$ is defined by

$$I_K(x) = \{r(y - x) + x | y \in K, r \geq 1\}.$$

A map $T : K \to X$ is called inward if for all $x \in K$, $T(x) \in I_K(x)$, and $T$ is said to be weakly inward if for all $x \in K$, $T(x) \in \overline{I_K(x)}$.

In 1979, Deimling proved the following theorem which can be found in [3].

THEOREM 3. *Let $X$ be a Banach space, $D \subset X$ closed bounded convex, $F : D \to X$ a continuous, condensing and weakly inward map. Then $F$ has a fixed point.*

REMARK. Let $C$ be a nonempty bounded closed convex set in a Banach space $X$. Let $F : C \to X$ be a contraction map and $G : C \to X$ a compact map. Then $T = F + G$ is a continuous and condensing map. Indeed, if $B$ is a subset of $C$ and $\alpha(B) > 0$, then we have

$$\begin{aligned}
\alpha(T(B)) &= \alpha(F(B) + G(B)) \\
&\leq \alpha(F(B)) + \alpha(G(B)) \\
&\leq \alpha(F(B)) + \alpha(\overline{G(B)}) \\
&= \alpha(F(B)) \\
&< \alpha(B).
\end{aligned}$$

From Theorem 3 and Remark, Theorem 1 can be restated as follows; Let $C$ be a nonempty bounded closed convex set in a Banach space $X$. Let $F : C \to X$ be a contraction map, and let $G : C \to X$ be a compact map. If $T = F + G : C \to X$ is a weakly inward map, then $T$ has a fixed point.

The purpose of this paper generalizes the above result by replacing nonexpansive maps instead of contraction maps.

## 2. The results

A Banach space $X$ is said to satisfy Opial's condition if whenever a sequence $\{x_n\}$ in $X$ converges weakly to $x_0$, then for $x \neq x_0$,

$$\lim_{n\to\infty} \inf \|x_n - x_0\| < \lim_{n\to\infty} \inf \|x_n - x\| \quad ([5]).$$

Let $D$ be a subset of a Banach space $X$. A map $T : D \to X$ is said to be demiclosed if for any sequence $\{x_n\}$ in $D$ the following implication holds:

$$w - \lim_{n\to\infty} x_n = x \text{ and } \lim_{n\to\infty} \|T(x_n) - w\| = 0$$

implies

$$x \in D \text{ and } T(x) = w.$$

THEOREM 4. *Let $X$ be a Banach space which satisfies Opial's condition, and $C$ a nonempty weakly compact convex subset of $X$. Let $F : C \to X$ be a nonexpansive map and $G : C \to X$ a compact and demiclosed map. If $T = F + G : C \to X$ is a weakly inward map, then $T$ has a fixed point.*

*Proof.* Without loss of generality we may assume $0 \in C$. Let $0 < k < 1$. Then by the same way as in Remark we can show that, $kT = kF + kG$ is continuous, $k$-condensing. Also it can be easily shown that $kT : C \to X$ is a weakly inward map (See [3]). From Theorem 3, $kT$ has a fixed point, i.e., there exists $x_k$ in $C$ such that $kTx_k = x_k$. In this case we have

$$\|x_k - Tx_k\| = \|x_k - \frac{x_k}{k}\| = \frac{1-k}{k}\|x_k\| \to 0 \ as \ k \to 1^-.$$

Since $C$ is weakly compact and $G$ is a compact map we can take a sequence $\{x_n\}$ in $C$ such that $\{x_n\}$ converges weakly to $x$ for some $x \in C$, $\|Tx_n - x_n\| \to 0$ and $G(x_n) \to y$ for some $y \in X$. Then we have

$$\begin{aligned}\|Tx_n - Fx - y\| &\leq \|Fx_n - Fx\| + \|Tx_n - Fx_n - y\| \\ &\leq \|x_n - x\| + \|Tx_n - Fx_n - y\|.\end{aligned}$$

Hence we have

$$\begin{aligned}&\liminf \|x_n - (Fx + y)\| \\ &\leq \liminf(\|x_n - Tx_n\| + \|Tx_n - (Fx + y)\|) \\ &\leq \liminf(\|x_n - Tx_n\| + \|x_n - x\| + \|Tx_n - (Fx_n + y)\|) \\ &= \liminf \|x_n - x\|.\end{aligned}$$

Since $X$ satisfies Opial's condition, we have $Fx + y = x$. And since $G$ is demiclosed $Gx = y$ so that $Tx = Fx + Gx = x$. Hence $T$ has a fixed point. □

Corollary 5. *Let $X$ be a reflexive Banach space which satisfies Opial's condition, and $C$ a bounded closed convex subset of $X$. Let $F : C \to X$ be a nonexpansive map and $G : C \to X$ a compact and demiclosed map. If $T = F + G : C \to X$ is a weakly inward map, then $T$ has a fixed point.*

Applications of our results will be given sufficient conditions so that there exist solutions of difference equations which are asypmtotically constant ([6]).

## References

1. M. Krasnoselskii, *Two remarks on the method of successive approximations*, Uspekhi Math. Nauk **1** (1955), 123-127.
2. K. Kuratowski, *Sur les espaces complets*, Fund. Math **15** (1930), 301-309.
3. K. Deimling, *Fixed points of condensing maps, Lecture note in Math*, Springer-Verlag, Berlin **737** (1979), 67-82.
4. K. Deimling, *Nonlinear functional analysis*, Springer-Verlag, Berlin (1985).
5. K. Goebel and W. A. Kirk, *Topics in metric fixed point theory*, Cambridge University press, (1990).

6. G. Gonzalez and A. Jimeuez-Melado, *An application of Krasnoselskii fixed point theorem to the Asymptotic behavior of solutions of difference equations in Banach spaces*, J. Math. Anal. Appl **247** (2000), 290-299.

Department of Mathematics
College of Science
Myong Ji University
Yongin-Si, 449-728, Korea